IAPT CHROMOSOME DATA

IAPT chromosome data 28

Edited by Karol Marhold & Jaromír Kučera

Julio Rubén Daviña & Ana Isabel Honfi*

Programa de Estudios Florísticos y Genética Vegetal, Instituto de Biología Subtropical CONICET-Universidad Nacional de Misiones, nodo Posadas, Rivadavia 2370, 3300 Posadas, Argentina * Author for correspondence: ahonfi@gmail.com

This study was supported by Agencia Nacional de Promoción Científica y Técnica (ANPCyT) grant nos. PICT-2014-2218 and PICT-2016-1637, and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

All materials CHN; collectors: D = J.R. Daviña, H = A.I. Honfi, L = B. Leuenberger.

AMARYLLIDACEAE

Habranthus barrosianus Hunz. & Di Fulvio, 2n = 2x = 18; Argentina, Buenos Aires, *L* 3772 (B).

- *Habranthus brachyandrus* (Baker) Sealy, n = 12II, 2n = 4x = 24; Argentina, Corrientes, *D* 416 (B, MNES).
- *Habranthus chacoensis* Ravenna, n = 6II, 2n = 2x = 12; Argentina, Chaco, *D* 342 (B, CTES, MNES).
- *Habranthus coeruleus* (Griseb.) Traub, 2n = 2x = 30; Argentina, Misiones, *H* 985 (MNES).
- *Habranthus gracilifolius* Herb., 2n = 4x = 24; Argentina, Buenos Aires, *L* 3774 (B).
- *Habranthus pedunculosus* Herb., 2n = 2x = 14; Argentina, Chaco, *D* 334 (B, MNES).
- *Habranthus tubisphathus* (L'Hér.) Traub, 2n = 4x = 24; Argentina, Corrientes, *D 329* (B, CTES, MNES). 2n = 6x = 36; Argentina, Buenos Aires, *L 3771* (B).

Luis Delgado,^{1*} Blanca M. Rojas-Andrés,^{1,2,3} Noemí López-González,^{1,2} Nélida Padilla-García^{1,2} & M. Montserrat Martínez-Ortega^{1,2}

- 1 Departamento de Botánica y Fisiología Vegetal, University of Salamanca, 37007 Salamanca, Spain
- 2 Biobanco de ADN Vegetal, University of Salamanca, Edificio Multiusos I+D+i, Calle Espejo s/n, 37007 Salamanca, Spain
- 3 Present address: Department of Molecular Evolution and Plant Systematics & Herbarium (LZ), Institute of Biology, Leipzig University, Leipzig, Germany
- * Author for correspondence: *ldelsan@usal.es*

DOI https://doi.org/10.12705/676.39

Chromosome numbers counted by L. Delgado and ploidy level estimated by B. Rojas-Andrés and N. López-González; collectors: AA = Antonio Abad, AT = Andreas Tribsch, BR = Blanca Rojas-Andrés, DGL = David Gutiérrez Larruscain, DP = Daniel Pinto, JASA= José Ángel Sánchez Agudo, JPG = Julio Peñas de Giles, LMC = Luz M^a Muñoz Centeno, MO = M. Montserrat Martínez-Ortega, MS= María Santos Vicente, NLG = Noemí López-González, NPG = Nélida Padilla-García, SA = Santiago Andrés, SB = Sara Barrios, VL= Víctor Lucía, XG = Ximena Giráldez.

This work has been supported by the Spanish Ministerio de Economía y Competitividad (projects CGL2009-07555, CGL2012-32574, *Flora iberica* VIII [CGL2008-02982-C03-02/CLI], *Flora iberica* IX [CGL2011-28613-C03-03], *Flora iberica* X [CGL2014-52787-C3-2-P]); the Spanish Ministerio de Ciencia e Innovación (Ph.D. grants to BR and NLG), and the University of Salamanca (Ph.D. grant to NPG financed jointly by Banco Santander). The authors are very grateful to all colleagues who helped with plant location and sampling.

PLANTAGINACEAE

- *Veronica angustifolia* (Vahl) Bernh., 2*n* = 64, CHN. Switzerland, *XG*, *MO* 6038 & *BR* (SALA 149416). 2*n* ~ 8*x* ~ 64, FCM. France, *XG*, *MO* & *BR* 167-1, *XG*, *MO* & *BR* 167-2, *XG*, *MO* & *BR* 167-3 (SALA 149398); Switzerland, *XG*, *MO* 6038-1 & *BR*, *XG*, *MO* 6038-2 & *BR*, *XG*, *MO* 6038-3 & *BR* (SALA 149416).
- *Veronica austriaca* subsp. *jacquinii* (Baumg.) Watzl, 2n = 32, CHN. Bosnia and Herzegovina, *MO 6122, XG, NPG & NLG* (SALA 157023); *MO 6123, XG, NPG & NLG* (SALA 157024). 2n = 48, CHN. Bosnia and Herzegovina, *MO 6120, XG, NPG & NLG* (SALA 157021); Bulgaria, *XG, MO, BR 21 & MS* (SALA 149368), *XG, MO 4577, BR & MS* (SALA 149376). $2n \sim 6x \sim 48$, FCM. Croatia, *MO 1404-1 & AT, MO 1404-2 & AT* (SALA 124609), *MO 1391-1 & AT, MO 1391-2 & AT* (SALA 124604).
- Veronica dalmatica N.Pad.Gar., Rojas-Andrés, López-González & M.M.Mart.Ort., 2n = 16, CHN. Montenegro, MO, XG, NPG & NLG 137 (SALA 157018), MO, XG, NPG & NLG 139 (SALA 157030).
- *Veronica kindlii* Adam., 2*n* = 16, CHN. Macedonia (F.Y.R.O.M.), *XG*, *MO* 6090, *BR* & *NLG* (SALA 157011).
- Veronica orsiniana Ten., 2n ~ 2x ~ 16, FCM. France, XG, MO & BR 203-1, XG, MO & BR 203-2, XG, MO & BR 203-3 (SALA 149303), XG, MO & BR 207-1, XG, MO & BR 207-2, XG, MO & BR 207-3 (SALA 149305), XG, MO & BR 210-1, XG, MO & BR 210-2, XG, MO & BR 210-3 (SALA 149308); Spain, MO, BR 235-1, AA & NLG, MO, BR 235-2, AA & NLG, MO, BR 235-3, AA & NLG (SALA 155093), MO,

All materials for the chromosome column should be submitted electronically to: Karol Marhold, karol.marhold@savba.sk (Institute of Botany, Slovak Academy of Sciences, SK-845 23 Bratislava, Slovakia, and Department of Botany, Charles University, CZ 128-01 Prague, Czech Republic). The full version of this contribution is available in the online edition of TAXON appended to this article. The following citation format is recommended: Baltisberger, M. & Voelger, M. 2006. *Sternbergia sicula*. In: Marhold, K. (ed.), IAPT/IOPB chromosome data 1. *Taxon* 55: 444, E2.

BR 235-22*, AA & NLG (SALA 155067), MO 6064-1, BR, XG & NLG, MO 6064-2, BR, XG & NLG, MO 6064-3, BR, XG & NLG (SALA 155119), MO, BR, XG & NLG 8-1, MO, BR, XG & NLG 8-2, MO, BR, XG & NLG 8-3 (SALA 155097), MO, BR 234-1, AA & NLG, MO, BR 234-2, AA & NLG, MO, BR 234-3, AA & NLG (SALA 155094), MO 6063-1, BR, AA, XG & NLG, MO 6063-2, BR, AA, XG & NLG, MO 6063-3, BR, AA, XG & NLG (SALA 155123), MO, BR 242-1, XG & NLG, MO, BR 242-2, XG & NLG, MO, BR 242-3, XG & NLG (SALA 155109), MO, BR 243-1, XG & NLG, MO, BR 243-2, XG & NLG, MO, BR 243-3, XG & NLG (SALA 155110).

- Veronica prostrata L., 2n ~ 2x ~ 16, FCM. Austria, MO 1055-1, MO 1055-2, MO 1055-3 (SALA 124616), MO 1337-1, MO 1337-2, MO 1337-3 (SALA 124615); Czech Republic, XG, MO & BR 184-1, XG, MO & BR 184-2, XG, MO & BR 184-3 (SALA 149310), MO 1445-1 (SALA 124617); France, XG, MO & BR 219-1, XG, MO & BR 219-2, XG, MO & BR 219-3 (SALA 149313); Romania, XG, MO 904-1 & JASA, XG, MO 904-2 & JASA (SALA 124620); Serbia, SA 428-1, XG, MO & BR, SA 428-2, XG, MO & BR, SA 428-3, XG, MO & BR, SA 428-4, XG, MO & BR (SALA 149319).
- *Veronica rosea* Desf., 2*n* ~ 2*x* ~ 16, FCM. Morocco, *DP*, *NLG 52-1* & *VL*, *DP*, *NLG 52-2* & *VL*, *DP*, *NLG 52-3* & *VL* (SALA 155072), *JPG JP-1*, *JPG JP-2*, *JPG JP-3* (GDA 59936).
- *Veronica sennenii* (Pau) M.M.Mart.Ort. & E.Rico, 2*n* ~ 8*x* ~ 64, FCM. Spain, *MO*, *BR* 248-1 & *NLG*, *MO*, *BR* 248-2 & *NLG*, *MO*, *BR* 248-3 & *NLG* (SALA 155081).
- Veronica tenuifolia subsp. javalambrensis (Pau) Molero & J.Pujadas, 2n ~ 2x ~ 16, FCM. Spain, MO, BR 250-1 & NLG, MO, BR 250-2 & NLG, MO, BR 250-3 & NLG (SALA 155086), DGL, NLG & DP 1278-1, DGL, NLG & DP 1278-2, DGL, NLG & DP 1278-3 (SALA 150521), DGL, NLG & DP 1280-1, DGL, NLG & DP 1280-2, DGL, NLG & DP 1280-3 (SALA 150519), MO 6074-1, BR & NLG, MO 6074-2, BR & NLG, MO 6074-3, BR & NLG (SALA 155084), MO & NLG 3-1, MO & NLG 3-2, MO & NLG 3-3 (SALA 155084), MO & NLG 3-1, MO & NLG 3-2, MO & NLG 3-3 (SALA 155084), DP 1315-1, DP 1315-2, DP 1315-3 (SALA 150484), DP 1311-1, DP 1311-2, DP 1311-3 (SALA 150488), DGL, NLG & DP 1287-1, DGL, NLG & DP 1287-2, DGL, NLG & DP 1287-3 (SALA 150512), DP 1307-1, DP 1307-2, DP 1307-3 (SALA 150492), DP 1322-1, DP 1322-2, DP 1322-3 (SALA 150477), SB, NLG, MO & BR 221-1, SB, NLG, MO & BR 221-2, SB, NLG, MO & BR 221-3 (SALA 149327).
- Veronica tenuifolia Asso subsp. tenuifolia, 2n ~ 2x ~ 16, FCM. Spain, MO, BR 237-1, AA & NLG, MO, BR 237-2, AA & NLG, MO, BR 237-3, AA & NLG (SALA 155065), MO, BR, AA & NLG 16-1, MO, BR, AA & NLG 16-2, MO, BR, AA & NLG 16-3 (SALA 155125), MO 6059-1, BR, XG & NLG, MO 6059-2, BR, XG & NLG, MO 6059-3, BR, XG & NLG (SALA 155099).
- *Veronica teucrium* L., 2*n* ~ 8*x* ~ 64, FCM. Slovakia, *MO 1551-1, XG* & *LMC*, *MO 1551-2, XG* & *LMC*, *MO 1551-3, XG* & *LMC* (SALA 124598).
- *Veronica thracica* Velen., $2n \sim 2x \sim 16$, FCM. Turkey, *XG*, *MO*, *BR* 37-1 & *MS*, *XG*, *MO*, *BR* 37-2 & *MS* (SALA 149289).

Aleksander A. Gnutikov,^{1,2} Marina V. Protopopova,^{3*} Vasiliy V. Pavlichenko³ & Victor V. Chepinoga^{4,5}

- 1 The N.I. Vavilov Institute of Plant Genetic Resources (VIR), Bolshaya Morskaya Str. 44, 190000 St.-Petersburg, Russia
- 2 Komarov Botanical Institute of the Russian Academy of Sciences, Professora Popova Str. 2, 197376 St.-Petersburg, Russia
- 3 Siberian Institute of Plant Physiology and Biochemistry SB RAS, Lermontov Str. 132, 664033 Irkutsk, Russia
- 4 The V.B. Sochava Institute of Geography SB RAS, Ulan-Batorskaya Str. 1, 664033 Irkutsk, Russia
- 5 Irkutsk State University, Karl Marx Str. 1, 664003 Irkutsk, Russia
- * Author for correspondence: marina.v.protopopova@gmail.com

All materials CHN; collectors: AK = A.D. Konovalov, EZ = E.D. Zolotovskaya, MP = M.V. Protopopova, VP = V.V. Pavlichenko; vouchers in IRKU.

The study was partially supported by the Russian Foundation for Basic Research (projects 16-34-60135-mol_a_dk, 16-05-00783-a, 18-04-01040, 17-00-00338) and done within the framework of the State Task 0126-2014-0028 "Karyological and molecular-phylogenetic investigation of the flowering plant taxa divergence". The chromosome counting for highland populations was proceeded under financial support of the Russian Science Foundation (project 17-74-10074).

AMARYLLIDACEAE

Allium chamarense M.M.Ivanova, 2*n* = 16; Russia, Irkutskaya Oblast', *MP, VP, EZ & AK Cl664.*

APIACEAE

Pleurospermum uralense Hoffm., 2n = 18; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl642.

ASPARAGACEAE

Maianthemum bifolium (L.) F.W.Schmidt, 2n = 36; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1641.

ASTERACEAE

Antennaria dioica (L.) Gaertn., 2n = 28; Russia, Irkutskaya Oblast', MP, VP & EZ Cl663.

Cirsium helenioides (L.) Hill, 2n = 34; Russia, Irkutskaya Oblast', *MP, VP, EZ & AK Cl635.*

Doronicum altaicum Pall., 2n = 60; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl647.

- Jacobaea nemorensis (L.) E.Wiebe, 2n = 40; Russia, Irkutskaya Oblast', *MP*, *VP*, *EZ* & *AK* C1673.
- Parasenecio hastatus (L.) H.Koyama, 2n = 60; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1672.
- Rhaponticum chamarense Peschkova (≡ Rh. carthamoides var. chamarense (Peschkova) O.S.Zhirova), 2n = 26; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1651.
- Saussurea latifolia Ledeb., 2n = 26; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl637.
- Solidago dahurica Kitag., 2n = 18; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1639, MP, VP & EZ C1667.

BORAGINACEAE

Myosotis palustris (L.) L., 2n = 22; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl657.

BRASSICACEAE

Cardamine macrophylla Willd., 2n = 64; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl656.

CARYOPHYLLACEAE

Dianthus superbus L., 2n = 30; Russia, Irkutskaya Oblast', MP, VP & EZ C1665.

CELASTRACEAE

Parnassia palustris L., 2n = 36; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1660.

GERANIACEAE

Geranium krylovii Tzvelev, 2n = 28; Russia, Irkutskaya Oblast', *MP*, *VP*, *EZ* & *AK* C1655.

LAMIACEAE

Prunella vulgaris L., 2n = 28; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1640.

MELANTHIACEAE

Veratrum lobelianum Bernh., 2n = 32; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1653.

ONAGRACEAE

Circaea alpina L., 2*n* = 22; Russia, Irkutskaya Oblast', *MP*, *VP*, *EZ* & *AK* C1670.

OROBANCHACEAE

Pedicularis incarnata L., 2n = 16; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl658.

POACEAE

Poa ×intricata Wein (P. nemoralis L. × P. palustris L.), 2n = 28; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1674.

POLYGONACEAE

Rumex alpestris Jacq., 2n = 14; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1652.

PRIMULACEAE

Primula pallasii Lehm., 2n = 22; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl644.

RANUNCULACEAE

Aconitum baicalense Turcz. ex Rapaics, 2n = 32; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1636.

Aconitum rubicundum Fisch. ex Steud., 2n = 16; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1638.

Anemone baicalensis Turcz., 2n = 28; Russia, Irkutskaya Oblast', MP, VP, EZ & AK Cl643; MP, VP, EZ & AK Cl645.

Aquilegia glandulosa Fisch. ex Link, 2n = 14; Russia, Irkutskaya Oblast', MP, VP & EZ C1668.

Caltha palustris L., 2*n* = 32; Russia, Irkutskaya Oblast', *MP*, *VP*, *EZ* & *AK* C1662.

VIOLACEAE

Viola biflora L., 2n = 12; Russia, Irkutskaya Oblast', MP, VP, EZ & AK C1649.

Iva Hodálová,^{1*} Lenka Mártonfiová,² Marilena Onete,³ Jaromír Kučera¹ & Pavol Mereďa, Jr.¹

- 1 Institute of Botany, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovak Republic
- 2 Pavol Jozef Šafárik University, Botanical Garden, Mánesova 23, 04352 Košice, Slovak Republic
- 3 Institute of Biology Bucharest, Romanian Academy, 296 Splaiul Independentei, 060031 Bucharest, Romania
- * Author for correspondence: iva.hodalova@savba.sk

Chromosomes counted by L. Mártonfiová. DNA ploidy levels estimated by P. Mereďa, Jr.; collectors: IH = I. Hodálová, MZ = M. Zaliberová, PM = P. Mereďa, Jr.; vouchers in SAV.

This study was supported by the Grant Agency of Ministry of Education, Science, Research and Sport of the Slovak Republic and Slovak Academy of Sciences VEGA (grant no. 2/0131/16).

POACEAE

Sesleria heufleriana Schur

2n = 4x = 28, CHN. Romania, IH, PM & MZ 137-1. 2n ~ 4x ~ 28, FCM. Romania, IH, PM & MZ 137-1, IH, PM & MZ 137-2, IH, PM & MZ 137-3, IH, PM & MZ 137-4, IH, PM & MZ 137-5, IH, PM & MZ 137-6, IH, PM & MZ 137-7, IH, PM & MZ 137-8, IH, PM & MZ 137-9, IH, PM & MZ 137-10, IH, PM & MZ 137-11, IH, PM & MZ 137-12, IH, PM & MZ 137-13, IH, PM & MZ 137-14, IH, PM & MZ 137-15, IH, PM & MZ 141-1, IH, PM & MZ 141-2, IH, PM & MZ 141-5, IH, PM & MZ 141-6, IH, PM & MZ 141-7, IH, PM & MZ 141-8, IH, PM & MZ 141-9, IH, PM & MZ 141-10, IH, PM & MZ 141-13, IH, PM & MZ 141-12, IH, PM & MZ 141-14, IH, PM & MZ 141-15, IH, PM & MZ 142-1, IH, PM & MZ 142-2, IH, PM & MZ 142-3, IH, PM & MZ 142-4, IH, PM & MZ 142-5, IH, PM & MZ 142-6, IH, PM & MZ 142-7, IH, PM & MZ 142-8, IH, PM & MZ 142-9, IH, PM & MZ 142-10, IH, PM & MZ 142-11, IH, PM & MZ 142-12, IH, PM & MZ 142-13, IH, PM & MZ 142-14, IH, PM & MZ 142-15.

Sesleria rigida Rchb.

2n = 4*x* = 28, CHN. Romania, *IH*, *PM & MZ 139-1*.

2n ~ 4x ~ 28, FCM. Romania, IH, PM & MZ 138-1, IH, PM & MZ 138-2, IH, PM & MZ 138-3, IH, PM & MZ 138-4, IH, PM & MZ 138-5, IH, PM & MZ 138-6, IH, PM & MZ 138-7, IH, PM & MZ 138-8, IH, PM & MZ 138-9, IH, PM & MZ 138-10, IH, PM & MZ 138-11, IH, PM & MZ 139-1, IH, PM & MZ 139-2, IH, PM & MZ 139-3, IH, PM & MZ 139-4, IH, PM & MZ 139-5, IH, PM & MZ 139-6, IH, PM & MZ 139-7, IH, PM & MZ 140-1, IH, PM & MZ 140-2, IH, PM & MZ 143-1, IH, PM & MZ 143-2, IH, PM & MZ 143-4, IH, PM & MZ 143-5, PM 144-1, PM 144-2, PM 144-3, PM 144-4, PM 144-5, PM 144-6, PM 144-7, PM 144-8, PM 144-9, PM 144-10, PM 144-11, PM 144-12, PM 144-13, PM 144-14.

Navjot Kaur* & Raghbir Chand Gupta

Department of Botany, Punjabi University Patiala, India * Author for correspondence: navjot21188@gmail.com

The study is supported by financial grant under DRS SAP I, SAP II and SAP III of UGC; UGC-BSR Indira Gandhi Single Girl Child Fellowship (Award letter no. F7-152/2007 BSR, dated 16/12/2013) to Navjot Kaur and IPLS-DBT project (Project no. BT/PR-4548/ INF/22/146/2012) sanctioned to Punjabi University, Patiala for using the facilities and financial support of this study.

All materials CHN. Collected in India, Rajasthan; collector: *NK* = Navjot Kaur; vouchers in PUN.

ASPARAGACEAE

Asparagus racemosus Willd., n = 30; NK 60936.

CANNACEAE

Canna indica L., n = 18; NK 61126.

COMMELINACEAE

Commelina attenuata Vahl, n = 12; NK 60938. Commelina diffusa Burm.f., n = 60; NK 60779. Commelina forskalaei Vahl, n = 45; NK 60937. Commelina hasskarlii C.B.Clarke (= C. caroliniana Walter), n = 60; NK 61120, NK 61121. Commelina paludosa Blume, n = 15; NK 61116, NK 61117. Commelina suffruticosa Blume, n = 15; NK 61116, NK 61117. Commelina suffruticosa Blume, n = 15; NK 61116, NK 61117. Murdannia nudiflora (L.) D.Don, n = 11; NK 61115. Murdannia nudiflora (L.) Brenan, n = 40; NK 61119.

POACEAE

Brachiaria deflexa (Schumach.) C.E.Hubb. ex Robyns, n = 18; NK 60211, NK 60217. Chloris prieurii Kunth, n = 10; NK 60164, NK 60186. Chloris quinquesetica Bhide, n = 10; NK 60208. Eragrostis aspera (Jacq.) Nees, n = 10; NK 60980. Leptothrium senegalense (Kunth) Clayton, n = 10; NK 60149. Sporobolus coromandelianus (Retz.) Kunth, n = 24; NK 60167.

Jaromír Kučera,^{1*} Eliška Štubňová,^{1,2} Lenka Mártonfiová,³ Filip Kolář,^{4,5} Jan Smyčka⁶ & Marek Slovák^{1,4}

- 1 Plant Science and Biodiversity Centre, Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovak Republic
- 2 Department of Botany, Natural History Museum, Slovak National Museum, Vajanského nábrežie 2, 810 06 Bratislava, Slovak Republic
- 3 Pavol Jozef Šafárik University, Botanical Garden, Mánesova 23, Košice, Slovak Republic
- 4 Department of Botany, Charles University, Benátská 2, 128 01 Praha 2, Czech Republic
- 5 Institute of Botany, The Czech Academy of Sciences, Zámek 1, 252 43 Průhonice, Czech Republic
- 6 Laboratoire d'Ecologie Alpine (LECA), Université Grenoble Alpes-CNRS, 2233 rue de la Piscine, 38000 Grenoble, France
- * Author for correspondence: jaromir.kucera@savba.sk

Genome size data (C-values) were measured by E. Štubňová and J. Kučera. Chromosome numbers were counted by L. Mártonfiová. Collectors: $E\check{S} = E$. Štubňová, FK = Filip Kolář, JK = J. Kučera, MS = M. Slovák, JS = J. Smyčka, MV = M. Valachovič; vouchers in SAV.

This study was supported by the Grant Agency VEGA, Bratislava, Slovak Republic (grant no. VEGA 2/0088/15).

PRIMULACEAE

Soldanella alpina L. subsp. alpina

2n = 4x = 40, CHN. Austria, JK & MS AT39/2; Switzerland, JK & MS SW5/5.

2n ~ 4x ~ 40, 2C = 3.52–3.68 pg, FCM. Austria, JK & MS AT37/1, JK & MS AT37/2, JK & MS AT37/5, JK & MS AT38/1, JK & MS AT38/2, JK & MS AT38/3, JK & MS AT39/1, JK & MS AT39/2, JK & MS AT39/3, FK AT41/1, FK AT41/2, FK AT41/3; France, JS FR2/1, JS FR2/2, JS FR2/3, JS FR3/1, JS FR3/2, JS FR3/3; Italy, JK & MS IT22/1, JK & MS IT22/2, JK & MS IT22/3; Switzerland, JK & MS SW2/1, JK & MS SW2/2, JK & MS SW2/4, JK & MS SW4/1, JK & MS SW4/2, JK & MS SW4/3, JK & MS SW5/2, JK & MS SW5/3, JK & MS SW5/5.

Soldanella angusta L.B.Zhang

- 2n = 4x = 40, CHN. Ukraine, *MV UA8/13*.
- $2n \sim 4x \sim 40$, 2C = 3.31-3.38 pg, FCM. Ukraine, *MV UA6/2*, *MV UA6/5*, *MV UA6/15*, *MV UA7/4*, *MV UA7/7*, *MV UA7/9*, *MV UA8/11*, *MV UA8/13*, *MV UA8/14*.

Soldanella ×hybrida A.Kern

2n = 4x = 40, CHN. Switzerland, JK & MS SW4/4.

2*n*~4*x*~40, 2C = 3.57–3.62 pg, FCM. Switzerland, *JK* & *MS SW*4/4,

JK & MS SW4/5, JK & MS SW4/6.

Soldanella major (Nielr.) Vierh.

2n = 4x = 40, CHN. Romania, *EŠ RO23/3*.

2*n* ~ 4*x* ~ 40, 2C = 3.28–3.32 pg, FCM. Romania, *EŠ RO23/1, EŠ RO23/2, EŠ RO23/3*.

Soldanella minima Hoppe

 $2n \sim 4x \sim 40$, 2C = 3.63-3.68 pg, FCM. Austria, *FK AT42/1*, *FK AT42/2*, *FK AT42/3*.

Soldanella pusilla subsp. alpicola (F.K.Mey.) Chrtek

2n = 4x = 40, CHN. Switzerland, JK & MS SW3/2.

2n~4x~40, 2C = 3.43–3.52 pg, FCM. Austria, JK AT35/1, JK AT35/2, JK AT35/3, JK AT36/1, JK AT36/2, JK AT36/3; Switzerland, JK & MS SW1/1, JK & MS SW1/2, JK & MS SW1/2, JK & MS SW3/2, JK & MS SW3/2, JK & MS SW3/3, JK & MS SW4/7, JK & MS SW4/8, JK & MS SW4/9, JK & MS SW6/1, JK & MS SW6/2, JK & MS SW6/3, JK & MS SW7/1, JK & MS SW7/3, JK & MS SW7/5, JK & MS SW8/1, JK & MS SW8/2, JK & MS SW8/2, JK & MS SW8/3.

Soldanella ×wettsteinii Vierh.

2*n*~4*x*~40, 2C = 3.51–3.56 pg, FCM. Austria, *JK*, *MS* & *EŠ AT*40/1, *JK*, *MS* & *EŠ AT*40/2, *JK*, *MS* & *EŠ AT*40/3, *JK*, *MS* & *EŠ AT*40/4.

Rohit Kumar, Vandna Kumari & Vijay Kumar Singhal*

Department of Botany, Punjabi University, Patiala-147002, Punjab, India

* Author for correspondence: vksinghal53@gmail.com

The study was supported by financial grant under DRS SAP I, SAP II and SAP III of University Grant Commision (UGC), New Delhi, India; Women Scientist Fellowship Scheme (WOS-A) of Department of Science and Technology, Government of India (GOI) to Vandna Kumari, Rajiv Gandhi National Fellowship Scheme of UGC to Rohit Kumar (award letter no. F1-17.1/2012-13 /RGNF-2012-13-SC HIM-18573) and IPLS project of Department of Biotechnology, GOI (BT/PR4548/INF/22/146/2012) sanctioned to Punjabi University Patiala, Punjab for using the facilities and financial support of this study.

All materials CHN; collectors: RK = Rohit Kumar; VK = Vandna Kumari; vouchers in PUN.

BRASSICACEAE

Turritis glabra L., *n* = 7; India, Uttarakhand, *RK 34335*.

CARYOPHYLLACEAE

Arenaria festucoides Benth., n = 11; India, Uttarakhand, RK 34336.

FABACEAE

Astragalus uttaranchalensis L.B.Chowdhary & Z.H.Khan, *n* = 8; India, Uttarakhand, *RK 34862*.

Medicago edgeworthii Širj., n = 8; India, Uttarakhand, *RK 34903. Thermopsis barbata* Benth., n = 9; India, Uttarakhand, *RK 34916.*

HYPERICACEAE

Hypericum elodeoides Choisy, n = 10; India, Uttarakhand, RK 34358.

PAPAVERACEAE

Corydalis stracheyi Duthie ex Prain, n = 8; India, Uttarakhand, *RK 34310.*

POACEAE

Digitaria abludens (Roem. & Schult.) Veldkamp, n = 36; India, Himachal Pradesh, *VK 32435*.

Festuca sibirica Hack. ex Boiss., n = 21; India, Himachal Pradesh, *VK 34676*.

Hordeum murinum L., n = 7; India, Himachal Pradesh, VK 34704.

Leptochloa chinensis (L.) Nees, *n* = 9; India, Himachal Pradesh, *VK* 34712.

Microstegium vimineum (Trin.) A.Camus, n = 10; India, Tosh, *VK* 34711.

Poa setulosa Bor, n = 14; India, Himachal Pradesh, *VK 34740*.

Puccinellia himalaica Tzvelev, *n* = 14; India, Himachal Pradesh, *VK* 34689.

Stipa jacquemontii Jaub. & Spach, *n* = 10; India, Himachal Pradesh, *VK 34694*.

SAXIFRAGACEAE

Saxifraga pedemontana subsp. cymosa Engl., n = 8; India, Uttarakhand, *RK 34981*.

TAMARICACEAE

Myricaria elegans Royle, *n* = 8; India, Uttarakhand, *RK 34988. Myricaria rosea* W.W.Sm., *n* = 8; India, Uttarakhand, *RK 34990.*

M. Laura Las Peñas

Instituto Multidisciplinario de Biología Vegetal (IMBIV), CONICET-UNC, Universidad Nacional de Córdoba, CC 495 Córdoba 5000, Argentina; laulaspenas@yahoo.com.ar

All materials CHN; collectors: DU = Diego Uñates; FC = Franco Chiarini, JU = Juan Urdampilleta, MLLP = M. Laura Las Peñas; vouchers in CORD.

Funding was provided by CONICET, FONCyT, and SECyT-UNC (FCEFyN) from Argentina.

CACTACEAE

Arrojadoa rhodantha (Gürke) Britton & Rose, 2n = 22; Brasil, Rio de Contas, *JU & al. 379*.

- Cereus aethiops Haw., 2n = 22; Argentina, Mendoza, FC 614.
- Cleistocactus smaragdiflorus (F.A.C.Weber) Britton & Rose, 2n = 22; Argentina, Salta, *MLLP & DU 279*.
- Denmoza rhodacantha (Salm-Dyck) Britton & Rose, 2n = 22; Argentina, La Rioja, MLLP & FC 212.

Echinopsis aurea var. *dobeana* (Dölz) J.Ullmann, 2*n* = 22; Argentina, Catamarca, *MLLP 125*.

Echinopsis aurea var. *falax* (Oehme) J.Ullmann, 2*n* = 22; Argentina, La Rioja, *MLLP & FC 203*.

- *Echinopsis leucantha* (Gillies ex Salm-Dyck) Walp., 2n = 22; Argentina, San Juan, *MLLP & DU 87*.
- *Gymnocalycium andreae* (Boed.) Backeb. & F.M.Knuth, 2n = 44; Argentina, Córdoba, *MLLP & DU 231*.

Gymnocalycium monvillei subsp. *achirasense* (H.Till & Schatzl) H.Till, 2n = 22; Argentina, Cordoba, *MLLP & al. 363*.

Gymnocalycium monvillei (Lem.) Britton & Rose subsp. *monvillei*, 2n = 44; Argentina, Córdoba, *MLLP & DU 232*.

Harrisia hahniana (Backeb.) Kimnach & Hutchison ex Kimnach, 2*n* = 22; Paraguay, Presidente Hayes, *MLLP 588*.

- Harrisia pomanensis (F.A.C.Weber ex K.Schum.) Britton & Rose, 2n = 44; Argentina, San Luis, *MLLP & DU 100.*
- Lobivia saltensis (Speg.) Britton & Rose, 2n = 22; Argentina, Salta, MLLP 138.
- Maihueniopsis darwinii (Hensl.) F.Ritter, 2n = 55; Argentina, Chubut, MLLP & DU 349.

Maihueniopsis glomerata (Haw.) R.Kiesling, 2n = 44; Argentina, San Juan, *MLLP & DU 178*.

- Maihueniopsis ovata (Pfeiff.) F.Ritter, 2n = 44; Argentina, San Juan, MLLP & DU 181.
- *Melocactus bahiensis* (Britton & Rose) Luetzelb, 2n = 44; Brasil, Rio de Contas, *JU & al. 387*.
- *Opuntia quimilo* K.Schum., 2n = 22; Argentina, La Rioja, *MLLP & al. 539.*
- *Oreocereus celsianus* (Lem. ex Salm-Dyck) Riccob., 2n = 44; Argentina, Jujuy, *MLLP & DU 274*.
- *Oreocereus trollii* (Kupper) Backeb., 2n = 22; Argentina, Jujuy, *MLLP* & *DU* 339.
- *Pachycereus pringlei* (S.Watson) Britton & Rose, 2*n* = 22; México, Dpt. La Paz, *MLLP 714*.
- *Quiabentia verticillata* (Vaupel) Vaupel ex A.Berger, 2n = ca. 187; Bolivia, Dpt. Mizque, *JU & al.* 542.
- *Rhipsalis lumbricoides* (Lem.) Lem. ex Salm-Dyck, 2n = 22; Argentina, Misiones, *MLLP & DU 56*.

- *Tacinga inamoena* (K.Schum.) Taylor & Stuppy, 2*n* = 44; Brasil, Rio de Contas, *JU & al. 385*.
- *Trichocereus arboricola* Kimnach, 2*n* = 22; Argentina, Salta, *MLLP* & *FC* 715.
- *Trichocereus atacamensis* (Phil.) W.T.Marshall & T.M.Bock, 2n = 22; Argentina, Salta, *MLLP & al. 319*.
- Trichocereus macrogonus Britton & Rose, 2n = 22; Perú, MLLP 122.
- *Trichocereus thelegonus* (F.A.C.Weber ex K.Schum.) Britton & Rose, 2n = 22; Argentina, Salta, *MLLP & al. 283*.

Rohit N. Mane,¹ Kumar Vinod C. Gosavi,^{1,2*} Avinash R. Gholave^{1,3} & Shrirang R. Yadav¹

- 1 Department of Botany, Shivaji University, Kolhapur (MS) 416 004, India
- 2 Department of Botany, HPT Arts & RYK Science College, Nashik (MS) 422 002, India
- 3 Life Science Research Laboratory, Walchand College of Arts and Science, Solapur (MS) 413 006, India
- * Author for correspondence: kumarvinodgosavi@gmail.com

All materials CHN; vouchers in SUK.

Authors express thanks to Head, Department of Botany, Shivaji University, Kolhapur for providing laboratory facilities, SERB (File No.: SB/FT/LS-130/2012).

ACANTHACEAE

Barleria gibsonii Dalzell, 2*n* = 44; India, Maharashtra, *K.V.C. Gosavi* 126.

- *Barleria prattensis* Santapau, 2*n* = 44; India, Maharashtra, *K.V.C. Gosavi* 125
- *Pleocaulus sessilis* (Nees) Bremek., 2n = 30; India, Maharashtra, *R.N. Mane 56*.

Ruellia urens Roth, 2n = 30; India, Maharashtra, R.N. Mane 54.

Strobilanthes callosa Nees, 2n = 30; India, Maharashtra, K.V.C. Gosavi 452.

ARECACEAE

Trachycarpus takil Becc., 2*n* = 36; India, Uttarakhand, *S.R. Yadav* 226.

DIPTEROCARPACEAE

Vatica chinensis L., 2n = 22; India, Karnataka, R.N. Mane 67.

FABACEAE

Clitoria biflora Dalzell, 2n = 14; India, Maharashtra, K.V.C. Gosavi 186.

POACEAE

Hemarthria protensa Nees ex Steud., *n* = 9; India, Assam, *K.V.C. Gosavi 171.*

Lasiurus scindicus Henrard, n = 18; India, Rajasthan, *K.N. Koli 167. Manisuris myurus* L., n = 8; India, Tamil Nadu, *K.V.C. Gosavi 196. Mnesithea striata* (Nees ex Steud.) de Koning & Sosef, n = 18; India,

Meghalaya, K.V.C. Gosavi 235.

SAPOTACEAE

Chrysophyllum roxburghii G.Don, 2*n* = 26; India, Karnataka, *R.N. Mane 61.*

José Ignacio Márquez-Corro,^{1#*} Modesto Luceño,^{1#} Pedro Jiménez-Mejías,² Marcial Escudero,³ Santiago Martín-Bravo,¹ Andrew L. Hipp,^{4,5} Kyong-Sook Chung,⁶ A. Muthama Muasya,⁷ Paul E. Rothrock,⁸ Jaime A. Weber⁴ & Robert Naczi⁹

- 1 Department of Molecular Biology and Biochemical Engineering, Universidad Pablo de Olavide, Carretera de Utrera km 1, 41013 Seville, Spain
- 2 Biology (Botany), Universidad Autónoma de Madrid, Ciudad Universitaria de Cantoblanco, Calle Francisco Tomás y Valiente, 7, 28049 Spain
- 3 Department of Plant Biology and Ecology, University of Seville, Reina Mercedes sn, 41012 Seville, Spain
- 4 The Morton Arboretum, 4100 Illinois Route 53, Lisle, Ilinois 60532, U.S.A.
- 5 The Field Museum, Integrative Research Center, 1400 S Lake Shore Dr., Chicago, Illinois 60605, U.S.A.
- 6 Department of Medicinal Plant Science, Jungwon University, Goesan-gun, Chungbuk 28024, Korea
- 7 Department of Biological Sciences, University of Cape Town, Private Bag X3, Rondebosch, 7701 Cape Town, South Africa
- 8 Department of Biology, Indiana University, Bloomington, Indiana 47405, U.S.A.
- 9 New York Botanical Garden, 2900 Southern Boulevard, Bronx, New York 10458-5126, U.S.A.
- [#] The two first authors have equally contributed to this article
- * Author for correspondence: jimarcorr@gmail.com

All material CHN.

CYPERACEAE SUBFAMILY CYPEROIDEAE TRIBE ABILDGAARDIEAE

Fimbristylis spadicea (L.) Vahl, 2*n* = 20; Brazil, Pernambuco, 15 Aug 1996, *Luceño & al. s.n.* (UFP).

TRIBE BISBOECKELEREAE

Calyptrocarya glomerulata (Brongn.) Urb., 2*n* = 20; Brazil, Pernambuco, *Luceño & al. s.n.* (UFP).

TRIBE CARICEAE

Carex acidicola Naczi, 2*n* = 50; U.S.A., Georgia, *Naczi 1783* (MICH, NY).

Carex aethiopica Schkuhr, 2*n* = 74, 76; South Africa, Cape Town, 12 Sep 2009, *Luceño & al. s.n.* (BOL).

- *Carex amphibola* Steud., 2n = 56; U.S.A., Arkansas, *Naczi 2829* (MICH, NY).
- *Carex asynchrona* Naczi, 2*n* = 52; Mexico, Tamaulipas, *Naczi 2949* (MICH).
- *Carex bolanderi* Olney, 2*n* = 50; U.S.A., Oregon, *Zika* 24689 (MOR). 2*n* = 58; U.S.A., New Mexico, *Norris, Kleinman & Walkup* 2008-7-23-43 (SNM).
- Carex borbonica Lam., 2n = 68; France, La Réunion, Luceño & Guzmán 1ML09 (UPOS-3635).
- *Carex borbonica* Lam. × *C. boryana* Schkuhr, 2*n* = 68; France, La Réunion, *Luceño & Guzmán 20ML09* (UPOS-3724).
- Carex boryana Schkuhr, 2n = 68; France, La Réunion, Luceño & Guzmán 14ML09 (UPOS-3641).
- *Carex brevior* (Dewey) Mack. ex Lunell, 2n = 60; U.S.A., Iowa, *Norris & Thompson s.n.* (MOR).

- *Carex brysonii* Naczi, 2*n* = 50; U.S.A., Alabama, *Naczi 2950* (MICH, NY).
- *Carex bulbostylis* Mack., 2*n* = 54; U.S.A., Texas, *Naczi 2328* (MICH, NY).
- *Carex calcifugens* Naczi, 2*n* = 52; U.S.A., South Carolina, *Naczi 2059* (MICH, NY, PH).
- Carex cespitosa L., 2n = 78; Spain, Jiménez-Mejías & al. 99PJM06 (UPOS-1930).
- *Carex conoidea* Schkuhr ex Willd., 2*n* = 68; Canada, New Brunswick, *Naczi 2347* (MICH, NY).
- *Carex corrugata* Fernald, 2*n* = 58; U.S.A., Virginia, *Naczi 2238* (MICH, NY).
- *Carex cusickii* Mack. ex Piper & Beattie, 2*n* = 63; U.S.A., California, *Janeway & Gog 3197C* (MOR).
- Carex demissa Hornem., 2n = 70; Portugal, Jiménez-Mejías & al. 06PJM05 (UPOS), Jiménez-Mejías & al. 185PJM05 (UPOS-2866); Spain, Jiménez-Mejías & al. 201PJM04 (UPOS-3518), Jiménez-Mejías & al. 28PJM03 (UPOS).
- Carex demissa Hornem. × C. lepidocarpa Tausch, 2n = 68; Spain, Marín 15504JMM (UPOS-3577), Jiménez-Mejías & al. 451PJM05 (UPOS-2615). 2n = 70; Spain, Luceño & al. 9205MLbis (UPOS-1405).
- *Carex densa* (L.H.Bailey) L.H.Bailey, 2n = 54; U.S.A., California, *Janeway & Gog 3183A* (MOR).
- *Carex distans* L., 2*n* = 70; Turkey, *Martín-Bravo & al. 112SMB10* (UPOS-7401).
- *Carex ecklonii* Nees, 2*n* = 64; South Africa, 12 Sep 2009, *Luceño & al. s.n.* (UPOS-6503).
- *Carex edwardsiana* E.L.Bridges & Orzell, 2*n* = 52; U.S.A., Texas, *Naczi 2321* (MICH, NY).
- *Carex flaccosperma* Dewey, 2*n* = 46; U.S.A., Arkansas, *Naczi 2733* (MICH, NY).
- *Carex fracta* Mack., 2n = 70; U.S.A., Oregon, *Wilson & Otting 15143* (MOR).
- *Carex glaucodea* Tuck. ex Olney, 2*n* = 44; Canada, Ontario, *Naczi* 2954 (MICH, NY).
- *Carex glomerabilis* V.I.Krecz., 2*n* = 60; South Africa, *Márquez-Corro* & *al.* 196JMC17 (UPOS-9968).
- *Carex godfreyi* Naczi, 2*n* = 50; U.S.A., Florida, *Naczi* 2740 (MICH, NY).
- *Carex grisea* Wahlenb., 2*n* = 56; Canada, Québec, *Naczi 1768* (MICH, NY).
- *Carex hitchcockiana* Dewey, 2*n* = 54; Canada, Ontario, *Naczi 1786* (MICH, NY).
- *Carex impressinervia* Bryson, Kral & Manhart, 2*n* = 46; U.S.A., Mississippi, *Naczi 2956* (MICH, NY).
- *Carex leporinella* Mack., 2*n* = 82; U.S.A., Idaho, *Wilson & al. 15858* (MOR).
- *Carex mariposana* L.H.Bailey ex Mack., 2*n* = 81; U.S.A., California, *Taylor & al. 20752* (MOR).
- *Carex mesochorea* Mack., 2n = 48; U.S.A., Michigan, 14 Jul 2011, *Reznicek s.n.* (MOR).
- *Carex molesta* Mack. ex Bright, 2*n* = 72; U.S.A., Wisconsin, *Hipp* & Gog 3137 (MOR).
- *Carex nervina* L.H.Bailey, 2*n* = 52; U.S.A., California, *Janeway & Gog 3195C* (MOR).
- Carex nigra (L.) Reichard, 2n = 83; Portugal, Martín-Bravo 216SMB06 (UPOS); Spain, Marín & Luceño 40JMM04 (UPOS), Jiménez-Mejías & al. 31PJM06 (UPOS-2253), Jiménez-Mejías & Escudero 160PJM06 (UPOS), Martín-Bravo & al. 153SMB07 (UPOS). 2n =

84; France, Corse, Escudero & Luceño 89ME07 (UPOS); Greece, Vargas & Luceño 284PV04 (UPOS-831); Portugal, Martín-Bravo 216SMB06 (UPOS); Spain, Jiménez-Mejías & García 153PJM06 (UPOS), Jiménez-Mejías 36PJM06 (UPOS), Jiménez-Mejías & al. 136PJM06 (UPOS-2220), Jiménez-Mejías & Escudero 157PJM06 (UPOS), Jiménez-Mejías & al. 137PJM06 (UPOS-2221). 2n = 85; Morocco, Chaparro & al. 07AJC05 (UPOS-1632), Jiménez-Mejías & al. 88PJM07 (UPOS-4750); Spain, Jiménez-Mejías & García 153PJM06 (UPOS), Jiménez-Mejías & al. 135PJM06 (UPOS-2218). 2n = 86; Morocco, Jiménez-Mejías & al. 88PJM07 (UPOS-4750).

- Carex oedipostyla Duval-Jouve, 2n = 68; Spain, Luceño & al. 1ML11 (UPOS-6228), Maguilla & al. 1EMS13 (UPOS-5127).
- *Carex oligocarpa* Schkuhr ex Willd., 2*n* = 54; U.S.A., Illinois, *Naczi* 2902 (MICH, NY).
- *Carex oronensis* Fernald, 2*n* = 68, 71, 74; U.S.A., Maine, *Reznicek 11968* (MOR).
- *Carex ouachitana* Kral, Manhart & Bryson, 2*n* = 52; U.S.A., Arkansas, *Naczi 1774* (MICH, NY).
- *Carex pachycarpa* Mack., 2*n* = 70; U.S.A., Oregon, *Otting & Wilson* 1945B (MOR), *Wilson & Otting* 14947 (MOR), *Wilson & Otting* 14948 (MOR), *Wilson & N. Otting* 14949 (MOR), *Wilson & Otting* 15017 (MOR). 2*n* = 74; U.S.A., Oregon, *Wilson* 15018 (MOR).
- *Carex pachystachya* Cham. ex Steud., 2n = 82; U.S.A., Oregon, *Wilson 15052* (MOR).
- *Carex paeninsulae* Naczi, E.L.Bridges & Orzell, 2*n* = 48; U.S.A., Florida, *Naczi 2958* (MICH, NY).
- *Carex petasata* Dewey, 2n = 74; U.S.A., Oregon, *Wilson 15019* (MOR). *Carex pigra* Naczi, 2n = 56; U.S.A., Alabama, *Naczi 2735* (MICH,
- NY).
- *Carex planispicata* Naczi, 2*n* = 50; U.S.A., Alabama, *Naczi 2060* (MICH, NY).
- *Carex projecta* Mack., 2n = 62; U.S.A., Michigan, *Rothrock 4070* (MOR). 2n = 64; U.S.A., Ontario, *Rothrock 3971* (MOR). 2n = 66; U.S.A., Maine, *Rothrock 3970* (MOR). U.S.A., Wisconsin, *Rothrock 4289* (MOR).
- *Carex radiata* (Wahlenb.) Small, 2*n* = 58; U.S.A., Wisconsin, *Hipp* & *Gog 3133* (MOR).
- *Carex rainbowii* Luceño, Jim.Mejías, M.Escudero & Martín-Bravo, 2n = 58; South Africa, KwaZulu-Natal, *Martín-Bravo & Luceño* 120SMB11 (UPOS-5030).
- *Carex retroflexa* Muhl. ex Willd., 2*n* = 42; U.S.A. Michigan, 14 Jun 2009, *Reznicek & Walters s.n.* (MOR).
- *Carex reuteriana* subsp. *mauritanica* (Boiss. & Reut.) Jim.Mejías & Luceño, 2n = 74; Spain, *Jiménez-Mejías & al. 22PJM05* (UPOS), *Escudero & al. 4ME05* (UPOS).
- *Carex simulata* Mack., 2*n* = 64; U.S.A., Colorado, *Chung & Lederer* 1039 (MOR).
- *Carex subbracteata* Mack., 2*n* = 80; U.S.A., Washington, *Zika* 24641 (MOR).
- *Carex subfusca* W.Boott, 2*n* = 82; U.S.A., Oregon, *Wilson & al. 15068* (MOR). 2*n* = 86; U.S.A., Nevada, *Zika 24619* (MOR).
- *Carex thornei* Naczi, 2n = 50; U.S.A., Georgia, *Naczi 1779* (MICH, NY).
- *Carex tincta* (Fernald) Fernald, 2*n* = 72; U.S.A., Maine, *Reznicek* 10919A (MOR).
- *Carex tribuloides* Wahlenb., 2*n* = 70; U.S.A., Indiana, *Rothrock 4243* (MOR), *Rothrock 3839C* (MOR). 2*n* = 72; U.S.A., Indiana, *Rothrock 3839A* (MOR).
- *Carex trinervis* Degl., 2*n* = 82, 84; Portugal, Coimbra, *Jiménez-Mejías* & *al.* 10PJM05 (UPOS).

Carex vernacula L.H.Bailey, 2*n* = 52; U.S.A., Colorado, *Chung 1063* (MOR).

Carex viridula Michx., 2n = 70; Greece, *Luceño 1404ML* (UPOS-691). *Carex vixdentata* (Kük.) G.A.Wheeler, 2n = 62; Argentina, Buenos Aires, *Marín 2102JMM* (UPOS-849).

TRIBE CARPHEAE

Trianoptiles capensis (Steud.) Harv., 2n = 52; South Africa, Cape Town, *Muasya & al. 4549* (BOL).

TRIBE CRYPTANGIEAE

Lagenocarpus guianensis Nees, 2*n* = 34; Brazil, Pernambuco, *Luceño* & *al. s.n.* (UFP).

TRIBE CYPEREAE

Afroscirpoides dioeca (Kunth) García-Madr. & Muasya (≡ Scirpoides dioeca (Kunth) J.Browning), 2n = 35; South Africa, Muasya & al. 4557 (BOL).

Cyperus fuscus L., 2n = 36; Spain, M. Luceño s.n. (UPOS).

Cyperus salzmannianus (Steud.) Bauters (= Lipocarpha salzmanniana Steud.), 2n = 38; Brazil, Rio Grande do Norte, M. Luceño & al. s.n. (UFP). 2n = 36, 38, 40; Brazil, Pernambuco, Luceño & Guerra s.n. (UFP).

Cyperus sesquiflorus (Torr.) Mattf. & Kük. (= *Kyllinga odorata* Vahl), 2*n* = 108; Brazil, Pernambuco, *Luceño & al. s.n.* (UFP).

Cyperus subsquarrosus (Muhl.) Bauters (= Hemicarpha micrantha (Vahl) Britton), 2n = 52; Brazil, Pernambuco, Luceño & al. s.n. (UFP). 2n = 54; Brazil, Pernambuco, Luceño & al. s.n. (UFP).

Dracoscirpoides ficinioides (Kunth) Muasya, 2n = 104; South Africa, Maguilla & al. 79EMS12 (UPOS).

Isolepis marginata (Thunb.) A.Dietr., 2n = 8; South Africa, *Muasya & al. 4516* (BOL).

TRIBE ELEOCHARIDEAE

Eleocharis confervoides (Poir.) Steud. (= *Websteria submersa* (C.Wright) Britton), 2n = 38, 40, 41; Brazil, Pernambuco, *Luceño & al. s.n.* (UFP).

TRIBE RHYNCHOSPOREAE

Rhynchospora brownii Roem. & Schult., 2n = 36; South Africa, *Luceño & al. 125ML08* (UPOS-3707).

Rhynchospora modesti-lucennoi Castrov., 2n = 84; Portugal, Jiménez-Mejías & al. 113PJM05 (UPOS-1394).

TRIBE SCHOENEAE

Schoenus nigricans L., 2n = 76; Spain, Jiménez-Mejías & al. 3PJM07 (UPOS-2857).

SUBFAMILY MAPANIOIDEAE TRIBE CHRYSITRICHEAE

Chrysitrix capensis L., 2*n* = 46; South Africa, *Muasya & al. 4540* (BOL).

Antonina V. Natyaganova,¹ Elena V. Mincheva,¹ Yurii S. Bukin,¹ Lyubov S. Kravtsova,¹ Tatiana E. Peretolchina,¹ Tatiana I. Triboy¹ & Denis A. Krivenko^{2,3*}

- 1 Limnological Institute of the Siberian Branch of the Russian Academy of Sciences, Ulan-Batorskaya Str. 3, 664033 Irkutsk, Russia
- 2 Siberian Institute of Plant Physiology & Biochemistry of the Siberian Branch of the Russian Academy of Sciences, Lermontov Str. 132, 664033 Irkutsk, Russia
- 3 Western Baikal Protected Areas, Baikalskaya Str. 291B, 664050 Irkutsk, Russia
- * Author for correspondence: krivenko.irk@gmail.com

All materials CHN; collectors: AVN = Antonina V. Natyaganova, EVM = Elena V. Mincheva, YuSB = Yurii S. Bukin; vouchers in IRK.

The research was supported by the state project LIN SB RAS #AAAA-A16-116122110060-9.

HYDROCHARITACEAE

Elodea canadensis Michx., 2n = 48; Russia, Irkutskaya Oblast', AVN & YuSB 50785, AVN 50789, AVN 50786, Yu.R. Zakharova & M.V. Bashenkhaeva 50791, AVN 50787, EVM 50788, M.P. Tskhovrebova 50795, AVN 50796, D.A. Krivenko & AVN 50771; Russia, Republic of Buryatiya, AVN 50790, AVN 50792, L.S. Kravtsova 50793; Russia, Zabaikal'skii Krai, EVM 50794; Russia, Moskovskaya Oblast', YuSB 51398.

Egeria densa Planch., 2*n* = 48; Russia, Moskovskaya Oblast', *AVN* 51053.

Saroj Kumar Pradhan,^{1*} Raghbir Chand Gupta¹ & Rajesh Goel²

- 1 Department of Botany, Punjabi University, Patiala, Punjab, 147002, India
- 2 Department of Pharmaceutical Sciences and Drug Research, Punjabi University, Patiala, Punjab, 147002, India
- * Author for correspondence: pradhan_sarojkumar@yahoo.com

All materials CHN; collected in India; collector: *SKP* = *Saroj Kumar Pradhan*; vouchers in PUN.

The study is supported by financial grant under DBT-IPLS Project (Reference number: BT/PR 4548/INF/22/146/2012).

ASTERACEAE

Arctium lappa L., n = 17; SKP 32112. Carduus edelbergii Rech.f., n = 16; SKP 32149. Cremanthodium arnicoides (DC. ex Royle) R.D.Good, n = 9; SKP 32150.

Saussurea auriculata (DC.) Sch.Bip., n = 16; SKP 32121. Saussurea costus (Falc.) Lipsch., n = 17; SKP 32104. Angeline M.S. Santos,¹ Felipe Nollet,² Alessandro Rapini,³ Roxana C. Barreto,⁴ Joel M.P. Cordeiro,¹ Erton M. Almeida,¹ José L. Silva,¹ Fabiane R.C. Batista⁵ & Leonardo P. Felix¹

1 Laboratório de Citogenética Vegetal, Departamento de Ciências Biológicas, Centro de Ciências Agrárias, Universidade Federal da Paraíba, Campus II, 58397-000 Areia, Paraíba Brazil

- 2 Laboratório de Sistemática e Florística de Angiospermas, Departamento de Biologia, Universidade Federal Rural de Pernambuco, 52171-900 Recife, Pernambuco, Brazil
- 3 Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas, 44036-900 Feira de Santana, Bahia, Brazil
- 4 Universidade Federal de Pernambuco, Departamento de Botânica, 50670-901 Recife, Pernambuco, Brazil
- 5 Instituto Nacional do Semiárido, 58434700, Campina Grande, Paraíba, Brazil
- * Author for correspondence: angelinemssantos@gmail.com

All materials CHN; vouchers in EAN (Herbarium Prof. Jayme Coelho de Moraes).

Financial support from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), and INSA (Instituto Nacional do Semiárido).

APOCYNACEAE

Allamanda doniana Müll.Arg., 2n = 27; Brazil, Ceará, E.M. Almeida 1137.

- Aspidosperma pyrifolium Mart., 2n = 34; Brazil, Paraíba, A.M.S. Santos 24.
- *Mandevilla bahiensis* (Woodson) M.F.Sales & Kin.-Gouv., 2*n* = 20; Brazil, Bahia, *L.P. Felix 15520*.
- Mandevilla catimbauensis Souza-Silva, Rapini & J.F.Morales, 2n = 26; Brazil, Pernambuco, J.M.P. Cordeiro 379.
- Mandevilla dardanoi M.F.Sales, Kin.-Gouv. & A.O.Simões, 2n = 20; Brazil, Alagoas, J.M.P. Cordeiro 481.
- Mandevilla leptophylla (A.DC.) K.Schum., 2n = 18; Brazil, Ceará, L.P. Felix 15060.
- Mandevilla tenuifolia (J.C.Mikan) Woodson, 2n = 20; Brazil, Ceará, E.M. Almeida 1165.

Marsdenia caatingae Morillo, 2n = 22; Brazil, Paraíba, A.S. Santos 26.

- Marsdenia hilariana E.Fourn., 2n = 22; Brazil, Ceará, L.P. Felix 15063.
- Marsdenia megalantha Goyder & Morillo, 2n = 22; Brazil, Ceará, E.M. Almeida 1169.
- *Peltastes peltatus* (Vell.) Woodson, 2*n* = 18; Brazil, Pernambuco, *L.P. Felix* 13771.
- Petalostelma dardanoi Fontella, 2n = 20; Brazil, Paraíba, J.M.P. Cordeiro 258.
- Tabernaemontana catharinensis A.DC., 2n = 22; Brazil, Ceará, L.P. Felix 14943.
- *Tabernaemontana solanifolia* A.DC., 2n = 22; Brazil, Bahia, *J.M.P. Cordeiro* 654.
- *Temnadenia stellaris* (Lindl.) Miers, 2*n* = 18; Brazil, Paraíba, *J.M.P. Cordeiro 591.*

ARACEAE

Anthurium affine Schott, 2n = 30; Brazil, Paraíba, E.M. Almeida 1069.

COMMELINACEAE

Aneilema brasiliense C.B.Clarke, 2n = 24; Brazil, Paraíba, L.P. Felix 15908.

Commelina diffusa Burm.f., 2n = 30; Brazil, Paraíba, A.S. Santos 32. Commelina diffusa Burm.f., 2n = 60; Brazil, Paraiba, E.M. Almeida 1067.

- *Commelina obliqua* Vahl, 2*n* = 60; Brazil, Pernambuco, *L.P. Felix* 15665.
- *Dichorisandra hexandra* (Aubl.) Standl., 2*n* = 76; Brazil, Pernambuco, *L.P. Felix 15736*.

Tradescantia ambigua Mart., 2*n* = 24; Brazil, Paraíba, *A.S. Santos* 18.

COSTACEAE

Chamaecostus subsessilis (Nees & Mart.) C.D.Specht & D.W.Stev., 2n = 18; Brazil, Paraíba, *E.M. Almeida 1074*.

EUPHORBIACEAE

Cnidoscolus urens (L.) Arthur, 2n = 72; Brazil, Paraíba, E.M. Almeida 1076.

MELASTOMATACEAE

Tibouchina heteromalla (D.Don) Cogn., 2n = 36; Brazil, Paraiba, *E.M. Almeida 1077.*

SOLANACEAE

Solanum polytrichum Moric., 2n = 24; Brazil, Bahia, E.M. Almeida 1409.

VELLOZIACEAE

Vellozia plicata Mart., 2n = 18; Brazil, Paraíba, L.P. Felix 14525.

VITACEAE

Cissus decidua Lombardi, 2*n* = 34; Brazil, Paraíba, *J.M.P. Cordeiro* 260.

Julia V. Shner, Tatiana V. Alexeeva & Michael G. Pimenov*

Botanical Garden, Moscow State University, 119234 Moscow, Russia

* Author for correspondence: mgpimenov@mail.ru

All materials CHN; collector: MP = M.G. Pimenov; vouchers in MW.

This study was supported by the grants 13-04-00648 and 16-04-00525 from the Russian Foundation for Basic Research (RFBR)

UMBELLIFERAE/APIACEAE

Athamanta sicula L., 2*n* = 22; Italy, Sicily, *MP 12-4*, 22 Jul 2012, *MP s.n.*, 13 Jul 2012, *MP s.n.*

Berula erecta (Huds.) Coville, n = 9; Greece, MP 15-30.

Bupleurum glumaceum Sm., n = 8; Greece, MP 15-7.

Chaerophyllum temulum L., n = 7; Greece, MP 15-39.

Conium divaricatum Boiss. & Orph., 2n = 22; Greece, MP 15-4.

Daucus carota subsp. *gummifer* (Syme) Hook.f. = *D. carota* subsp. *hispanicus* (Gouan) Thell.), *n* = 9; France, 03 Jun 2017, *MP s.n.*

Daucus involucratus Sm., 2n = 22; Greece, MP 15-17.

- *Daucus maximus* Desf. (\equiv *D.carota* subsp. *maximus* (Desf.) Ball), 2*n* = 18; Turkey, 7 Aug 2014, *MP s.n.*
- *Eryngium campestre* var. *virens* (Link) Weins, *n* = 14; Greece, *MP* 15-9.
- *Eryngium creticum* Lam., *n* = 7; Greece, *MP* 15-29.
- Ferula tingitana L., 2n = 22; Turkey, MP 14-4.
- *Ferulago humilis* Boiss., 2n = 22; Turkey, *MP 14-8*.
- Ferulago mughlae Peşmen, 2n = 66; Turkey, MP 14-9.
- *Ferulago trojana* Akalın & Pimenov, 2*n* = 22; Turkey, 6 Aug 2014, *MP* & *E. Akalın s.n.*
- Helosciadium nodiflorum (L.) W.D.J.Koch, 2n = 22; Italy, Sicily, MP 12-13.
- Orlaya grandiflora (L.) Hoffm., 2n = 20; Greece, MP 15-25.
- Pastinaca yildizii Dirmenci, n = 11; Turkey, MP & T. Dirmenci 14-10. Pimpinella cretica Poir., n = 9; Greece, MP 15-3; 2n = 18; Greece, 25 Jun 2015, MP s.n.
- *Ridolfia segetum* (Guss.) Moris, 2n = 22; Italy, Sicily, *MP 12-8*.
- Scaligeria moreana Engstrand, 2n = 22; Greece, E.A. Zakharova & U.A. Ukrainskaya 48.
- Scaligeria napiformis (Willd. ex Spreng.) Grande, 2n = 20; Greece, MP 15-20, MP 15-32, E.A. Zakharova & U.A. Ukrainskaya 47.
- Seseli varium Trevir. (= S. pallasii Besser), n = 9; Greece, MP 15-23.
- *Smyrnium olusatrum* L., 2*n* = 22; Greece, 25 Jun 2015, *MP s.n.*; Italy, Sicily, 13 Jul 2012, *MP s.n.*; Turkey, 27 Jul 2014, *MP s.n.*

Smyrnium rotundifolium Mill., 2n = 22; Greece, 23 Jun 2015, *MP s.n.*; Italy, Sicily, *MP 12-15*.

Thapsia garganica L., 2n = 22; Italy, Sicily, MP 12-1.

Tordylium apulum L., 2*n* = 20; Greece, *MP 15-2*.

Torilis triradiata Boiss. & Heldr., 2n = 12; Turkey, MP 14-3.

Mayco Werllen dos Santos Sousa,^{1*} Marisa Toniolo Pozzobon,² José Francisco Montenegro Valls^{2,3} & Regina Célia de Oliveira¹

- 1 Programa de Pós-Graduação em Botânica da Universidade de Brasília, Campus Darcy Ribeiro, Brasília, DF, Brazil, CEP 70910-900
- 2 Embrapa Recursos Genéticos e Biotecnologia, Parque Estação Biológica - PqEB, Brasília, DF, Brazil, CEP 70770-917
- 3 Research Productivity Fellowship/CNPq, Brazil (Proc. 312215/2013-4)
- * Author for correspondence: maycowerllen@gmail.com

All materials CHN.

Financial support from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior/CAPES, Conselho Nacional de Desenvolvimento Científico e Tecnológico/ CNPq (Proc. 312215/2013-4, 562340/2010-6 and 561768/2010-2), and Fundação de Amparo à Pesquisa do Distrito Federal/FAP-DF (Proc. 0193.000.979/2015). Brasília, Brazil.

POACEAE

Paspalum centrale Chase, 2n = 20; Brazil, Acre, *R.C. Oliveira 3405* (UB).

Paspalum convexum Humb. & Bonpl. ex Flüggé, 2n = 32; Brazil, Distrito Federal, J.F.M. Valls & M.W.S. Sousa 16788 (CEN, UB), J.F.M. Valls & M.W.S. Sousa 16789 (CEN, UB); Brazil, Goiás, J.F.M. Valls & al. 16187 (CEN, UB), R.C. Oliveira & M.W.S. Sousa 3005 (UB).
Paspalum cordaense Swallen, 2n = 40; Brazil, Bahia, J.F.M. Valls

- & al. 16694 (CEN, UB); Brazil, Goiás, J.F.M. Valls & al. 16611 (CEN, UB).
- Paspalum macranthecium Parodi, 2n = 20; Brazil, Bahia, J.F.M. Valls & al. 16693 (CEN, UB).
- Paspalum marmoratum Kuhlm., 2n = 18; Brazil, Mato Grosso, C.O. Moura & Y.F.F. Soares 98 (UB).
- Paspalum melanospermum Desv. ex Poir., 2n = 20; Brazil, Amazonas, R.C. Oliveira & J.M. Mendoza 3025 (UB); Brazil, Bahia, R.C. Oliveira 3362 (UB); Brazil, Goiás, Valls & al. 16671 (CEN, UB); 2n = 40, Brazil, Amazonas, R.C. Oliveira & J.M. Mendoza 3010 (UB); Brazil, Distrito Federal, R.C. Oliveira & al. 3443 (UB); Brazil, Goiás, J.F.M. Valls. & al. 16188 (CEN, UB), J.F.M. Valls. & al. 16195 (CEN, UB), J.F.M. Valls. & al. 16622 (CEN, UB); R.C. Oliveira & M.W.S. Sousa 3003 (UB); Brazil, Mato Grosso, C.O. Moura & Y.F.F. Soares 90 (UB); Brazil, Tocantins, M.W.S. Sousa 87 (UB).
- Paspalum plicatulum Michx., 2n = 40; Brazil, Distrito Federal, A.P.M.L. Gouveia 232 (UB), R.C. Oliveira & M.W.S. Sousa 3423 (UB).
- Paspalum riparium Nees, 2n = 40; Brazil, Amazonas, R.C. Oliveira & J.M. Mendoza 3008 (UB), R.C. Oliveira & J.M. Mendoza 3016 (UB).
 Paspalum rojasii Hack., 2n = 40; Brazil, Distrito Federal, J.F.M. Valls & M.W.S. Sousa 16777 (CEN, UB).

Saeedeh Sadat Mirzadeh Vaghefi* & Adel Jalili

Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), P.O. Box 13185-116, Tehran, Iran Author for correspondence: Mirzadeh@rifr-ac.ir

The study was financially supported by Research Institute Forests and Rangeland of Iran (grant no. 12-09-09-7701-89001). The authors are grateful to several collectors for material donation.

All materials CHN; collectors: A.A. = Ardeshir Ashrafi, J.M. = Javad Mohebi, S.M. = Saeedeh Mirzadeh; vouchers in PUN.

APIACEAE

Ferulago angulata (Schltdl.) Boiss., 2n = 22; Iran, A.A. 106062.

ASTERACEAE

Achillea talagonica Boiss., 2n = 18; Iran, J.M. 106054. Anthemis tinctoria L., 2n = 18; Iran, A.A. 106056. Tanacetum abrotanifolium (L.) Druce, 2n = 36; Iran, J.M. 106059. Tanacetum pinnatum Boiss., 2n = 18; Iran, J.M. 106057. Varthemia persica DC., 2n = 16; Iran, J.M. 106058.

CAPRIFOLIACEAE

Centranthus longiflorus Steven, 2n = 32; Iran, J.M. 106047.

CARYOPHYLLACEAE

Gypsophila bicolor Grossh., 2n = 34; Iran, A.A. 106063.

ELAEAGNACEAE

Hippophae rhamnoides L., 2n = 24; Iran, S.M. 105207.

LAMIACEAE

Lallemantia canescens Fisch. & C.A.Mey., 2n = 14; Iran, A.A. 106048. Phlomis olivieri Benth., 2n = 20; Iran, J.M. 106050. Salvia hypoleuca Benth., 2n = 22; Iran, S.M. 106052. Teucrium hircanicum L., 2n = 32; Iran, J.M. 106051. Thymus persicus (Ronniger ex Rech.f.) Jalas, 2n = 30; Iran, J.M. 106049.

MALVACEAE

Sida rhombifolia L., 2*n* = 14; Iran, *J.M.* 106061.

PAPAVERACEAE

Papaver bracteatum Lindl., 2*n* = 14; Iran, *J.M.* 106060.

PAPILIONACEAE/FABACEAE

Astragalus mesoleios Boiss. & Hohen., 2n = 16; Iran, A.A. 106053. Oreophysa triphylla (Bunge ex Boiss.) Bornm., 2n = 16; Iran, S.M. 106055.

IAPT CHROMOSOME DATA

IAPT chromosome data 28 [extended online version]

Edited by Karol Marhold & Jaromír Kučera

Julio Rubén Daviña & Ana Isabel Honfi*

Programa de Estudios Florísticos y Genética Vegetal, Instituto de Biología Subtropical CONICET-Universidad Nacional de Misiones, nodo Posadas, Rivadavia 2370, 3300 Posadas, Argentina * Author for correspondence: ahonfi@gmail.com

Methods are described in Daviña & Fernández (1989) and Daviña (2001).

- * First chromosome count for the species.
- First gametic chromosome count for the species.
- ▼ New cytotype for the species.
- First meiotic behaviour report for the species.

This study was supported by Agencia Nacional de Promoción Científica y Técnica (ANPCyT) grant nos. PICT-2014-2218 and PICT-2016-1637, and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

AMARYLLIDACEAE

Amaryllidaceae family comprises 860 species and 59 genera with mainly tropical distribution (Meerow & Snijman, 1998). The Hippeastreae Herb. ex Sweet tribe has a major center of diversification in central Chile and western Andean Argentina and a second center in eastern Brazil and northeastern Argentina (Meerow & Snijman, 1998; Arroyo-Leuenberger & Dutilh, 2008). In this opportunity, species belonging to *Habranthus* Herb. from Argentina were cytogenetically studied.

Habranthus barrosianus Hunz. & Di Fulvio

2n = 2x = 18, CHN. Argentina, Buenos Aires Province, Tandil Department, South City Park, 37°20'S, 59°08'W, 27 Jan 1987, *Leuenberger & Arroyo 3772* (B) [Fig. 1A].

The somatic chromosome number of this accession agrees and confirms the findings made by Naranjo (1969, 1974).

Habranthus brachyandrus (Baker) Sealy

• n = 12, 2n = 4x = 24, CHN. Argentina, Corrientes Province, General Paz Department, National Route nº 12, Km 1099, 27°28′S, 58°47′W, 17 Dec 1997, *Daviña 416* (B, MNES) [Fig. 1E].

This chromosome number is in agreement with a previous report (Flory & Flagg, 1958). Chromosomes behave regularly at meiosis, with 12 II at metaphase I (Fig. 2A). Sometimes, in pollen mother cells (PMCs) at anaphase I and telophase I, chromosomes bridges and few laggards were observed (Fig. 2B, C). The chromosome pairing suggest allotetraploidy for the taxon.



Fig. 1. Mitotic metaphase. **A**, *Habranthus barrosianus*, 2n = 18; **B**, *H*. *gracilifolius*, 2*n* = 24; **C**, *H*. tubispathus, 2n = 24; **D**, *H*. tubispathus, 2n = 36; **E**, *H*. *brachyandrus*, 2n = 24, arrowhead points out to chromosome carrying satellite at the short arm; **F**, *H*. coeruleus, 2n = 30, arrowhead points out to chromosome carrying satellite at the long arm; **G**, *H*. *chacoensis*, 2n = 12, arrowhead points out to chromosome carrying satellite at the short arm; **H**, *H*. pedunculosus, 2n = 14. Scale bars = $5 \mu m$.



Fig. 2. Meiotic behaviour: **A**, *Habranthus brachyandrus*, PMC at metaphase I with 12 bivalents; **B**, *H. brachyandrus*, anaphase I with chromosome bridges without fragments suggesting translocation rearrangements; **C**, *H. brachyandrus*, PMC at telophase I with lagard chromosomes; **D**, *H. chacoensis*, PMC at metaphase I with 6 bivalents. — Scale bars = 5 μ m.

Habranthus chacoensis Ravenna

*•• n = 6, 2n = 2x = 12, CHN. Argentina, Chaco Province, Iro. de Mayo Department, Paraje tres Horquetas, 27°27′S, 59°00′W, 4 Mar 1996, *Daviña & Lavia 342* (B, CTES, MNES) [Fig. 1G].

Regularly, chromosomes associate in 6 II at diakinesis and metaphase I (Fig. 2D) in all analyzed PMCs of this diploid species.

Habranthus coeruleus (Griseb.) Traub

* 2n = 2x = 30, CHN. Argentina, Misiones Province, Capital Department, Parada Leis, 27°36'S, 55°52'W, 16 Feb 1999, *Honfi & Daviña 985* (MNES) [Fig. 1F].

Habranthus gracilifolius Herb.

2n = 4x = 24, CHN. Argentina, Buenos Aires Province, Tandil Department, South City Park, 37°20'S, 59°08'W, 27 Jan 1987, *Leuenberger & Arroyo 3774* (B) [Fig. 1B].

This is a species with two ploidy levels, diploids (2n = 12; Naranjo, 1974) and tetraploids as found here, in agreement with a previous report of Flory & Flagg (1958).

Habranthus pedunculosus Herb.

2n = 2x = 14, CHN. Argentina, Chaco Province, 1º de Mayo Department, Colonia Benítez, 27°27′S, 59°00′W, 21 Feb 1996, *Daviña, Dematteis, Lavia & López 334* (B, MNES) [Fig. 1H].

We confirm the chromosome number and ploidy level of the species, previously reported by Naranjo (1974) and Flory & Flagg (1958).

Habranthus tubispathus (L'Hér.) Traub

2n = 4x = 24, CHN. Argentina, Corrientes Province, Capital Department, Molina Punta, 27°27'S, 58°47'W, 22 Feb 1996, *Daviña, Dematteis & Schinini 329* (B, CTES, MNES) [Fig. 1C].

Our count confirms the tetraploid level for the species in agreement with Flory (1948) and Flory & Flagg (1958).

▼ 2n = 6x = 36, CHN. Argentina, Buenos Aires Province, Tandil Department, South City Park, 37°20'S, 59°08'W, 27 Jan 1987, *Leuenberger & Arroyo 3771* (B) [Fig. 1D].

For the first time, we report a new hexaploid cytotype for *H. tubi-sphathus* from Argentina.

LITERATURE CITED

- Arroyo-Leuenberger, S.C. & Dutilh, J. 2008. Amaryllidaceae. Pp. 203–226 in: Zuloaga, F.O., Morrone, O. & Belgrano. M.J. (eds.), Catálogo de las plantas vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay), vol. 1. Monographs in Systematic Botany from the Missouri Botanical Garden 107. St. Louis: Missouri Botanical Garden Press.
- Daviña, J.R. 2001. Estudios citogenéticos en algunos géneros Argentinos de Amaryllidaceae. Tesis doctoral, Universidad Nacional de Córdoba, Argentina.
- Daviña, J.R. & Fernández, A. 1989. Karyotype and meiotic behavior in Zephyranthes (Amaryllidaceae) from South America. Cytologia 54: 269–274. https://doi.org/10.1508/cytologia.54.269
- Flory, W.S. 1948. Chromosome studies, and their bearing on phylogeny, in the Amaryllidaceae. I *Habranthus. Amer. J. Bot.* 35: 791–792.
- Flory, W.S. & Flagg, R.O. 1958. A cytological study of the genus Habranthus. Nucleus (Calcutta) 1: 267–280.
- Meerow, A. & Snijman, D.A. 1998. Amaryllidaceae. Pp. 83–110 in: Kubitzki, K. (ed.), *The families and genera of vascular plants*, vol. 3, *Flowering plants: Monocotyledons; Lilianae (except Orchidaceae)*. Berlin: Springer.
- Naranjo, C.A. 1969. Cariotipo de nueve especies Argentina de Rhodophiala, Hippeastrum, Zephyranthes y Habranthus (Amaryllidaceae). Kurtziana 5: 67–87.
- Naranjo, C.A. 1974. Karyotypes of four Argentine species of Habranthus and Zephyranthes (Amaryllidaceae). Phyton (Buenos Aires) 32: 61–71.

Luis Delgado,^{1*} Blanca M. Rojas-Andrés,^{1,2,3} Noemí López-González,^{1,2} Nélida Padilla-García^{1,2} & M. Montserrat Martínez-Ortega^{1,2}

- 1 Departamento de Botánica y Fisiología Vegetal, University of Salamanca, 37007 Salamanca, Spain
- 2 Biobanco de ADN Vegetal, University of Salamanca, Edificio Multiusos I+D+i, Calle Espejo s/n, 37007 Salamanca, Spain
- 3 Present address: Department of Molecular Evolution and Plant Systematics & Herbarium (LZ), Institute of Biology, Leipzig University, Leipzig, Germany
- * Author for correspondence: ldelsan@usal.es

▲ First chromosome count for the species.

This work has been supported by the Spanish Ministerio de Economía y Competitividad (projects CGL2009-07555, CGL2012-32574, *Flora iberica* VIII [CGL2008-02982-C03-02/CLI], *Flora iberica* IX [CGL2011-28613-C03-03], *Flora iberica* X [CGL2014-52787-C3-2-P]); the Spanish Ministerio de Ciencia e Innovación (Ph.D. grants to BR and NLG), and the University of Salamanca (Ph.D. grant to NPG financed jointly by Banco Santander). The authors are very grateful to all colleagues who helped with plant location and sampling.

CHN

Material CHN, collected in Bosnia and Herzegovina, Bulgaria, Montenegro, Macedonia (F.Y.R.O.M.) and Switzerland. All cytological investigations have been carried out on anthers and gynoecia. Material was fixed in 3:1 absolute ethanol-glacial acetic acid and stained in 2% acetic orcein (Cour, 1945).

FCM

Material FCM, collected in Austria, Czech Republic, Croatia, France, Morocco, Romania, Serbia, Slovakia, Spain, Switzerland and Turkey. DNA ploidy levels were estimated from silica-gel-dried leaves. Samples were prepared according to the procedure described in Rojas-Andrés & al. (2015). Leaf samples of *Pisum sativum* L. 'Ctirad' (2C DNA = 9.09 pg; Dolezel & al., 1998), *Solanum pseudocapsicum* L. (2C DNA = 2.589 pg; Temsch & al., 2010) and *Zea mays* L. 'CE-777' (2C DNA = 5.43 pg; Lysak & Dolezel, 1998) were used as internal reference standards according to sample C-value and standard availability. Results on DNA ploidy level were acquired using a CyFlow SL system (Partec, Münster, Germany) equipped with a blue 488 nm solid state laser. For each individual, the ratio of the G_0/G_1 peak positions of samples and internal standards were recorded. DNA ploidy level was estimated based on the genome size value found and the available chromosome number for each particular taxon. In many cases FCM measurements were directly compared with CHN corresponding to the same samples. The coefficient of variation (CV) was calculated for each sample and the standard used. All data were suitable for DNA ploidy level estimation because CV values did not exceed the 10% threshold.

TAXONOMIC TREATMENT

Material determination and nomenclatural treatment follow Rojas-Andrés & Martínez-Ortega (2016) and Rojas-Andrés & al. (2016), respectively, with the modifications proposed by Padilla-García & al. (2018).

PLANTAGINACEAE

Veronica angustifolia (Vahl) Bernh.

2n = 64, CHN. Switzerland, Valais, on the ascent to Tanay lake, 46°20'36.0"N, 06°50'33.4"E, 1418 m, 11 Jul 2011, X. Giráldez, M.M. Martínez-Ortega 6038 & B. Rojas-Andrés (SALA 149416) [Fig. 31].

 $2n \sim 8x \sim 64$, FCM. France, Departement of Hautes-Alpes, Gap. Chorges, near Gap. On the ascent to Monte Chabrières from Les Andrieux, at the margins of the road, 44°34'35.3"N, 06°16'35.3"E,



Fig. 3. Mitotic metaphases: **A–D**, *Veronica austriaca* subsp. *jacquinii*: **A**, 2n = 32 (SALA 157023); **B**, 2n = 32 (SALA 157024); **C**, 2n = 48 (SALA 157021); **D**, 2n = 48 (SALA 149376); **E & F**, *V. dalmatica*: **E**, 2n = 16 (SALA 157018); **F**, 2n = 16 (SALA 157030); **G & H**, *V. kindlii*, 2n = 16 (SALA 157011); **I**, *V. angustifolia*, 2n = 64 (SALA 149416). — Scale bars, 10 μm.

1393 m, 10 Jul 2011, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 167-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 167-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 167-3 (SALA 149398); Switzerland, Valais, on the ascent to Tanay lake, 46°20'36.0"N, 06°50'33.4"E, 1418 m, 11 Jul 2011, X. Giráldez, M.M. Martínez-Ortega 6038-1 & B. Rojas-Andrés, X. Giráldez, M.M. Martínez-Ortega 6038-2 & B. Rojas-Andrés, X. Giráldez, M.M. Martínez-Ortega 6038-3 & B. Rojas-Andrés (SALA 149416).

Based on AFLP data, this species name has been recently resurrected to name plants that were previously included under the variation of *V. teucrium* L. (Padilla-García & al., 2018). Simonet (1934) and Brandt (1952) counted the chromosomes of plants from this group from France and Switzerland, but we have not been able to revise the corresponding herbarium vouchers and do therefore not know for sure whether they correspond to *V. angustifolia* or *V. teucrium*.

Veronica austriaca subsp. jacquinii (Baumg.) Watzl

2n = 32, CHN. Bosnia and Herzegovina, Trebinje, between Gacko and Tjentiste, subalpine meadows on limestone soils, 43°11′04.92″N, 18°33′56.808″E, 1076 m, 10 Jun 2015, *M.M. Martínez-Ortega 6122, X. Giráldez, N. Padilla-García & N. López-González* (SALA 157023) [Fig. 3A], *M.M. Martínez-Ortega 6123, X. Giráldez, N. Padilla-García & N. López-González* (SALA 157024) [Fig. 3B].

2n = 48, CHN. Bosnia and Herzegovina, Trebinje, between Gacko and Tjentiste, subalpine meadows on limestone soils, 43°11′04.92″N, 18°33′56.808″E, 1076 m, 10 Jun 2015, *M.M. Martínez-Ortega 6120, X. Giráldez, N. Padilla-García & N. López-González* (SALA 157021) [Fig. 3C]; Bulgaria, Pazardzhik, between Batak and Beglika, perennial grassland, 41°50′27″N, 24°08′48″E, 1522 m, 17 Jun 2009, *X. Giráldez, M.M. Martínez-Ortega, B. Rojas-Andrés 21* & *M. Santos Vicente* (SALA 149368); Bulgaria, Pernik, between Staychovtsi and Dolna Melna, 42°41′04″N, 22°31′57″E, 950 m, 14 Jun 2009, *X. Giráldez, M.M. Martínez-Ortega 4577, B. Rojas-Andrés & M. Santos Vicente* (SALA 149376) [Fig. 3D].

 $2n \sim 6x \sim 48$, FCM. Croatia, Licausko Polje, meadows on limestone soils, 704 m, 17 Jun 2001, *M.M. Martínez-Ortega 1404-1 & A. Tribsch, M.M. Martínez-Ortega 1404-2 & A. Tribsch* (SALA 124609); Croatia, Primorje-Gorski Kotar, Velika Kapela, Gornje Jelenje pass, at the crossroad towards Crikvanica, meadows on limestone soils, 45°21′58″N, 14°37′09″E, 880 m, 16 Jun 2001, *M.M. Martínez-Ortega 1391-1 & A. Tribsch*, *M.M. Martínez-Ortega 1391-2 & A. Tribsch* (SALA 124604).

Numerous chromosome counts are known for this taxon, 2n = 32, 48, 64 (cf. Scheerer, 1937; Moore, 1973, 1977; Peev, 1978; Goldblatt & Johnson, 1979+; Goldblatt, 1981, 1988; Májovský & Murín, 1987), but 2n = 48 is the most frequent one. The tetraploid cytotype 2n = 32 is rare and has been reported only once from Bulgaria (Peev, 1972). The population studied here from Bosnia and Hercegovina contains tetraploid (2n = 32; SALA 157023 and SALA 157024) and hexaploid individuals (2n = 48; SALA 157021). The morphology of the tetraploid individuals is also close to that of *Veronica orbiculata* A.Kern.

▲ *Veronica dalmatica* N.Pad.Gar., Rojas-Andrés, López-González & M.M.Mart.Ort.

2n = 16, CHN. Montenegro, Kotor, Lovcen, shrub clearings on limestones, $42^{\circ}25'04.9''N$, $18^{\circ}47'39''E$, 904 m, 9 Jun 2015, *M.M. Martínez-Ortega, X. Giráldez, N. Padilla-García & N. López-González 137* (SALA 157018) [Fig. 3E]; Montenegro, Žabljak, Meždo, dry meadows on limestone soils with *Juniperus*, $43^{\circ}09'49.824''N$, $19^{\circ}08'56.688''E$, 1390 m, 12 Jun 2015, *M.M. Martínez-Ortega*, X. Giráldez, N. Padilla-García & N. López-González 139 (SALA 157030) [Fig. 3F].

It has not been possible to revise herbarium material corresponding to a previous chromosome count (2n = 16) from an Albanian population (Baltisberger, 1988) identified as *V. jacquinii*. Padilla-García & al. (2018) have recently shown that the diploid representatives traditionally assigned to *V. austriaca* subsp. *jacquinii* represent a distinct species named *V. dalmatica*. Here, material clearly assigned to this species collected in Montenegro is investigated for the first time.

▲ *Veronica kindlii* Adam.

2*n* = 16, CHN. Macedonia (F.Y.R.O.M.), Bitola, on the ascent to Pelister summit, subalpine meadows on granite soils, 40°59′40.416″N, 21°10′42.492″E, 2355 m, 8 Jul 2014, *X. Giráldez, M.M. Martínez-Ortega 6090, B. Rojas-Andrés & N. López-González* (SALA 157011) [Fig. 3G, H].

The name *V. kindlii* has been recently resurrected to designate those populations from the Balkan Peninsula previously identified as *V. orsiniana* (Rojas-Andrés & al., 2015). A previous chromosome count for *V. kindlii* was carried out on material collected in the Macedonian locality of Barbaros (Makedonski Brod) (Sopova & al., 1983 in Goldblatt, 1988). However, AFLP data showed that the latter population corresponds to the closely related species *V. linearis* (Padilla-García & al., 2018). Thus, the chromosome count published here would represent the first one for *V. kindlii*.

Veronica orsiniana Ten.

 $2n \sim 2x \sim 16$, FCM. France, Department of Lozère, Cévennes, 5 kilometers from Aven Armand cave, at the crossroad towards Masde-la-Font, montane meadows with Buxus on acid soil, 44°12'22"N, 03°23'31"E, 945 m, 8 Jun 2012, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 203-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 203-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 203-3 (SALA 149303); France, Department of Lozère, Cévennes, Aven Armand cave, meadows on limestone soils, 44°13'29"N, 03°21'34"E, 1006 m, 8 Jun 2012, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 207-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 207-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 207-3 (SALA 149305); France, Department of Var, on the ascent to Sainte Baume massif, Braque pathway, meadows on limestone soils, 43°19'21"N, 05°42'09"E, 693 m, 9 Jun 2012, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 210-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 210-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 210-3 (SALA 149308); Spain, Barcelona, Collsuspina, between Tona and Moyá, grazed pastures near the road, grassy slopes, 41°49'39"N, 02°10'45"E, 916 m, 14 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 235-1, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 235-2, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 235-3, A. Abad & N. López-González (SALA 155093), M.M. Martínez-Ortega, B. Rojas-Andrés 235-22*, A. Abad & N. López-González (SALA 155067); Spain, Barcelona, Espunyola, towards Montclar village, clearings in Quercus ilex and Q. faginea forest on limestone soils, 42°01'15"N, 01°46'11"E, 757 m, 16 Jun 2013, M.M. Martínez-Ortega 6064-1, B. Rojas-Andrés, X. Giráldez & N. López-González, M.M. Martínez-Ortega 6064-2, B. Rojas-Andrés, X. Giráldez & N. López-González, M.M. Martínez-Ortega 6064-3, B. Rojas-Andrés, X. Giráldez & N. López-González (SALA 155119); Spain, Barcelona, Montserrat, on the road to the monastery El Bruc, to Manresa, Can Macana, meadows on limestone soils behind the farmhouse,

41°36'36"N, 01°46'01"E, 724 m, 13 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 8-1, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 8-2, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 8-3 (SALA 155097); Spain, Barcelona, Tona, on the road to Castell and the Lourdes chapel, dry grassy meadows at the side of the pathway, marly limestones, 41°51'16"N, 02°13'18"E, 661 m, 14 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 234-1, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 234-2, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 234-3, A. Abad & N. López-González (SALA 155094); Spain, Gerona, from Viladray to Sta. Fe del Montseny, at the crossroad to Mas el Martí, meadows near the road, 41°50'24"N, 02°24'52"E, 966 m, 16 Jun 2013, M.M. Martínez-Ortega 6063-1, B. Rojas-Andrés, A. Abad & N. López-González, M.M. Martínez-Ortega 6063-2, B. Rojas-Andrés, A. Abad & N. López-González, M.M. Martínez-Ortega 6063-3, B. Rojas-Andrés, A. Abad & N. López-González (SALA 155123); Spain, Huesca, Sta. Cruz de la Serós, San Juan de la Peña, meadows at the picnic area, near the interpretation center and the new Monastery, 42°30'26"N, 00°39'55"W, 1214 m, 20 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 242-1, X. Giráldez & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 242-2, X. Giráldez & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 242-3, X. Giráldez & N. López-González (SALA 155109); Spain, Huesca, pathway between Binacua and the crossroad to the road N-240, meadows on limestone soils and slopes with Buxus, 42°33'09"N, 00°41'47"W, 718 m, 20 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 243-1, X. Giráldez & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 243-2, X. Giráldez & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 243-3, X. Giráldez & N. López-González (SALA 155110).

Veronica prostrata L.

 $2n \sim 2x \sim 16$, FCM. Austria, Burgenland, Leithagebirge, between Donnerskirchen and Franz Josef-Warte, dry grasslands, Pannonian vegetation, 47°55'59"N, 16°40'03"E, 250 m, 28 May 2001, M.M. Martínez-Ortega 1055-1, M.M. Martínez-Ortega 1055-2, M.M. Martínez-Ortega 1055-3 (SALA 124616); Austria, Niederösterreich, Falkenstein, near the castle ruins, 48°43'28"N, 16°34'45"E, 390 m, 11 Jun 2001, M.M. Martínez-Ortega 1337-1, M.M. Martínez-Ortega 1337-2, M.M. Martínez-Ortega 1337-3 (SALA 124615); Czech Republic, South Moravia, Břeclav District, Mikulov, near the Austrian border, Šibeničník Nature reserve, dry meadows, Pannonian vegetation, 48°47'25"N, 16°37'56"E, 236 m, 23 Jul 2011, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 184-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 184-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 184-3 (SALA 149310); Czech Republic, South Moravia, Břeclav District, between Mikulov and Klentnice, dry meadows on limestone soils, in an environment of Pannonian vegetation, 48°49'33"N, 16°38'26"E, 345 m, 24 Jul 2001, M.M. Martínez-Ortega 1445-1 (SALA 124617); France, Departement of Hautes-Alpes, Gap, Col du Noyer, between Le Noyer and L'Enclus, subalpine meadows on limestone soils, 44°41'35"N, 05°59'13"E, 1671 m, 14 Jun 2012, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 219-1, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 219-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés 219-3 (SALA 149313); Romania, Cluj, Fanatele Clujului, 8 km from Cluj-Napoca, meadows on limestone soils, 46°51'08"N, 23°37'01"E, 550 m, 30 Jun 2000, X. Giráldez, M.M. Martínez-Ortega 904-1 & J.A. Sánchez Agudo, X. Giráldez, M.M. Martínez-Ortega 904-2 & J.A. Sánchez Agudo (SALA 124620); Serbia, Južnobanatski okrug

(South Banat), Devojački Bunar, Vladimirovac, dry sandy meadows, 45°00'58"N, 20°57'17"E, 163 m, 27 Jul 2010, S. Andrés 428-I, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés, S. Andrés 428-2, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés, S. Andrés 428-3, X. Giráldez, M.M. Martínez-Ortega & B. Rojas-Andrés (SALA 149319).

Ploidy level estimations for the species in Romania are here given for the first time. No previous chromosome count or ploidy level estimation was available for *V. prostata* in this country (cf. Bolkhovskikh & al., 1969; Goldblatt & Johnson, 1979+; Albach & al., 2008).

Veronica rosea Desf.

 $2n \sim 2x \sim 16$, FCM. Morocco, Ifrane, Meknès-Tafilalet, Jebel Hebri, grazed pastures with *Thymelaea*, 33°21'13"N, 05°08'43"W, 1931 m, 18 Jul 2013, *D. Pinto*, *N. López-González 52-1 & V. Lucía García*, *D. Pinto*, *N. López-González 52-2 & V. Lucía García*, *D. Pinto*, *N. López-González 52-3 & V. Lucía García* (SALA 155072); Morocco, Ifrane, in front of Aquelmame de Si-Ali, road between Azrou and Midelt (between Timahdite and Ait-Oufella), scrubland in clearings of Juniperus thurifera subsp. *africana*, on limestone soils, 33°04'50.988"N, 05°01'13.094"W, 2168 m, 1 May 2013, J. Peñas de Giles 1, J. Peñas de Giles 2, J. Peñas de Giles 3 (GDA 59936).

Veronica sennenii (Pau) M.M.Mart.Ort. & E.Rico

 $2n \sim 8x \sim 64$, FCM. Spain, León, Lois, near the road towards Liegos/Anciles, meadows on limestone soils, $42^{\circ}59'04''N$, $05^{\circ}07'05''W$, 1278 m, 2 Jul 2013, *M.M. Martínez-Ortega, B. Rojas-Andrés 248-1 & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 248-2* & *N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 248-3* & *N. López-González* (SALA 155081).

Veronica tenuifolia subsp. javalambrensis (Pau) Molero & J.Pujadas

 $2n \sim 2x \sim 16$, FCM. Spain, Burgos, Amaya, on the ascent to Peña Amaya, meadows on limestone soils, 42°39'02"N, 04°10'41"W, 1167 m, 5 Jul 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 250-1 & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 250-2 & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 250-3 & N. López-González (SALA 155086); Spain, Burgos, Ciruelos de Cervera, Briongos de Cervera, Raposa pathway, near the road BU-914, clearings of Juniperus sabina, limestone soil, 41°54'50"N, 03°29'48"W, 1085 m, 2 Jun 2013, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1278-1, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1278-2, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1278-3 (SALA 150521); Spain, Burgos, Espinosa de Cervera, near the old football ground, short pastures in clearings of Juniperus sabina close to the pathway, 41°54'01.3"N, 03°27'53.9"W, 1074 m, 2 Jun 2013, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1280-1, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1280-2, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1280-3 (SALA 150519); Spain, Santander, road between Espinama and the Aliva refuge, subalpine meadows on limestone soils, 43°09'55.86"N, 04°46'25.92"W, 1565 m, 3 Jul 2013, M.M. Martínez-Ortega 6074-1, B. Rojas-Andrés & N. López-González, M.M. Martínez-Ortega 6074-2, B. Rojas-Andrés & N. López-González, M.M. Martínez-Ortega 6074-3, B. Rojas-Andrés & N. López-González (SALA 155084); Spain, Segovia, Moral de Hornuez, Sabinar de Hornuez (picnic area near the chapel), dry and short meadows on limestone soils, 41°29'00"N, 03°37'32.8"W, 25 May 2013, M.M. Martínez-Ortega & N. López-González 3-1, M.M. Martínez-Ortega & N. López-González 3-2, M.M. Martínez-Ortega

& N. López-González 3-3 (SALA 155104); Spain, Soria, El Burgo de Osma, Torralba del Burgo, Arroyo de los Barranquillos, in environments of Juniperus thurifera, on clay and gravel soils, 41°37'44.1"N, 02°55'00.2"W, 974 m, 8 Jun 2013, D. Pinto 1315-1, D. Pinto 1315-2, D. Pinto 1315-3 (SALA 150484); Spain, Soria, Herrera de Soria, Camino del Oropar, Barranco de la Covatilla, streambed in Juniperus thurifera forest, 41°46'16.8"N, 03°02'05.0"W, 1093 m, 8 Jun 2013, D. Pinto 1311-1, D. Pinto1311-2, D. Pinto 1311-3 (SALA 150488); Spain, Soria, Langa de Duero, Alcozar, near Cerro Hestilla, short meadows and scrublands of Thymus sp. on limestone soils, 41°37'29.9"N, 03°19'28.1"W, 904 m, 2 Jun 2013, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1287-1, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1287-2, D. Gutiérrez-Larruscain, N. López-González & D. Pinto 1287-3 (SALA 150512); Spain, Soria, Recuerda, on the ascent to La Muela from the water source el Cepo, Juniperus thurifera forest on limestones, 41°28'01.0"N, 02°58'54.1"W, 950 m, 8 Jun 2013, D. Pinto 1307-1, D. Pinto 1307-2, D. Pinto 1307-3 (SALA 150492); Spain, Soria, Villaciervos, El Santo, streambed on Genista scorpius scrublands in environment of Juniperus sabina, 41°46'08.1"N, 02°38'54.6"W, 1228 m, 8 Jun 2013, D. Pinto 1322-1, D. Pinto 1322-2, D. Pinto 1322-3 (SALA 150477); Spain, Zamora, Vezdemarbán, hills near the village on limestone soils, table hills, 41°39'03.8"N, 05°21'50.0"W, 793 m, 19 Jun 2012, S. Barrios, N. López-González, M.M. Martínez-Ortega & B. Rojas-Andrés 221-1, S. Barrios, N. López-González, M.M. Martínez-Ortega & B. Rojas-Andrés 221-2, S. Barrios, N. López-González, M.M. Martínez-Ortega & B. Rojas-Andrés 221-3 (SALA 149327).

Veronica tenuifolia Asso subsp. tenuifolia

 $2n \sim 2x \sim 16$, FCM. Spain, Barcelona, Collsuspina, meadows on limestones near the village, on the left side along the pathway to Sta. Coloma de Castellterçol (GR177), 41°49'24"N, 02°10'36.24"E, 905 m, 14 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés 237-1, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 237-2, A. Abad & N. López-González, M.M. Martínez-Ortega, B. Rojas-Andrés 237-3, A. Abad & N. López-González (SALA 155065); Spain, Barcelona, on the ascent to Sta. Perpetua chapel, near the road, 41°59′54.9″N, 02°12′09.9″E, 663 m, 15 Jun 2013, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 16-1, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 16-2, M.M. Martínez-Ortega, B. Rojas-Andrés, A. Abad & N. López-González 16-3 (SALA 155125); Spain, Teruel, Bordón, on the road to Calanda, before the crossroad to Luco de Bordón and Bordón river, marly slopes, 40°41'36.6"N, 00°19'09.5"W, 769 m, 10 Jun 2013, M.M. Martínez-Ortega 6059-1, B. Rojas-Andrés, X. Giraldez & N. López-González, M.M. Martínez-Ortega 6059-2, B. Rojas-Andrés, X. Giraldez & N. López-González, M.M. Martínez-Ortega 6059-3, B. Rojas-Andrés, X. Giraldez & N. López-González (SALA 155099).

Veronica teucrium L.

2n ~ 8x ~ 64, FCM. Slovakia, Košice, distr. Košice-okolie, Dvorníky-Včeláre, Zádiel, dry meadows on limestones, 48°36'46"N, 20°50'32"E, 300 m, 3 Jul 2002, *M.M. Martínez-Ortega 1551-1,* X. Giráldez & Muñoz Centeno, M.M. Martínez-Ortega 1551-2, X. Giráldez & Muñoz Centeno, M.M. Martínez-Ortega 1551-3, X. Giráldez & Muñoz Centeno (SALA 124598).

Veronica thracica Velen.

 $2n \sim 2x \sim 16$, FCM. Turkey, Kirklareli, Dereköy, road towards Geçitagzi, mixed forest and meadows on limestone soils, 41°56'15"N, 27°21'11"E, 462 m, 23 Jun 2009, X. Giráldez, M.M. Martínez-Ortega, B. Rojas-Andrés 37-1 & M. Santos Vicente, X. Giráldez, M.M. Martínez-Ortega, B. Rojas-Andrés 37-2 & M. Santos Vicente (SALA 149289).

Veronica thracica was previously included within the variation of *V. crinita* Velen. (Peev, 1972; Rojas-Andrés & al., 2015) but has been recently combined at the specific rank based on AFLP data (Padilla-García & al., 2018). Previous chromosome counts (Peev, 1975, 1976) and ploidy level estimations (Padilla-García & al., 2018) concern plants distributed in Bulgaria. Consequently, these are the first ploidy level estimations for this species from Turkey.

LITERATURE CITED

- Albach, D.C., Martínez-Ortega, M.M., Delgado, L., Weiss-Schneeweiss, H., Özgökce, F. & Fischer, M.A. 2008. Chromosome numbers in Veroniceae (Plantaginaceae): Review and several new counts. Ann. Missouri Bot. Gard. 95: 543–566. https://doi.org/10.3417/2006094
- Baltisberger, M. 1988. Chromosomenzahlen einiger Pflanzen aus Albanien. II = Chromosome numbers of some plants from Albania. II. Ber. Geobot. Inst. E. T. H. Stiftung Rübel 54: 42–50.
- Bolkhovskikh, Z., Grif, V., Matvejeva, T. & Zakharyeva, O. 1969. Chromosome numbers of flowering plants. Leningrad: Nauka.
- Brandt, J.P. 1952. Contribution à la cytologie du genre Veronica. Bull. Soc. Neuchâtel. Sci. Nat. 75: 179–188.
- Cour, L.F. 1945. Smear and squash techniques in plant cytology. *Lab. Pract.* 3: 326–330.
- Dolezel, J., Greilhuber, J., Lucrettiii, S., Meister, A., Lysak, M.A., Nardi, L. & Obermayer, R. 1998. Plant genome size estimation by flow cytometry: Inter-laboratory comparison. *Ann. Bot. (Oxford)* 82: 17–26. https://doi.org/10.1093/oxfordjournals.aob.a010312
- Goldblatt, P. 1981. Index to plant chromosome numbers 1975–1978. Monogr. Syst. Bot. Missouri Bot. Gard. 5: 1–553.
- Goldblatt, P. 1988. Index to plant chromosome numbers 1984–1985. Monogr. Syst. Bot. Missouri Bot. Gard. 23: 1–264.
- Goldblatt, P. & Johnson, D.E. (eds.) 1979+. Index to plant chromosome numbers. http://www.tropicos.org/Project/IPCN (accessed Jul 2018).
- Lysak, M.A. & Dolezel, J. 1998. Estimation of nuclear DNA content in *Sesleria* (Poaceae). *Caryologia* 51: 123–132. https://doi.org/10. 1080/00087114.1998.10589127
- Májovský, J. & Murín, A. 1987. Karyotaxonomický prehľad flóry Slovenska. Bratislava: Veda Vydavateľsto Slovenskej Akadémie vied.
- Moore, R.J. 1973. Index to plant chromosome numbers for 1967– 1971. Regnum Vegetabile 90. Utrecht: Oosthoek, Scheltema & Holkema.
- Moore, R.J. 1977. Index to plant chromosome numbers for 1973–1974. Regnum Vegetabile 96. Utrecht: Oosthoek, Scheltema & Holkema.
- Padilla-García, N., Rojas-Andrés, B.M., López-González, N., Castro, M., Castro, S., Loureiro, J., Albach, D.C., Machon, N. & Martínez-Ortega, M.M. 2018. The challenge of species delimitation in the diploid-polyploid complex *Veronica* subsection *Pentasepalae*. *Molec. Phylogen. Evol.* 119: 196–209. https:// doi.org/10.1016/j.ympev.2017.11.007
- Peev, D. 1972. New taxa and ploidy levels of some Bulgarian Veronica species. Dokl. Bulg. Akad. Nauk. 25: 811–814.
- **Peev, D.** 1975. Genus *Veronica* in Bulgaria Some new taxa and chromosome numbers. *Fitologiya* 2: 42–46.
- Peev, D. 1976. [Report]. In: Löve, A. (ed.), IOPB Chromosome number reports LIV. *Taxon* 25: 631–649.

- Peev, D. 1978. Taxonomija i microevoljucija na divorastjastite predstaviteli na rod Veronica L. (velikdence) v Balgarija. Pp. 72–106 in: Kozuharov, S.I. & Kuzmanov, B.A. (ed.), Evolution of flowering plants and florogenesis, vol. 1, Fabaceae, Scrophulariaceae, Rubiaceae. Sofia: Bulgarian Academy of Sciences.
- Rojas-Andrés, B.M. & Martínez-Ortega, M.M. 2016. Taxonomic revision of Veronica subsection Pentasepalae (Veronica, Plantaginaceae sensu APG III). Phytotaxa 285: 1–100. https://doi. org/10.11646/phytotaxa.285.1.1
- Rojas-Andrés, B.M., Albach, D.C. & Martínez-Ortega, M.M. 2015. Exploring the intricate evolutionary history of the diploid– polyploid complex *Veronica* subsection *Pentasepalae* (Plantaginaceae). *Bot. J. Linn. Soc.* 179: 670–692. https://doi.org/10.1111/ boj.12345
- Rojas-Andrés, B.M., Rico, E. & Martínez-Ortega, M.M. 2016. A nomenclatural treatment for *Veronica* subsect. *Pentasepalae* (Plantaginaceae sensu APG III) and typification of several names. *Taxon* 65: 617–627. https://doi.org/10.12705/653.14
- Scheerer, H. 1937. Experimentelle und zytologische Untersuchungen innerhalb der Veronica-Gruppe Pentasepala. Flora 131: 287–323.
- Simonet, M. 1934. Contribution à l'étude caryologique et génétique des Veronica. Compt.-Rend. Hebd. Séances Mém. Soc. Biol. 117: 1153–1156.
- Temsch, E., Greilhuber, J. & Krisai, R. 2010. Genome size in liverworts. *Preslia* 82: 63–80.

Aleksander A. Gnutikov,^{1,2} Marina V. Protopopova,^{3*} Vasiliy V. Pavlichenko³ & Victor V. Chepinoga^{4,5}

- 1 The N.I. Vavilov Institute of Plant Genetic Resources (VIR), Bolshaya Morskaya Str. 44, 190000 St.-Petersburg, Russia
- 2 Komarov Botanical Institute of the Russian Academy of Sciences, Professora Popova Str. 2, 197376 St.-Petersburg, Russia
- 3 Siberian Institute of Plant Physiology and Biochemistry SB RAS, Lermontov Str. 132, 664033 Irkutsk, Russia
- 4 The V.B. Sochava Institute of Geography SB RAS, Ulan-Batorskaya Str. 1, 664033 Irkutsk, Russia
- 5 Irkutsk State University, Karl Marx Str. 1, 664003 Irkutsk, Russia
- * Author for correspondence: marina.v.protopopova@gmail.com

The study was partially supported by the Russian Foundation for Basic Research (projects 16-34-60135-mol_a_dk, 16-05-00783-a, 18-04-01040, 17-00-00338) and done within the framework of the State Task 0126-2014-0028 "Karyological and molecular-phylogenetic investigation of the flowering plant taxa divergence". The chromosome counting for highland populations was proceeded under financial support of the Russian Science Foundation (project 17-74-10074).

* First chromosome count for the species.

AMARYLLIDACEAE

Allium chamarense M.M.Ivanova

2n = 16, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), SE of Mt. Osinovskii Golets, alpine meadow, 1677 m, 51.41106°N, 104.16936°E, 31 Jul 2017, *M. Protopopova, V. Pavlichenko & E. Zolotovskaya Cl664* (IRKU).

APIACEAE

Pleurospermum uralense Hoffm.

2n = 18, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl642* (IRKU).

ASPARAGACEAE

Maianthemum bifolium (L.) F.W.Schmidt

2n = 36, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl641* (IRKU).

ASTERACEAE

Antennaria dioica (L.) Gaertn.

2n = 28, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), SE of Mt. Osinovskii Golets, alpine meadow, 1677 m, 51.41106°N, 104.16936°E, 31 Jul 2017, *M. Protopopova, V. Pavlichenko & E. Zolotovskaya Cl663* (IRKU).

Cirsium helenioides (L.) Hill

2n = 34, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl635* (IRKU).

Doronicum altaicum Pall.

2n = 60, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl647* (IRKU).

Jacobaea nemorensis (L.) E.Wiebe

2n = 40, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 1 Aug 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl673* (IRKU).

Parasenecio hastatus (L.) H.Koyama

2n = 60, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 1 Aug 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov C1672* (IRKU). Rhaponticum chamarense Peschkova (\equiv Rhaponticum carthamoides var. chamarense (Peschkova) O.S.Zhirova)

2n = 26, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl651* (IRKU).

Saussurea latifolia Ledeb.

2n = 26, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl637* (IRKU).

Solidago dahurica Kitag.

2n = 18, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl639* (IRKU); Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), SE of Mt. Osinovskii Golets, alpine meadow, 1677 m, 51.41106°N, 104.16936°E, 31 Jul 2017, *M. Protopopova, V. Pavlichenko & E. Zolotovskaya Cl667* (IRKU).

BORAGINACEAE

Myosotis palustris (L.) L.

2n = 22, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 30 Jul 2017, *M. Protopopova*, *V. Pavlichenko*, *E. Zolotovskaya & A. Konovalov Cl657* (IRKU).

BRASSICACEAE

Cardamine macrophylla Willd.

2n = 64, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 30 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl656* (IRKU).

CARYOPHYLLACEAE

Dianthus superbus L.

2n = 30, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), SE of Mt. Osinovskii Golets, alpine meadow, 1677 m, 51.41106°N, 104.16936°E, 31 Jul 2017, *M. Protopopova, V. Pavlichenko & E. Zolotovskaya Cl665* (IRKU).

CELASTRACEAE

Parnassia palustris L.

2n = 36, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 30 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl660* (IRKU).

GERANIACEAE

Geranium krylovii Tzvelev

2n = 28, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, the upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl655* (IRKU).

LAMIACEAE

Prunella vulgaris L.

2n = 28, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl640* (IRKU).

MELANTHIACEAE

Veratrum lobelianum Bernh.

2n = 32, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl653* (IRKU).

ONAGRACEAE

Circaea alpina L.

2n = 22, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 1 Aug 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl670* (IRKU).

OROBANCHACEAE

Pedicularis incarnata L.

2n = 16, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 30 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl658* (IRKU).

POACEAE

*Poa ×intricata Wein (P. nemoralis L. × P. palustris L.)

2n = 28, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 1 Aug 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov C1674* (IRKU). Plants determined by M. Olonova (Tomsk State University, Russia).

POLYGONACEAE

Rumex alpestris Jacq.

2n = 14, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl652* (IRKU).

PRIMULACEAE

Primula pallasii Lehm.

2n = 22, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova*, *V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl644* (IRKU).

RANUNCULACEAE

Aconitum baicalense Turcz. ex Rapaics

2n = 32, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl636* (IRKU).

Aconitum rubicundum Fisch. ex Steud.

2n = 16, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl638* (IRKU).

Anemone baicalensis Turcz.

2n = 28, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, middle course of the Bolshaya Osinovka River (near the confluence of the Pravaya Osinovka and Levaya Osinovka rivers), 7.5 km S of the Lake Baikal, mountain forest, 791 m, 51.43661°N, 104.22295°E, 26 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov C1643* (IRKU); Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov C1645* (IRKU).

Aquilegia glandulosa Fisch. ex Link

2n = 14, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), SE of Mt, Osinovskii Golets, alpine meadow, 1677 m, 51.41106°N, 104.16936°E, 31 Jul 2017, *M. Protopopova, V. Pavlichenko & E. Zolotovskaya Cl668* (IRKU).

Caltha palustris L.

2n = 32, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 30 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl662* (IRKU).

VIOLACEAE

Viola biflora L.

2n = 12, CHN. Russia, Eastern Siberia, Irkutskaya Oblast', Sludyanskii Raion, the Khamar-Daban Ridge, upper course of the Irkut River (tributary of the Khara-Murin River), wet meadow at a stream bank, 1384 m, 51.40333°N, 104.15485°E, 29 Jul 2017, *M. Protopopova, V. Pavlichenko, E. Zolotovskaya & A. Konovalov Cl649* (IRKU).

Iva Hodálová,^{1*} Lenka Mártonfiová,² Marilena Onete,³ Jaromír Kučera¹ & Pavol Mereďa, Jr.¹

- 1 Institute of Botany, Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovak Republic
- 2 Pavol Jozef Šafárik University, Botanical Garden, Mánesova 23, 04352 Košice, Slovak Republic
- 3 Institute of Biology Bucharest, Romanian Academy, 296 Splaiul Independentei, 060031 Bucharest, Romania
- * Author for correspondence: iva.hodalova@savba.sk

Chromosomes were counted in 2018 by L. Mártonfiová. DNA ploidy levels were estimated in 2018 by P. Mered'a, Jr. Authors thank to M. Zaliberová for assistance in the field.

This study was supported by the Grant Agency of Ministry of Education, Science, Research and Sport of the Slovak Republic and Slovak Academy of Sciences VEGA (grant no. 2/0131/16).

POACEAE

Sesleria heufleriana Schur

2*n* = 4*x* = 28, CHN. Romania, Apuseni Mts, Cluj County, Petreștii de Jos village, Turzii Gorges, 46°34'12"N, 23°40'16"E, 530–560 m, 17 May 2018, *I. Hodálová*, *P. Mereďa*, *Jr. & M. Zaliberová 137-1* (SAV) [Fig. 4A].

 $2n \sim 4x \sim 28$, FCM. Romania, Apuseni Mts, Cluj County, Petreștii de Jos village, Turzii Gorges, 46°34'12"N, 23°40'16"E, 530-560 m, 17 May 2018, I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 137-1 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-2 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-3 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 137-4 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 137-5 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-7 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-8 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-9 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-10 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-11 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 137-12 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 137-13 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-14 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 137-15 (SAV). Romania, Apuseni Mts, Alba County, Colțești village, Piatra Secuiului Mt., 46°25′56″N, 23°34′56″E, 740-840 m, 18 May 2018, I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 141-1 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-2 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 141-5 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 141-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-7 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-8 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-9 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-10 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-11 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 141-12 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 141-14 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 141-15 (SAV). Romania, Southern Carpathians, Sibiu County, Tălmaciu town, Măgura hill, 45°38'38"N, 24°16'55"E, ca. 500 m (type locality of S. heufleriana), 18 May 2018, I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 142-1 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 142-2 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-3 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-4 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-5 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-7 (SAV),
I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-8 (SAV), I. Hodálová,
P. Mereďa, Jr. & M. Zaliberová 142-9 (SAV), I. Hodálová, P. Mereďa,
Jr. & M. Zaliberová 142-10 (SAV), I. Hodálová, P. Mereďa, Jr. &
M. Zaliberová 142-11 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-12 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-13 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-14 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-14 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 142-15 (SAV).

Sesleria rigida Rchb.

2n = 4x = 28, CHN. Romania, Apuseni Mts, Alba County, Poşaga de Sus village, Poşegii Gorges, 46°27′36″N, 23°23′52″E, ca. 560 m, 17 May 2018, *I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 139-1* (SAV) [Fig. 4B].

2n ~ 4x ~ 28, FCM. Romania, Apuseni Mts, Alba County, Runc village, Runcului Gorges, 46°30'27"N, 23°26'29"E, 550–560 m, 17 May 2018, I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-1 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-2 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-3 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-4 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-5 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-7 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-8 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-9 (SAV), I. Hodálová, P. Mereďa,



Fig. 4. Mitotic metaphase. **A**, *Sesleria heufleriana* Schur, 2n = 28; **B**, *S. rigida* Rchb., 2n = 28.

Jr. & M. Zaliberová 138-10 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 138-11 (SAV). Romania, Apuseni Mts, Alba County, Poşaga de Sus village, Poşegii Gorges, 46°27'36"N, 23°23'52"E, ca 560 m, 17 May 2018, I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 139-1 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 139-2 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 139-3 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 139-4 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 139-5 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 139-6 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 139-7 (SAV). Romania, Apuseni Mts, Alba County, Colțești village, Piatra Secuiului Mt., 46°25'56"N, 23°34'56"E, ca. 750 m, 18 May 2018, I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 140-1 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 140-2 (SAV). Romania, Apuseni Mts. Alba County, Vălișoara, Vălișoarei Gorges, 46°22'31"N, 23°35'02"E, ca. 460 m, 18 May 2018, I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 143-1 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 143-2 (SAV), I. Hodálová, P. Mereďa, Jr. & M. Zaliberová 143-4 (SAV), I. Hodálová, P. Mered'a, Jr. & M. Zaliberová 143-5 (SAV). Romania, Southern Carpathians, Braşov County, Braşov town, Stejărișul Mare Protected Area from N-W Postăvaru Massif, 45°38'02"N, 25°33'08"E, ca. 930 m, 19 May 2018, P. Mered'a, Jr. 144-1 (SAV), P. Mered'a, Jr. 144-2 (SAV), P. Mered'a, Jr. 144-3 (SAV), P. Mered'a, Jr. 144-4 (SAV), P. Mered'a, Jr. 144-5 (SAV), P. Mered'a, Jr. 144-6 (SAV), P. Mered'a, Jr. 144-7 (SAV), P. Mered'a, Jr. 144-8 (SAV), P. Mered'a, Jr. 144-9 (SAV), P. Mered'a, Jr. 144-10 (SAV), P. Mered'a, Jr. 144-11 (SAV), P. Mered'a, Jr. 144-12 (SAV), P. Mered'a, Jr. 144-13 (SAV), P. Mered'a, Jr. 144-14 (SAV).

CHN

For the chromosome counts, root tip meristems of potted plants were employed. The actively growing root tips were pre-treated in a 0.002 M water solution of 8-hydroxyquinoline at 4°C for about 16 hours (overnight), fixed in a 1:3 mixture of 98% acetic acid and 96% ethanol for 1-24 hours, washed in distilled water and then in citrate buffer with the pH 5.8. They were macerated in the mixture of enzymes: 0.5% cellulase ("Onozuka R-10" from Trichoderma viride 1.3 U/mg) and 0.5% pectinase (from Aspergillus niger 0.2 U/ mg; Serva, Germany) in citrate buffer (pH 5.8) at 37°C for 4 hours. Then the roots were washed in the citrate buffer overnight at 4°C. After washing the root-tips were hydrolyzed in 1 N HCl at 60°C for 8 min and washed in distilled water. Squashes were made using the cellophane squash technique (Murín, 1960). The slides were stained by a 7% solution of Giemsa Stain, Modified Solution, Fluka Analytical, in Sörensen phosphate buffer, dried and observed in a drop of immersion oil using a Leica DM 2500 microscope equipped with camera HDCE-X5 and software ScopeImage v.9.0.

FCM

DAPI. Fresh leaf tissue were used. The sample preparation and FCM procedure followed that of Budzáková & al. (2014). Internal reference: *Pisum sativum* 'Ctirad', 2C = 8.76 pg (Trávníček & al., 2015). Fluorescence intensity for *S. heufleriana* varied from 0.748 to 0.778 (mean 0.758) and for *S. rigida* from 0.686 to 0.705 (mean 0.692). Coefficients of variation (CVs) of samples and internal standard were 1.27%–3.09% (mean 2.35%) and 1.05%–3.18% (mean 1.75%), respectively.

LITERATURE CITED

Budzáková, M., Hodálová, I., Mered'a, P., Jr., Somlyay, L., Bisbing, S.M. & Šibík, J. 2014. Karyological, morphological and ecological differentiation of *Sesleria caerulea* and *S. tatrae* in the Western Carpathians and adjacent regions. *Preslia* 86: 245–277.

- Murín, A. 1960. Substitution of cellophane for glass covers to facilitate preparation of permanent squashes and smears. *Stain Tech*nol. 35: 351–353.
- Trávníček, P., Ponert, J., Urfus, T., Jersáková, J., Vrána, J., Hřibová, E., Doležel, J. & Suda, J. 2015. Challenges of flowcytometric estimation of nuclear genome size in orchids, a plant group with both whole-genome and progressively partial endoreplication. *Cytometry A* 87: 958–966. https://doi. org/10.1002/cyto.a.22681

Navjot Kaur* & Raghbir Chand Gupta

Department of Botany, Punjabi University Patiala, India * Author for correspondence: navjot21188@gmail.com

The study is supported by financial grant under DRS SAP I, SAP II and SAP III of UGC; UGC-BSR Indira Gandhi Single Girl Child Fellowship (Award letter no. F7-152/2007 BSR, dated 16/12/2013) to Navjot Kaur and IPLS-DBT project (Project no. BT/PR-4548/ INF/22/146/2012) sanctioned to Punjabi University, Patiala for using the facilities and financial support of this study.

* First chromosome count for the species.

- ** New cytotype for the species.
- ▼ First chromosome report for an Indian accession.

ASPARAGACEAE

** Asparagus racemosus Willd.

n = 30, CHN. India, Rajasthan, Udaipur, 24°34′31.58″N, 73°41′35.97″E, 600 m, Gulab bagh, 16 Sep 2015, *N. Kaur 33823* (PUN 60936) [Fig. 5A].

Previously, this species was known to have chromosome number of n = 10 (Kamble, 1997), 2n = 20 (Sastry, 1981) from India and also 2n = 20, 40 (Sheidai & Inamdar, 1997) from outside of India.

CANNACEAE

** Canna indica L.

n = 18, CHN. India, Rajasthan, Sriganganagar, 29°56'50.57"N, 73°53'52.58"E, 178 m, 26 Oct 2015, *N. Kaur 33893* (PUN 61126) [Fig. 5B].

Previously, this species was known to have chromosome number of n = 9 (Khoshoo & Mukherjee, 1970) from India and 2n = 18 from outside India (Hanson & al., 2001).

COMMELINACEAE

***Commelina attenuata* Vahl

n = 12, CHN. India, Rajasthan, Udaipur, Gulab bagh, 24°34'31.58"N, 73°41'35.97"E, 600 m, 15 Sep 2015, *N. Kaur 33825* (PUN 60938) [Fig. 5C].

Previously, this species was reported to have 2n = 44 by Renugadevi & Sampathkumar (1983, 1986) from India.

** Commelina diffusa Burm.f.

n = 60, CHN. India, Rajasthan, Mount Abu, Naki lake, 24°35′40.41″N, 72°42′22.41″E, 1220 m, 30 Feb 2016, *N. Kaur 33801* (PUN 60779) [Fig. 5D].

Previously, the species was reported with n = 15 (Alam & Sharma, 1984; Patwary & al., 1987; Umoh & al., 1991), 2n = 18 (Zheng & al., 1989), 2n = 28 (Palomino & al., 1990), 2n = 30 (Cristobal de Hinojo & al., 1998), n = 45 (Grabiele, 2005) and 2n = 72 (Faden & Suda, 1980).

▼ *Commelina forskalaei* Vahl.

n = 45, CHN. India, Rajasthan, Jhalawar, Gagron fort, 24°37′40.99″N, 76°11′09.76″E, 312 m, 21 Aug 2014, *N. Kaur 33824* (PUN 60937) [Fig. 5E].

For this species, it was previously reported 2n = 28 from West Africa (Morton 1967) and 2n = 30 from United States by Faden (1993).

** Commelina hasskarlii C.B.Clarke (= C. caroliniana Walter)

n = 60, CHN. India, Rajasthan, Mount Abu, Naki lake, 24°35′40.41″N, 72°42′22.41″E, 1220 m, 31 Aug 2016, *N. Kaur 33887* (PUN 61120) [Fig. 5F]; India, Rajasthan, Udaipur, 24°35′07.60″N, 73°42′44.92″E, 600 m, 16 Aug 2015, *N. Kaur 33888* (PUN 61121).

Previously, this species was reported with n = 15 (Patwary & al., 1987) and 2n = 90 (Raghavan & Rao, 1961) from different parts of India.

**Commelina paludosa Blume

n = 15, CHN. India, Rajasthan, Pali, 25°46'25.77"N, 73°19'27.75"E, 218 m, 30 Aug 2014, *N. Kaur 33883* (PUN 61116) [Fig. 5G]; India, Rajasthan, 24°35'35.99"N, 73°38'21.45"E, 850 m, 15 Sep 2015, *N. Kaur 33884* (PUN 61117).

Previously, this species was reported with n = 30 by Patwary & al. (1987), 2n = 60 by Renugadevi & Sampathkumar (1986), n = 56, 60 by Bhattacharya (1975) and 2n = 75 by Baquar & Saeed (1977).

** Commelina suffruticosa Blume

n = 15, CHN. India, Rajasthan, Mount Abu, 24°35′34.98″N, 72°42′58.23″E, 1220 m, 30 Aug 2016, *N. Kaur 33889* (PUN 61122) [Fig. 5H].

Previously, this species was recorded only with 2n = 64 by Alam & Sharma (1984).

**Cyanotis cristata (L.) D.Don

n = 11, CHN. India, Rajasthan, Jhalawar, Bheem Sagar dam, 24°33'46.43"N, 76°20'05.52"E, 312 m, 24 Aug 2014, *N. Kaur 33882* (PUN 61115) [Fig. 51].

Previously, this species was observed with n = 12 by Alam & Sharma (1984) and 2n = 24 (Bhattacharya, 1975; Renugadevi & Sampathkumar, 1986; Lalithambika Bai & Kuriachan, 1997) from different parts of India.

**Murdannia nudiflora (L.) Brenan

n = 40, CHN. India, Rajasthan, Mount Abu, Dilwara Temple, 24°36′33.95″N, 72°43′22.71″E, 1230 m, 1 Sep 2016, *N. Kaur 33886* (PUN 61119) [Fig. 5J].

This species was previously reported with chromosome numbers of n = 10 (Bhattacharya, 1975), 2n = 20 (Bhatt, 1976; Alam & Sharma, 1984; Renugadevi & Sampathkumar, 1986) from India and from outside of India.

POACEAE

**Brachiaria deflexa (Schumach.) C.E.Hubb. ex Robyns

n = 18, CHN. India, Rajasthan, Jhalawar, Jhalrapatan, 24°33'02.25"N, 76°10'15.15"E, 312 m, 13 Aug 2013, *N. Kaur 31979* (PUN 60211) [Fig. 5K]; India, Rajasthan, Udaipur, Railway station, 24°34'35.29"N, 73°42'03.21"E, 600 m, 15 Sep 2015, *N. Kaur 31985* (PUN 60217).

This species was previously reported with chromosome numbers of n = 9 by Basappa & al. (1987) from India and by Hoshino & Davidse (1988) from southern Africa, n = 18 from Pakistan (Ahsan & al., 1994), and 2n = 18 (Basappa & Muniyamma, 1981) from India. ▼ *Chloris prieurii* Kunth (≡ *Enteropogon prieurii* (Kunth) Clayton)

n = 10, CHN. India, Rajasthan, Udaipur, Monsoon Palace, 24°35'35.90"N, 73°38'21.23"E, 850 m, 27 Aug 2014, *N. Kaur 31932* (PUN 60164) [Fig. 5L]; India, Rajasthan, Churu, 28°17'32.71"N, 74°57'44.12"E, 292 m, 19 Oct 2013, *N. Kaur 31954* (PUN 60186).

Previously, this species was reported with chromosome number of 2n = 40 by Miege & Josserand (1972) from outside of India.

*Chloris quinquesetica Bhide

n = 10, CHN. India, Rajasthan, Pali, 25°44′59.45″N, 73°20′03.81″E, 214 m, 15 Mar 2015, *N. Kaur 31976* (PUN 60208) [Fig. 5M].

▼ *Eragrostis aspera* (Jacq.) Nees,

n = 10, CHN. India, Rajasthan, Mount Abu, 24°35′33.88″N, 72°42′58.84″E, 1220 m, 30 Aug 2016, *N. Kaur 33848* (PUN 60980) [Fig. 5N].



Fig. 5. A, Asparagus racemosus, meiotic anaphase-I, n = 30 (PUN 60936); **B**, Canna indica, meiotic metaphase-I, n = 18 (PUN 61126); **C**, Commelina attenuata, meiotic metaphase-I, n = 12 (PUN 60938); **D**, C. diffusa, meiotic diakinesis n = 60 (PUN 60779); **E**, C. forskalaei, meiotic diakinesis, n = 45 (PUN 60937); **F**, C. hasskarlii, meiotic diakinesis, n = 60 (PUN 61120); **G**, C. paludosa, meiotic anaphase-I, n = 15 (PUN 61116); **H**, C. suffruticosa, meiotic diakinesis, n = 15 (PUN 61122); **I**, Cyanotis cristata, meiotic metaphase-I, n = 11 (PUN 61115); **J**, Murdannia nudiflora, meiotic diakinesis, n = 40 (PUN 61119); **K**, Brachiaria deflexa, meiotic diakinesis, n = 18 (PUN 60211); **L**, Chloris prieurii, meiotic metaphase-I, n = 10 (PUN 60164); **M**, C. quinquesetica, meiotic metaphase-I, n = 10 (PUN 60208); **N**, Eragrostis aspera, meiotic metaphase-I, n = 10 (PUN 60980); **O**, Leptothrium senegalense, meiotic metaphase-I, n = 10 (PUN 60149); **P**, Sporobolus coromandelianus, meiotic metaphase-I, n = 24 (PUN 60167)

For this species the chromosome number of 2n = 20 was reported by Moffett & Hurcombe (1949) from outside of India.

▼ Leptothrium senegalense (Kunth) Clayton

n = 10, CHN. India, Rajasthan, Churu, wild life sanctuary, 27°47′55.91″N, 74°27′40.97″E, 292 m, 20 Oct 2014, *N. Kaur 31917* (PUN 60149) [Fig. 50].

Previously, this species was recorded with n = 10 by Moinuddin & al. (1994) from Pakistan.

** Sporobolus coromandelianus (Retz.) Kunth

n = 24, CHN. India, Rajasthan, Pali, Hemawas Dam, 25°24'42.82"N, 73°15'18.50"E, 218 m, 30 Aug 2014, *N. Kaur 31935* (PUN 60167) [Fig. 5P].

This species was reported previously with n = 18 (Bir & Sahni, 1985; Bir & al., 1988) from India and from outside of India by Moinuddin & al. (1994). Gupta & al. (2014) observed also 2n = 36 from India.

LITERATURE CITED

- Ahsan, S.M.N., Vahidy, A.A. & Ali, S.I. 1994. Chromosome numbers and incidence of polyploidy in Panicoideae (Poaceae) from Pakistan. Ann. Missouri Bot. Gard. 81: 775–783. https://doi. org/10.2307/2399922
- Alam, N. & Sharma, A.K. 1984. Trends of chromosome evolution in family Commelinaceae. *Nucleus (Calcutta)* 27: 231–241.
- Baquar, S.R. & Saeed, V.A. 1977. Aneusomaty in *Commelina* sp. from southern West Pakistan. *Kromosomo* 6: 163–169.
- Basappa, G.P. & Muniyamma, M. 1981. [Reports]. In: Löve, Á. (ed.), Chromosome number reports LXXII. *Taxon* 30: 703.
- Basappa, G.P., Muniyamma, M. & Chinnappa, C.C. 1987. An investigation of chromosome numbers in the genus *Brachiaria* (Poaceae: Paniceae) in relation to morphology and taxonomy. *Canad. J. Bot.* 65: 2297–2309. https://doi.org/10.1139/b87-313
- Bhatt, R.P. 1976. [Reports]. In: Löve, Å. (ed.), IOPB chromosome number reports LIII. *Taxon* 25: 494.
- Bhattacharya, B. 1975. Cytological studies on some Indian members of Commelinaceae. *Cytologia* 40: 285–299. https://doi. org/10.1508/cytologia.40.285
- Bir, S.S. & Sahni, M. 1985. Cytological investigations on some grasses from Punjab Plain, North India. *Proc. Indian Natl. Sci. Acad. B* 51: 609–626.
- Bir, S.S., Sahni, M. & Singh, C.P. 1988. Cytology of genus Sporobolus R. Br. from North India (Punjab Plain). Cytologia 53: 53–57. https://doi.org/10.1508/cytologia.53.53
- Cristobal de Hinojo, M.E., González Villarreal, L.M. & Frías de Fernández, A. M. 1998. Estudios citogenéticos en el género Commelina L. (Commelinaceae). I. Análisis cariotípico de Commelina erecta L. y Commelina diffusa Burm. Lilloa 39: 157–164.
- Faden, R.B. 1993. The misconstrued and rare species of *Commelina* (Commelinaceae) in the Eastern United States. *Ann. Missouri Bot. Gard.* 80: 208–218. http://doi.org/10.2307/2399824
- Faden, R.B. & Suda, Y. 1980. Cytotaxonomy of Commelinaceae: Chromosome counts of some African and Asiatic species. *Bot. J. Linn. Soc.* 81: 301–325. https://doi.org/10.1111/j.1095-8339.1980. tb01681.x
- Grabiele, M. 2005. Chromosomes of four species of *Commelina* (Commelinaceae). *Bot. J. Linn. Soc.* 148: 207–218. https://doi. org/10.1111/j.1095-8339.2005.00401.x
- Gupta, R.C., Chauhan, H.S., Saggoo, M.I.S. & Kaur, N. 2014.

Cytomorphology of some grasses (Poaceae) from Lahaul-Spiti (Himachal Pradesh), India. *Biolife* 2: 1234–1247.

- Hanson, L., McMahon, K.A., Johnson, M.A.T. & Bennett, M.D. 2001. First nuclear DNA C-values for another 25 angiosperm families. Ann. Bot. (Oxford) 88: 851–858. https://doi.org/10.1006/ anbo.2001.1528
- Hoshino, T. & Davidse. G. 1988. Chromosome numbers of grasses (Poaceae) from southern Africa. I. Ann. Missouri Bot. Gard. 75: 866–873. https://doi.org/10.2307/2399375
- Kamble, S.Y. 1997. Studies on the genus Asparagus L. (sensu lato) in India: Endemic species and their uses. J. Econ. Taxon. Bot. 21: 167–174.
- Khoshoo, T.N. & Mukherjee, I. 1970. Genetic-evolutionary studies on cultivated Cannas. 3. Variation in meiotic system. *Cellule* 68: 107–134.
- Lalithambika Bai, K. & Kuriachan, P.I. 1997. Cytological evolution among south Indian Cyanotis D. Don (Commelinaceae). Caryologia 50: 163–174. https://doi.org/10.1080/00087114.1997.10797395
- Miege, J. & Josserand. N. 1972. Nombres chromosomiques d'especes africaines et malgaches. *Candollea* 27: 283–292.
- Moffett, A.A. & Hurcombe, R. 1949. Chromosome numbers in South African grasses. *Heredity* 3: 369–373. https://doi.org/10.1038/ hdy.1949.27
- Moinuddin, M., Vahidy, A.A. & Ali, S.I. 1994. Chromosome counts in Arundinoideae, Chloridoideae, and Poideae (Poaceae) from Pakistan. Ann. Missouri Bot. Gard. 81: 784–791. https://doi. org/10.2307/2399923
- Morton, J.K. 1967. The Commelinaceae of West Africa: A biosystematic survey. *Bot. J. Linn. Soc.* 60: 167–221. https://doi. org/10.1111/j.1095-8339.1967.tb00085.x
- Palomino, H.G., Benítez, B.V., Almeraya P.M. & Ruaro P.M. 1990. Estudios cromosómicos en seis especies Mexicanas de Comelináceas. Bol. Soc. Bot. México 50: 77–87.
- Patwary, M.U., Mirza S.H. & Zaman. M.A. 1987. Meiotic behaviour of chromosomes and the nature of ploidy in four species of *Commelina* (Commelinaceae) from Bangladesh. *Cytologia* 52: 175–181. https://doi.org/10.1508/cytologia.52.175
- Raghavan, R.S. & Rao, S.R. 1961. Cytological observations on the Indian species of Commelinaceae. *Curr. Sci.* 30: 310–311.
- Renugadevi, K. & Sampathkumar, R. 1983. [Reports]. In: Löve, Á. (ed.), IOPB chromosome number reports LXXIX. *Taxon* 32: 320–321.
- Renugadevi, K. & Sampathkumar, R. 1986. On the karyomorphological delineations in some taxa of Commelinaceae. J. Cytol. Genet. 21: 115–132.
- Sastry, M. 1981. Chromosome studies and DNA estimation in Asparagus sp. Proc. Indian Sci. Congr. Assoc. 68 (Sect. VI): 85.
- Sheidai, M. & Inamdar, A.C. 1997. Cytomorphology of Asparagus taxa using multivariate statistical analysis. Nucleus (Calcutta) 40: 7–12.
- Umoh, E.O., Hossain, G.M. & Ene-Obong, E.E. 1991. On interchange complexes in *Commelina diffusa* Burm. f. from south eastern Nigeria. *Cytologia* 56: 173–179. https://doi.org/10.1508/ cytologia.56.173
- Zheng, J.Y., Gu C.Y. & Chen. R.Y. 1989. Cytotaxonomical studies on Commelinaceae in China I. Chromosome numbers and karyotypes of some Chinese species. Pp. 363–368 in: Hong, D. (ed.), Plant chromosome research 1987: Proceedings of the Sino-Japanese Symposium on Plant Chromosomes Held at Beijing, China, October 6–11, 1987. Beijing: Organizing Committee of the Symposium.

Jaromír Kučera,^{1*} Eliška Štubňová,^{1,2} Lenka Mártonfiová,³ Filip Kolář,^{4,5} Jan Smyčka⁶ & Marek Slovák^{1,4}

- 1 Plant Science and Biodiversity Centre, Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 9, 845 23 Bratislava, Slovak Republic
- 2 Department of Botany, Natural History Museum, Slovak National Museum, Vajanského nábrežie 2, 810 06 Bratislava, Slovak Republic
- 3 Pavol Jozef Šafárik University, Botanical Garden, Mánesova 23, Košice, Slovak Republic
- 4 Department of Botany, Charles University, Benátská 2, 128 01 Praha 2, Czech Republic
- 5 Institute of Botany, The Czech Academy of Sciences, Zámek 1, 252 43 Průhonice, Czech Republic
- 6 Laboratoire d'Ecologie Alpine (LECA), Université Grenoble Alpes-CNRS, 2233 rue de la Piscine, 38000 Grenoble, France
- * Author for correspondence: jaromir.kucera@savba.sk
 - * The first chromosome count and genome size (C-values) for the taxon.

This study was supported by the Grant Agency VEGA, Bratislava, Slovak Republic (grant no. VEGA 2/0088/15).

PRIMULACEAE

Soldanella angusta L.B.Zhang

The coefficient of variation (CV) of samples and internal standard ranged from 1.66% to 3.09% (mean 2.52%) and from 1.64% to 3.14% (mean 2.62%), respectively.

2n = 4x = 40, CHN. Ukraine, Zakarpatskaya oblasť, Polonina Borzhava Mts., Gemba Mt., 48°38′03.9″N, 23°15′52.3″E, 1400 m, 10 May 2018, *M. Valachovič UA8/I3* (SAV) [Fig. 6A].

 $2n \sim 4x \sim 40$, 2C = 3.32-3.36 pg, FCM. Ukraine, Zakarpatskaya oblasť, Skhidni Beskidi Mts., Pikui Mt., 48°49'56.4"N, 22°59'47.8"E, 1400 m, 8 May 2018, *M. Valachovič UA6/2* (SAV), *M. Valachovič UA6/5* (SAV), *M. Valachovič UA6/15* (SAV).

2*n* ~ 4*x* ~ 40, 2C = 3.32–3.38 pg, FCM. Ukraine, Zakarpatskaya oblasť, Gorgany Mts., Horb Mt., 48°27'38.2"N, 23°41'43.4"E, 1700 m, 9 May 2018, *M. Valachovič UA7/4* (SAV), *M. Valachovič UA7/7* (SAV), *M. Valachovič UA7/9* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.31-3.35 pg, FCM. Ukraine, Zakarpatskaya oblasť, Polonina Borzhava Mts., Gemba Mt., 48°38′03.9″N, 23°15′52.3″E, 1400 m, 10 May 2018, *M. Valachovič UA8/11* (SAV), *M. Valachovič UA8/13* (SAV), *M. Valachovič UA8/14* (SAV).

Soldanella alpina L. subsp. alpina

The CV of samples and internal standard ranged from 1.49% to 3.52% (mean 2.42%) and from 1.5% to 3.49% (mean 2.49%), respectively.

2n = 4x = 40, CHN. Austria, Vorarlberg, Damüls village, Furkajoch saddle, 47°16'00.0"N, 09°49'59.3"E, 1784 m, 13 Aug 2018, *J. Kučera & M. Slovák AT39/2* (SAV) [Fig. 6B]; Switzerland, Valais, Conthey town, Col du Sanetsch saddle, 46°19'55.4"N, 07°17'00.4"E, 2258 m, 15 Aug 2018, *J. Kučera & M. Slovák SW5/5* (SAV) [Fig. 6C].

 $2n \sim 4x \sim 40$, 2C = 3.61-3.67 pg, FCM. Austria, Nordtirol, Imst town, Hahntennjoch saddle, $47^{\circ}17'08.7''$ N, $10^{\circ}40'23.0''$ E, 1768 m, 13 Aug 2018, *J. Kučera & M. Slovák AT37/1* (SAV), *J. Kučera & M. Slovák AT37/2* (SAV), *J. Kučera & M. Slovák AT37/5* (SAV).

2*n* ~ 4*x* ~ 40, 2C = 3.59–3.64 pg, FCM. Austria, Nordtirol, Arlberg village, Flexenpass saddle, 47°09′55.9″N, 10°09′50.2″E, 1733 m, 13 Aug 2018, *J. Kučera & M. Slovák AT38/1* (SAV), *J. Kučera & M. Slovák AT38/2* (SAV), *J. Kučera & M. Slovák AT38/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.62-3.65 pg, FCM. Austria, Vorarlberg, Damüls village, Furkajoch saddle, 47°16′00.0″N, 09°49′59.3″E, 1784 m, 13 Aug 2018, *J. Kučera & M. Slovák AT39/1* (SAV), *J. Kučera & M. Slovák AT39/2* (SAV), *J. Kučera & M. Slovák AT39/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.52-3.60 pg, FCM. France, Rhône Alpes, Grenoble town, Lac du Crozet, $45^{\circ}10'23.0''$ N, $05^{\circ}56'11.3''$ E, 1985 m, 21 Aug 2018, *J. Smyčka FR2/1* (SAV), *J. Smyčka FR2/2* (SAV), *J. Smyčka FR2/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.65-3.68 pg, FCM. France, Rhône Alpes, Saligniere village, Le Taillefer, along the path from Lac de Poursollet to refuge de Taillefer, $45^{\circ}03'48.3''$ N, $05^{\circ}55'43.9''$ E, 2035 m, 24 Aug 2018, J. Smyčka FR3/1 (SAV), J. Smyčka FR3/2 (SAV), J. Smyčka FR3/3 (SAV).

 $2n \sim 4x \sim 40, 2C = 3.56-3.57$ pg, FCM. Italy, Südtirol, Trafoi village, Stilfserjoch saddle, 46°31'43.7"N, 10°27'41.5"E, 2469 m, 16 Aug 2018, J. Kučera & M. Slovák IT22/1 (SAV), J. Kučera & M. Slovák IT22/2 (SAV), J. Kučera & M. Slovák IT22/3 (SAV).

2*n*~4*x*~40, 2C = 3.56–3.61 pg, FCM. Switzerland, Ticino, Hospental village, St. Gotthard pass saddle, 46°35'38.1"N, 08°33'43.6"E, 1745 m, 14 Aug 2018, *J. Kučera & M. Slovák SW2/1* (SAV), *J. Kučera & M. Slovák SW2/2* (SAV), *J. Kučera & M. Slovák SW2/4* (SAV).

 $2n \sim 4x \sim 40, 2C = 3.63 - 3.64$ pg, FCM. Switzerland, Valais, Wichel village, Nufenen pass saddle, 46°28′20.8″N, 08°22′57.0″E, 2307 m, 14



Fig. 6. Mitotic metaphases of *Soldanella* species. **A**, *S. angusta* (pop. UA8), 2n = 4x = 40; **B**, *S. alpina* subsp. *alpina* (pop. AT39), 2n = 4x = 40; **C**, *S. alpina* subsp. *alpina* (pop. SW5), 2n = 4x = 40; **D**, *S. major* (pop. RO23), 2n = 4x = 40; **E**, *S. pusilla* subsp. *alpicola* (pop. SW3), 2n = 4x = 40; **F**, *S. ×hybrida* (pop. SW4), 2n = 4x = 40. — Scale bars 10 µm.

Aug 2018, J. Kučera & M. Slovák SW4/I (SAV), J. Kučera & M. Slovák SW4/2 (SAV), J. Kučera & M. Slovák SW4/3 (SAV) [Fig. 7D].

 $2n \sim 4x \sim 40$, 2C = 3.58-3.63 pg, FCM. Switzerland, Valais, Conthey town, Col du Sanetsch saddle, $46^{\circ}19'55.4''$ N, $07^{\circ}17'00.4''$ E, 2258 m, 15 Aug 2018, *J. Kučera & M. Slovák SW5/2* (SAV), *J. Kučera & M. Slovák SW5/5* (SAV), *J. Kučera & M. Slovák SW5/5* (SAV).

Soldanella major (Nielr.) Vierh.

CVs of samples and internal standard ranged from 2.08% to 2.99% (mean 2.59%) and from 2.15% to 2.99% (mean 2.71%), respectively. Our results confirm a previously published chromosome number (2n = 40) for this taxon (Kress, 1984; Štubňová & al., 2017).

2*n* = 4*x* = 40, CHN. Romania, Cozia Mts., near Călimănești village, 45°18'52.9"N, 24°20'51.4"E, 1400 m, 29 Aug 2018, *E. Štubňová RO23/3* (SAV) [Fig. 6D].

2*n* ~ 4*x* ~ 40, 2C = 3.28–3.32 pg, FCM. Romania, Cozia Mts., near Călimănești village, 45°18′52.9″N, 24°20′51.4″E, 1400 m, 29 Aug 2018, *E. Štubňová RO23/1* (SAV), *E. Štubňová RO23/2* (SAV), *E. Štubňová RO23/3* (SAV).

Soldanella minima Hoppe

The CV of samples and internal standard ranged from 2.51% to 3.16% (mean 2.97%) and from 2.14% to 2,89% (mean 2.51%), respectively.

 $2n \sim 4x \sim 40$, 2C = 3.63-3.68 pg, FCM. Austria, Nordtirol, Mutters village, Nockspitze, 47°11′26.8″N, 11°19′26.5″E, 2369 m, 14 Aug 2018, *F. Kolář AT42/1* (SAV), *F. Kolář AT42/2* (SAV).

Soldanella pusilla subsp. alpicola (F.K.Mey.) Chrtek

CVs of samples and internal standard ranged from 1.57% to 3.04% (mean 2.29%) and from 2.25% to 3.42% (mean 2.89%), respectively. Our results are in accordance with our previously published chromosome number (2n = 40) for this taxon (Štubňová & al., 2017).

2n = 4x = 40, CHN. Switzerland, Ticino, St. Gotthard pass saddle, near Lago di San Carlo, 46°33'29.3"N, 08°33'25.5"E, 2116 m, 14 Aug 2018, *J. Kučera & M. Slovák SW3/2* (SAV) [Fig. 6E].

 $2n \sim 4x \sim 40$, 2C = 3.48-3.49 pg, FCM. Austria, Nordtirol, Kaunertal valley, near Weisssee glacial lake, $46^{\circ}52'12.5''$ N, $10^{\circ}42'44.8''$ E, 2505 m, 25 Jul 2018, *J. Kučera AT35/1* (SAV), *J. Kučera AT35/2* (SAV), *J. Kučera AT35/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.45–3.47 pg, FCM. Austria, Nordtirol, Fiss village, Zwölferkopf Mt., 47°04′25.5″N, 10°34′49.1″E, 2546 m, 27 Jul 2018, *J. Kučera AT36/1* (SAV), *J. Kučera AT36/2* (SAV), *J. Kučera AT36/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.43-3.48 pg, FCM. Switzerland, Graubünden, Vals village, above Zervriela lake, $46^{\circ}34'16.8''$ N, $09^{\circ}06'45.8''$ E, 1948 m, 14 Aug 2018, *J. Kučera & M. Slovák SW1/1* (SAV), *J. Kučera & M. Slovák SW1/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.44-3.49 pg, FCM. Switzerland, Graubünden, Realp village, Furkapass saddle, $46^{\circ}34'22.8''$ N, $08^{\circ}24'49.9''$ E, 2467 m, 15 Aug 2018, *J. Kučera & M. Slovák SW6/1*, (SAV), *J. Kučera & M. Slovák SW6/2* (SAV), *J. Kučera & M. Slovák SW6/3* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.46-3.52 pg, FCM. Switzerland, Graubünden, Waseen village, Süstenpass saddle, $46^{\circ}44'11.4''$ N, $08^{\circ}26'59.5''$ E, 2203 m, 15 Aug 2018, *J. Kučera & M. Slovák SW7/1* (SAV), *J. Kučera & M. Slovák SW7/5* (SAV).

 $2n \sim 4x \sim 40$, 2C = 3.46-3.50 pg, FCM. Switzerland, Graubünden, Silvaplana village, Julierpass saddle, 46°28'13.3"N, 09°43'08.8"E, 2248 m, 16 Aug 2018, *J. Kučera & M. Slovák SW8/1* (SAV), *J. Kučera & M. Slovák SW8/2* (SAV), *J. Kučera & M. Slovák SW8/3* (SAV).

 $2n \sim 4x \sim 40, 2C = 3.47-3.49 \text{ pg}$, FCM. Switzerland, Ticino, St. Gotthard pass saddle, near Lago di San Carlo, 46°33'29.3"N, 08°33'25.5"E, 2116 m, 14 Aug 2018, *J. Kučera & M. Slovák SW3/1* (SAV), *J. Kučera & M. Slovák SW3/2* (SAV), *J. Kučera & M. Slovák SW3/3* (SAV).

 $2n \sim 4x \sim 40, 2C = 3.48-3.52$ pg, FCM. Switzerland, Valais, Wichel village, Nufenen pass saddle, 46°28′20.8″N, 08°22′57.0″E, 2307 m, 14 Aug 2018, *J. Kučera & M. Slovák SW4/7* (SAV), *J. Kučera & M. Slovák SW4/8* (SAV), *J. Kučera & M. Slovák SW4/9* (SAV) [Fig. 7F].

Soldanella ×hybrida A.Kern

The CV of samples and internal standard ranged from 2.05% to 3.01% (mean 2.61%) and from 2.33% to 3.14% (mean 2.71%), respectively.

* 2n = 4x = 40, CHN. Switzerland, Valais, Wichel village, Nufenen pass saddle, 46°28′20.8″N, 08°22′57.0″E, 2307 m, 14 Aug 2018, J. Kučera & M. Slovák SW4/4 (SAV) [Fig. 6F]

* 2n ~ 4x ~ 40, 2C = 3.57–3.62 pg, FCM. Switzerland, Valais, Wichel village, Nufenen pass saddle, 46°28′20.8″N, 08°22′57.0″E, 2307 m, 14 Aug 2018, *J. Kučera & M. Slovák SW4/4* (SAV), *J. Kučera & M. Slovák SW4/5* (SAV), *J. Kučera & M. Slovák SW4/6* (SAV) [Fig. 7E].

Soldanella ×wettsteinii Vierh.

The CV of samples and internal standard ranged from 2.02% to 2.88% (mean 2.52%) and from 2.10% to 3.16% (mean 2.78%), respectively.

* 2n ~ 4x ~ 40, 2C = 3.51–3.56 pg, FCM. Austria, Niederösterreich, Hochschneeberg Mt., Ochsenboden, 47°45′56.0″N, 15°48′36.4″E, 1909 m, 20 Jul 2017, J. Kučera, M. Slovák & E. Štubňová AT40/1 (SAV), J. Kučera, M. Slovák & E. Štubňová AT40/2 (SAV), J. Kučera, M. Slovák & E. Štubňová AT40/3 (SAV), J. Kučera, M. Slovák & E. Štubňová AT40/4 (SAV) [Fig. 7B].

Members of the genus *Soldanella* L. (snowbells, Primulaceae) are perennial, evergreen caespitose herbs with funnel- or bell-shaped flowers and leathery rounded leaves. The genus *Soldanella* is endemic to the European continent and its species occupy understory of mid-altitudinal forests or eventually higher alpine habitats in the large mountain ranges. Following the most recent taxonomic concepts, the genus harbours up to 24 taxa (Zhang & al., 2001; Zhang & Kadereit, 2002).

The karyological variation of the genus, including chromosome numbers and absolute genome size (AGS) was recently investigated by Štubňová & al. (2017). The study confirmed that snowbells are karyologically rather invariable since only two cytotypes, specifically the euploid 2n = 40 and dysploid one with 2n = 38, were detected within the whole genus (Štubňová & al., 2017). Rare aneuploid chromosome numbers reported elsewhere (e.g., Mattick, 1950; Kress, 1969, 1984; Pawłowska, 1963; Peev, 1976; Zhang & Kadereit, 2002; Rice & al., 2015) were, however, not confirmed. Flow cytometry analyses uncovered a limited absolute genome size (AGS) variation in snowbells ranging between 2.97 and 3.99 pg (25.6% overall variation). Beside this, significantly higher AGS (2C = 12.4 pg) has been reported for S. pusilla Baumg. by Veselý & al. (2012). Although authors attempted to analyse all taxa and to cover their distribution range within the study performed by Štubňová & al. (2017), certain parts of the distribution area of several taxa remain unsampled. This was especially a case of S. alpina subsp. alpina and S. pusilla subsp. alpicola from the central and western Alps. Furthermore, the study did not include any of the previously reported hybrids between snowbell species (Zhang & Kadereit, 2002).

The herein presented chromosomal report attempts to fill the gap in the knowledge of karyological variation of the genus with

respect to important but unsampled parts of the distribution area of selected taxa and morphologically intermediate, presumably hybrid individuals. Altogether we analysed 23 populations from the Alps and Carpathians, namely nine populations per S. alpina subsp. alpina, eight populations per S. pusilla subsp. alpicola originating from the south-western Austria, Switzerland, north Italy, and eastern France, one population S. minima s.l. from Austria, three populations of S. angusta from south-eastern part of Ukrainian Carpathians, and the single population of S. major from the Southern Carpathians. Along with pure taxa, we also investigated the karyological variation of two populations with morphologically intermediate, hybrid individuals, namely a single population S. ×hybrida A.Kern (S. alpina subsp. alpina \times S. pusilla subsp. alpicola; Fig. 7D-F) and one population of S. ×wettsteinii Vierh. (S. alpina subsp. alpina × S. minima subsp. austriaca; Fig. 7A-C)). We tested here whether the analysed populations harbour additional (unique) cytotypes beside the ones detected previously and/or possess AGS which would deviate significantly from already gathered and published data.

DIRECT CHROMOSOME COUNTING

Chromosome numbers were determined from the root tips of plants transplanted from the field and cultivated in experimental greenhouses in the Plant Science and Biodiversity Centre of the Slovak Academy of Sciences. The laboratory protocol was as follows: fresh root tips were pre-treated in 0.002 M aqueous solution of hydroxyquinoline overnight; fixed in a fresh solution of 96% ethanol and 100% acetic acid (3:1) for at least 1 hour; rinsed with distilled water; macerated in a mixture of 35% HCl and 96% ethanol (1:1) for about 3 minutes in ambient conditions and finally rinsed with distilled water. Permanent squashes were made using the cellophane squash method (Murín, 1960), stained with a 7% Giemsa solution in Sörensen phosphate buffer (Fluka Analytical, Munich, Germany), washed, dried, mounted, and observed with a high-power oil immersion objective (Mártonfi & al., 1999).

FLOW CYTOMETRY

To estimate the AGS in analysed snowbell populations, we used flow propidium iodide cytometry. All analyses were performed on a flow cytometer CyFlow SL (Partec, Münster, Germany) equipped with a green laser. *Solanum pseudocapsicum* (2C DNA = 2.59 pg, Temsch & al., 2010) was used as the internal standard. Samples were prepared from fresh, intact leaves in a two-step laboratory procedure. First, the sample and standard were co-chopped with a sharp razor blade in a Petri dish in 0.9 to 1 ml of ice-cold Otto I buffer solution and the cell suspension was filtered through a 42-µm nylon mesh. Then, the filtered solution was supplemented with Otto II buffer, propidium iodide as a fluorochrome, RNAse, and β-mercaptoethanol and analysed. For each measurement 5000 nuclei were recorded. To



Fig. 7. Investigated hybrids and their parental species of the genus *Soldanella*. **A**, *S. alpina* (pop. AT40), 2n = 4x = 40; **B**, *S.* ×*wettsteinii* (pop. AT40), $2n \sim 4x$ ~ 40; **C**, *S. minima* subsp. *austriaca* (pop. AT40), 2n = 4x = 40; **D**, *S. alpina* (pop. SW4), 2n = 4x= 40; **E**, *S.* ×*hybrida* (pop. SW4), 2n = 4x = 40; **F**, *S. pusilla* subsp. *alpicola* (pop. SW4), 2n = 4x =40. — Credits to J. Kučera.

avoid potential biases caused by random instrumental drift and/or presence of secondary metabolites we performed following tests: (i) the AGS of each individual was measured independently three times during different days; (ii) the coefficient of variation (CV) of sample and standard were accepted only if they did not exceed 3% in snowbell sample and 5% in standard; (iii) the between-day difference among particular measurements did not exceed critical threshold of 2% otherwise was measurement discarded and sample reanalysed.

NOTES ON KARYOLOGICAL VARIATION OF ANALYSED SOLDANELLA POPULATIONS

Chromosome numbers detected in analysed snowbells were in accordance with all chromosome number records published to date (Bogenrieder, 1972; Chichiricco & Tammaro, 1980; Kress, 1969, 1984; Larsen, 1954; Lippert, 2006; Štubňová & al., 2017). Although we did not obtain a chromosome count for S. ×wettsteinii, the detected AGS values indicate that a shift in chromosome number in this species is improbable. In general, the obtained AGS values fit essentially well to AGS value ranges previously reported for a specific taxon (cf. Štubňová & al., 2017; AGS ranges detected by Štubňová & al., 2017 are in brackets): S. alpina subsp. alpina - 2C = 3.52 - 3.67 pg (2C = 3.47 - 3.65 pg); S. angusta - 2C = 3.31 - 3.38 pg (2C = 3.29 - 3.41 pg); S. major – 2C = 3.28–3.32 pg (2C = 3.35–3.48 pg); S. pusilla subsp. alpicola - 2C = 3.43 - 3.52 pg (2C = 3.44 - 3.49 pg). The exception represent values for S. minima s.l. from southwestern Austria. The detected AGS values, 2C = 3.63-3.68 pg fit to the AGS range previously reported for S. minima subsp. austriaca (2C = 3.50-3.65 pg; Štubňová & al., 2017). However, based on published distribution data this subspecies should not occur in southwestern Austria and its distribution range is shifted more eastwards (Zhang & al., 2001; Zhang & Kadereit, 2002). On the other hand AGS range of S. minima subsp. minima, which is reported from this region were detected to be lower and ranged between 3.39-3.44 pg (Štubňová & al., 2017).

We paid particular attention to potential hybrids between two pairs of higher-alpine snowbell species (Soldanella ×hybrida and Soldanella ×wettsteinii). Although we have no obvious genetic evidence of their hybridity, the analysed plants originated from sympatric populations and their morphological intermediacy was inalienable. Soldanella alpina s.l. represents one of the potential parents for both hybrids and is a typical example of the member of traditionally recognised Soldanella sect. Soldanella. This section is morphologically characterised by the more robust habit, bearing of several (3 to 8) funnel-shaped pink-violet to blue-violet flowers, the presence of floral scales and well-developed stamen appendages. In contrast, another two potentially parental taxa, S. minima, and S. pusilla, are members of traditionally recognised Soldanella sect. Tubiflores Borb. which harbours dwarf plants with reduced floral morphology, i.e., they possess mostly single, bell-shaped, white or pinkish flower, without floral scales and rudimentary stamen appendages (Zhang & al., 2001; Zhang & Kadereit, 2002). Plants which we consider here hybrids were morphologically intermediate and share traits from both parental taxa. In the case of S. ×hybrida, AGS values of the potential hybrid ranged between 2C = 3.57 - 3.62 pg what clearly fits the AGS range detected for S. alpina subsp. alpina - 2C = 3.47 - 3.67. Another potential parent, S. pusilla subsp. alpicola possesses rather smaller AGS - 2C = 3.43-3.49 pg. In contrast, three individuals assignable to S. ×wettsteinii with AGS ranging between 2C = 3.51-3.56 pg perfectly fit AGS variation of both parental taxa (S. minima subsp. austriaca - 2C = 3.50 - 3.65pg and see above for S. alpina subsp. alpina). Although we analysed here the only limited number of morphologically intermediate hybrid

individuals, it seems that hybridisation did not cause significant shifts in AGS of hybrids compared to their parental taxa. The huge AGS values reported for *S. pusilla* (Veselý & al., 2012) were found neither in this taxon nor in whichever of other analysed (sub-)species or hybrids. Thus, the potential scenario involving the whole chromosome doubling and the evolution of allopolyploid entities was not evidenced.

LITERATURE CITED

- **Bogenrieder, A.** 1972. Vergleichende physiologisch-ökologische Untersuchen an Populationen subalpiner Pflanzen aus Schwarzwald und Alpen. Ph.D. dissertation, University of Freiburg, Switzerland.
- Chichiricco, G. & Tammaro, F. 1980. Numeri cromosomici per la flora italiana. *Inform. Bot. Ital.* 12: 161–165.
- Kress, A. 1969. Zytotaxonomische Untersuchungen an Primulaceae. *Phyton (Horn)* 13: 211–225.
- Kress, A. 1984. Chromosomenzählungen an verschiedenen Primulaceen: Soldanella. Primulaceen Studien 7, Teil B: 1–25.
- Larsen, K. 1954. Chromosome numbers of some European flowering plants. *Bot. Tidsskr.* 50: 163–174.
- Lippert, W. 2006. Chromosomenzahlen von Pflanzen aus Bayern und anderen Gebieten. *Ber. Bayer. Bot. Ges.* 76: 85–110.
- Mártonfi, P., Michálek, J., Hadinec, J., Mártonfiová, L. & Repčák, M. 1999. *Hypericum dubium* – A new species of the Czech flora. *Preslia* 71: 337–348.
- Mattick, F. 1950. Soldanella. P. 85 in: Tischler, G. (ed.), Die Chromosomenzahlen der Gefäßpflanzen Mitteleuropas. s'Gravenhage: Springer.
- Murín, A. 1960. Substitution of cellophane for glass covers to facilitate preparation of permanent squashes and smears. *Stain Tech*nol. 35: 351–353.
- Pawłowska, S. 1963. O polnocno karpackich gatunkach rodzaju Soldanella L. – De Soldanella quae in parte septentrionali Carpatorum crescunt. Fragm. Florist. Geobot. 9: 3–30.
- Peev, D. 1976. Chromosome numbers and critical notes on taxonomy of some Bulgarian Primulaceae species. *Fitologija* 4: 16–24.
- Rice, A., Glick, L., Abadi, S., Einhorn, M., Kopelman, N. M., Salman-Minkov, A., Mayzel, J., Chay, O. & Mayrose, I. 2015. The chromosome Counts Database (CCDB) – A community resource of plant chromosome numbers. *New Phytol.* 206: 19–26. https://doi.org/10.1111/nph.13191
- Štubňová, E., Hodálová, I., Kučera, J., Mártonfiová, L., Svitok, M. & Slovák, M. 2017. Karyological pattern in the European endemic genus *Soldanella* L.: Absolute genome size variation uncorrelated with cytotype chromosome numbers. *Amer. J. Bot.* 104: 1241–1253. https://doi.org/10.3732/ajb.1700153
- Temsch, E.M., Greilhuber, J. & Krisai, R. 2010. Genome size in liverworts. *Preslia* 82: 63–80.
- Veselý, P., Bureš, P., Šmarda, P. & Pavlíček, T. 2012. Genome size and DNA base composition of geophytes: The mirror of phenology and ecology? *Ann. Bot. (Oxford)* 109: 65–75. https://doi. org/10.1093/aob/mcr267
- Zhang, L.B. & Kadereit, J.W. 2002. The systematic of *Soldanella* L. (Primulaceae) based on morphological and molecular (ITS, AFLPs) evidence. *Nordic J. Bot.* 22: 129–169. https://doi. org/10.1111/j.1756-1051.2002.tb01360.x
- Zhang, L.B., Comes, H.P. & Kadereit, J.W. 2001. Phylogeny and quaternary history of the European montane/alpine endemic *Soldanella* (Primulaceae) based on ITS and AFLP variation. *Amer. J. Bot.* 88: 2331–2345.

Rohit Kumar, Vandna Kumari & Vijay Kumar Singhal*

Department of Botany, Punjabi University, Patiala-147002, Punjab, India

* Author for correspondence: vksinghal53@gmail.com

The study was supported by financial grant under DRS SAP I, SAP II and SAP III of University Grant Commision (UGC), New Delhi, India; Women Scientist Fellowship Scheme (WOS-A) of Department of Science and Technology, Government of India (GOI) to Vandna Kumari, Rajiv Gandhi National Fellowship Scheme of UGC to Rohit Kumar (award letter no. F1-17.1/2012-13 /RGNF-2012-13-SC HIM-18573) and IPLS project of Department of Biotechnology, GOI (BT/PR4548/INF/22/146/ 2012) sanctioned to Punjabi University Patiala, Punjab for using the facilities and financial support of this study.

- * First chromosome count for the species.
- ** New chromosome number (cytotype) for the species.
- ▼ First chromosome count for an Indian accession.

BRASSICACEAE

** Turritis glabra L.

n = 7, CHN. India, Uttarakhand, Uttarkashi, Gangotri-Gomukh Trek, 30°59'17.68"N, 79°00'17.57"E, 3500 m, found in open stony, partly shaded slopes, *R. Kumar 34335* (PUN 61593) [Fig. 8A].

The other chromosome number reports for this species are 2n = 12 (Titz, 1967; Taylor & Mulligan, 1968; Rodman & Bhargava, 1976; Arohonka, 1982; Dobeš & al., 1997; Koch & al., 1999; Lövkvist & Hultgård, 1999; Kumar & Singhal, 2011), 2n = 16 (Titz, 1968; Hill, 1982) and 2n = 16, 32 (Titz, 1966).

CARYOPHYLLACEAE

* Arenaria festucoides Benth.

n = 11, CHN. India, Uttarakhand, Uttarkashi, Nelang, 31°06'38.08"N, 78°59'59.42"E, 3500 m, found in moist, open or partly shaded places along streams in meadows, *Rohit Kumar 34336* (PUN 61594) [Fig. 8B].

FABACEAE

**Astragalus uttaranchalensis* L.B.Chowdhary & Z.H.Khan n = 8, CHN. India, Uttarakhand, Uttarkashi, Gangotri, 30°59′42.61″N, 78°56′24.04″E, 3500 m, rarely found in open dry slopes in particular areas, *R. Kumar* 34862 (PUN 61720) [Fig. 8C].

▼ Medicago edgeworthii Širj.

n = 8, CHN. India, Uttarakhand, Uttarkashi, Nelang, 31°06'38.08"N, 78°59'59.42"E, 3500 m, grows as isolated individuals in open and partly shaded slopes, *Rohit Kumar 34903* (PUN 61761) [Fig. 8D].

The current chromosome number report is in agreement with the previous one, 2n = 16 (Zhou & al., 2000).

▼ *Thermopsis barbata* Benth.

n = 9, CHN. India, Uttarakhand, Uttarkashi, Har-ki-Dun, 31°08'30.76"N, 78°24'37.43"E, 3600 m, grows as isolated individuals in open grassy slopes, *R. Kumar 34916* (PUN 61774) [Fig. 8E].

Diploid chromosome count of 2n = 18 for this species has been previously reported by Chen & al. (1992).

HYPERICACEAE

***Hypericum elodeoides* Choisy

n = 10, CHN. India, Uttarakhand, Uttarkashi, Taluka, 31°04′59.83″N, 78°15′31.50″E, 2100 m, found in along roadsides, rocky slopes, damp meadows and grassy grounds, *R. Kumar 34358* (PUN 61616) [Fig. 8G].

There are several different chromosome number reports for this species, e.g., 2n = 16 (Suguira, 1941, 1944; Al-Bermani & al., 1993; Kumar, 2012), 2n = 18 (Gupta & al., 2014; Kumar & Singhal, 2011; Jeelani & al., 2014; Kaur, 2017) and 2n = 32 (Sandhu & Mann, 1989; Kumar, 2012).

PAPAVERACEAE

* Corydalis stracheyi Duthie ex Prain

n = 8, CHN. India, Uttarakhand, Uttarkashi, Dodi Tal, 30°53'50.09"N, 78°31'38.74"E, 3050 m, found in moist grassy slopes along streams in meadows, *R. Kumar 34310* (PUN 61568) [Fig. 8F].

POACEAE

** Digitaria abludens (Roem. & Schult.) Veldkamp

n = 36, CHN. India, Himachal Pradesh, Kullu District, Sheela Village, 32°00′23″N, 77°22′44″E, 2650 m, found in weedy places and roadsides, *V. Kumari 32435* (PUN 60309) [Fig. 8H].

Previous chromosome number reports for this species from the Indian Himalayas were 2n = 36 (Sharma & Sharma, 1979; Mehra, 1982).

** Festuca sibirica Hack. ex Boiss.

n = 21, CHN. India, Himachal Pradesh, Kullu District, Gulaba, $32^{\circ}20'27''N$, $77^{\circ}13'06''E$, 3200 m, found on stony slopes, among rocks and sometimes in sandy areas, *V. Kumari* 34676 (PUN 61325) [Fig. 81].

Previous reports for the species were 2n = 28 recorded by (Chepinoga & al., 2008; Probatova & al., 2012).

▼ *Hordeum murinum* L.

n = 7, CHN. India, Himachal Pradesh, Kullu District, Jalori Pass, 31°32′13″N, 77°22′23″E, 3223 m, found in fields and along roadsides, *V. Kumari 34705* (PUN 61254) [Fig. 8J].

The current chromosome number report is in agreement with the previous ones of 2n = 14 as reported by Morrison (1959), Strid & Franzén (1981), Faruqi & al. (1987) and Nazarova (2004).

**Leptochloa chinensis (L.) Nees

n = 9, CHN. India, Himachal Pradesh, Kullu District, Bhuntar, 31°53′02″N, 77°08′42″E, 1100 m, found in the moist places, *V. Kumari 34712* (PUN 61261) [Fig. 8K].

Several other chromosome numbers were reported for this species, namely 2n = 36 (Mehra & Kalia, 1975), 2n = 40 (Avdulov, 1928; Chen & Hsu, 1962; Mehra, 1982; Bir & Sahni, 1986) and 2n = 54 (Christopher & Abraham, 1974).

** Microstegium vimineum (Trin.) A.Camus

n = 10, CHN. India, Tosh, 32°03′08″N, 77°26′58″E, 2400 m, found in forest margins, moist grassy places, *V. Kumari 34711* (PUN 61260) [Fig. 8L].

Previous reports for the species is 2n = 40 (Tateoka, 1967; Mehra & Kalia, 1976; Mehra, 1982). The present report of 2n = 20 adds a new diploid cytotype for the species at world level.



Fig. 8. Meiosis. A, Turritis glabra, PMC showing 7 bivalents at metaphase-I (M-I), n = 7 (PUN 61593); **B**, Arenaria festucoides, PMC showing 11 bivalents at diakinesis, n = 11 (PUN 61594); **C**, Astragalus uttaranchalensis, PMC showing 8 bivalents at M-I, n = 8 (PUN 61720); D, Medicago edgeworthii, PMC showing 8 bivalents at M-I, n = 8 (PUN 61761); E, Thermopsis barbata, PMC showing 9 bivalents at M-I, n = 9 (PUN 61774); F, Corydalis stracheyi, PMC showing 8 bivalents at M-I, n = 8 (PUN 61568); **G**, *Hypericum elodeoides*, PMC showing 10 bivalents at M-I, n = 10 (PUN 61616); H, Digitaria abludens, PMC showing equal distribution of 36:36 chromosomes at anaphase-I (A-I), n = 36 (PUN 60309); I, Festuca sibirica, PMC showing 21 bivalents at diakinesis, n = 21 (PUN 61325); J, Hordeum murinum, PMC showing 7 bivalents (4 bivalents in groups of two each + 3 bivalents) at M-I, n = 7(PUN 61254); K, Leptochloa chinensis, PMC showing 9 bivalents at M-I, n = 9(PUN 61261); L, Microstegium vimineum, PMC showing 10 bivalents at M-I, n = 10(PUN 61260); M, Poa setulosa, PMC showing 14 bivalents (3 large-sized and 11 normal-sized) at diakinesis n = 14(PUN 61188); N, Puccinellia himalaica, PMC showing 14 bivalents (2 large-sized ring-shaped and 12 normal-sized) at M-I, n = 14 (PUN 61238); **0**, *Stipa jacque*montii, PMC showing equal distribution of 10:10 chromosomes at A-I, n = 10(PUN 61243); P, Saxifraga pedemontana subsp. cymosa, PMC showing 8 bivalents at M-I, *n* = 8 (PUN 61854); **Q**, *Myricaria* elegans, PMC showing 12 bivalents at M-I, n = 12 (PUN 61862); **R**, *M*. rosea, PMC showing 12 bivalents at diakinesis, n = 12 (PUN 61863). — Scale bars = 10 µm.

**Poa setulosa Bor

n = 14, CHN. India, Himachal Pradesh, Kullu, Kothi, 32°18'52.77"N, 77°11'24.76"E, 2500 m, found on stony slopes, among rocks and sometimes in sandy areas, *V. Kumari 34740* (PUN 61188) [Fig. 8M].

Previous chromosome number report for this species was 2n = 14 (Kaur & al., 2011).

*Puccinellia himalaica Tzvelev

n = 14, CHN. India, Himachal Pradesh, Kullu District, Jalori Pass, 31°32′13″N, 77°22′23″E, 3223 m, found in moist and sandy places, *V. Kumari 34689* (PUN 61238) [Fig. 8N].

**Stipa jacquemontii Jaub. & Spach

n = 10, CHN. India, Himachal Pradesh, Kullu, Bijali Mahadev, 31°55′30″N, 77°09′01″E, 2435 m, found in arid soils and rock crevices, *V. Kumari 34694* (PUN 61243) [Fig. 80].

Previously different chromosome numbers were reported for this species: 2n = 24 (Mehra & Sharma, 1975) and 2n = 42 (Gupta & al., 2014).

SAXIFRAGACEAE

* Saxifraga pedemontana subsp. cymosa Engl.

n = 8, CHN. India, Uttarakhand, Uttarkashi, Har-ki-Dun Trek, 31°08'10.67"N, 78°23'58.06"E, 3460 m, rarely found in grassy slopes in meadows, *R. Kumar 34981* (PUN 61854) [Fig. 8P].

TAMARICACEAE

*Myricaria elegans Royle

n = 12, CHN. India, Uttarakhand, Uttarkashi, Chirwasa, 31°08'10.67"N, 78°23'58.06"E, 3560 m, grows in stony or boulderstrewn places along river beds or streams and open dry slopes, *R. Kumar 34988* (PUN 61862) [Fig. 8Q].

*Myricaria rosea W.W.Sm.

n = 12, CHN. India, Uttarakhand, Uttarkashi, Har-ki-Dun, 31°08′30.76″N, 78°24′37.43″E, 3600 m, rarely found in open dry, boulder-strewn slopes, along river banks, *R. Kumar 34990* (PUN 61863) [Fig. 8R].

LITERATURE CITED

- Al-Bermani, A.K., Al-Shammary, K.I.A, Bailey, J.P. & Gornall, R.J. 1993. Contribution to a cytological catalogue of the British and Irish flora, 3. *Watsonia* 19: 169–171.
- Arohonka, T. 1982. Chromosome counts of vascular plants of the island Seili in Nauvo, southwestern Finland. *Turun Yliopiston* Julkaisuja, Sar. A 2, Biol. Geogr. 3: 1–12.
- Avdulov, N.P. 1928. Sistematicheskaya kariologiya semeistva Gramineae. Pp. 65–67 in: Borodin, I.P. & Bush N.A. (eds.), Dnevnik Vsesojuznogo s'ezda botanikov v Leningrade v janvare 1928 goda, vol. 1. Leningrad: Russkoe Botanicheskoe Obshchestvo.
- Bir, S.S. & Sahni, M. 1986. SOCGI plant chromosome number report IV. J. Cytol. Genet. 21: 152–154.
- Chen, C.C. & Hsu, C.C. 1962. Cytological studies on Taiwan grasses II. Chromosome numbers of miscellaneous tribes. J. Jap. Bot. 37: 303–313.
- Chen, C.J., Zhu, X.Y. & Yuan, Y.M. 1992. Cytological studies on the tribe Thermopsideae (Fabaceae) I. Report on karyotypes of eleven species of four genera. *Cathaya* 4: 103–116.

Chepinoga, V.V., Gnutikov, A.A., Enushchenko, I.V. & Chepinoga,

A.V. 2008. [Report] in: Marhold, K. (ed.), IAPT/IOPB chromosome data 6. *Taxon* 57: 1267–1268.

- Christopher, J. & Abraham, A. 1974. Studies on the cytology and phylogeny of South Indian grasses. II. Sub-family Eragrostoideae. *Cytologia* 39: 561–571. https://doi.org/10.1508/cytologia.39.561
- Dobes, C., Hahn, B. & Morawetz, W. 1997. Chromosome numbers for the vascular plant flora of Austria. *Linzer Biol. Beitr*. 29: 5–43.
- Faruqi, S.A., Quraish, H.B. & Inamuddin, M. 1987. Studies in Libyan grasses X. Chromosome number and some interesting features. Ann. Bot. (Oxford) 45: 75–102.
- Gupta, R.C., Chauhan, H.S., Saggoo, M.I.S. & Kaur, N. 2014. Cytomorphology of some grasses (Poaceae) from Lahaul-Spiti (Himachal Pradesh), India. *Biolife* 2: 1234–1247.
- Hill, L.M. 1982. [Report]. In: Löve, Á. (ed.), IOPB chromosome number reports. LXXIV. *Taxon* 31: 119–128.
- Jeelani, S.M., Kumari, S., Gupta, R.C. & Siddique, M.A.A. 2014. Detailed cytomorphological investigations through male meiosis of polypetalous plants from the Kashmir Himalaya. *Pl. Syst. Evol.* 300: 1175–1198. https://doi.org/10.1007/s00606-013-0954-z
- Kaur, H., Gupta, R.C. & Kumari, S. 2011. [Reports] in: Marhold, K. (ed.), IAPT/IOPB chromosome data 12. *Taxon* 60: 1784–1796.
- Kaur, M. 2017. Evaluation of cytomorphological diversity in the dicots from Solang Valley (Himachal Pradesh). Ph.D. Thesis, Department of Botany, Punjabi University Patiala, India.
- Koch, M., Bishop, J. & Mitchell-Olds, T. 1999. Molecular systematics and evolution of *Arabidopsis* and *Arabis*. *Pl. Biol.* 1: 529–537. https://doi.org/10.1111/j.1438-8677.1999.tb00779.x
- Kumar, P. & Singhal, V.K. 2011. Chromosome number, male meiosis and pollen fertility in selected angiosperms of the cold deserts of Lahaul-Spiti and adjoining areas (Himachal Pradesh, India). *Pl. Syst. Evol.* 297: 271–297. https://doi.org/10.1007/s00606-011-0516-1
- Kumar, S. 2012. Exploration and evaluation of cytomorphological diversity in Polypetalae from district Sirmaur (Himachal Pradesh). Ph.D. Thesis. Punjabi University Patiala, India.
- Lövkvist, B. & Hultgård, U.M. 1999. Chromosome numbers in the South Swedish vascular plants. Opera Bot. 137: 1–42.
- Mehra, P.N. 1982. *Cytology of East Indian grasses*. New Delhi: Pramodh P. Kapur at Rajbandhu Industrial Company.
- Mehra, P.N. & Kalia, V. 1975. [Report]. In: Löve, A. (ed.), IOPB chromosome number reports. XLIX. *Taxon* 24: 501–516.
- Mehra, P.N. & Kalia, V. 1976. [Report]. In: Löve, Á. (ed.), IOPB chromosome number reports LIV. *Taxon* 25: 632.
- Mehra, P.N. & Sharma, M.L. 1975. Cytological studies in some Central and Eastern Himalayan grasses. III. The Agrostideae, Aveneae, Brachypodieae, Bromeae, Festuceae, Phalarideae and Triticeae. Cytologia 40: 441–452. https://doi.org/10.1508/cytologia.40.441
- Morrison, J.W. 1959. Cytogenetic studies in the genus *Hordeum*. I. Chromosome morphology. *Canad. J. Bot.* 37: 27–538. https://doi. org/10.1139/b59-046
- Nazarova, E.A. 2004. Karyological characterization of Armenian species of the genus *Hordeum* L. *Fl. Rastitel'nost' Rastitel'nye Resursy Armyansk. S.S.R.* 15: 57–63, 91–95.
- Probatova, N.S., Kazanovsky, S.G., Rudyka, E.G., Seledets, V.P. & Nechaev, V.A. 2012. [Report] in: Marhold, K. (ed.), IAPT/ IOPB Chromosome data 13. *Taxon* 61: 899–901.
- Rodman, J.E. & Bhargava, M. 1976. [Report]. In: Löve, Á. (ed.), IOPB chromosome number reports. LIII. *Taxon* 25: 483–500.

- Sandhu, P.S. & Mann, S.K. 1989. SOCGI plant chromosome number reports. VIII. J. Cytol. Genet. 24: 221–228.
- Sharma, M.L. & Sharma, K. 1979. Cytological studies in the North Indian grasses. *Cytologia* 44: 861–872. https://doi.org/10.1508/ cytologia.44.861
- Strid, A. & Franzen, R. 1981. [Report]. In: Löve, A. (ed.), IOPB Chromosome number reports LXXIII. *Taxon* 30: 829–842.
- Suguira, T. 1941. A list of chromosome numbers in angiospermous plants. VIII. Proc. Imp. Acad. Japan 17: 29–30. https://doi. org/10.2183/pjab1912.17.29
- Suguira, T. 1942. Studies on chromosome numbers in higher plants VI. Cytologia 13: 352–359. https://doi.org/10.1508/cytologia.13.352
- Tateoka, T. 1967. [Report]. In: Löve, Á. (ed.), IOPB chromosome number reports XIV. *Taxon* 16: 561–564.
- Taylor, R.L. & Mulligan, G.A. 1968. Flora of the Queen Charlotte Islands. Part 2. Cytological aspects of the vascular plants. *Monogr. Res. Branch Canada Dept. Agric.* 4(2): 1–148.
- Titz, W. 1966. New Austrian sites of Agropyron, Bromus and Arabis species as well as their chromosome numbers. Österr. Bot. Z. 113: 470–475. https://doi.org/10.1007/BF01373449
- Titz, W. 1967. Zur Cytologie und Systematik einiger österreichischer Arabis-Arten (einschliesslich Turlitis glabra L.). Ber. Deutsch Bot. Ges. 79: 474–488.
- Titz, W. 1968. Chromosomenzahlar europäischer Arabis-Arten (Cruciferae). Ber. Deutsch. Bot. Ges. 81: 217–220.
- Zhou, Z.W., Xue, S.M., Kui, J.X. & Wang, L. 2000. Karyotype analysis of *Melilotoides pubescens*. Grassl. China 5: 76–77, 79.

M. Laura Las Peñas

Instituto Multidisciplinario de Biología Vegetal (IMBIV), CONICET-UNC, Universidad Nacional de Córdoba, CC 495 Córdoba 5000, Argentina; laulaspenas@yahoo.com.ar

Funding was provided by CONICET, FONCyT, and SECyT-UNC (FCEFyN) from Argentina.

* First chromosome counts from the given regions.

CACTACEAE

**Arrojadoa rhodantha* (Gürke) Britton & Rose 2*n* = 22, *x* = 11, CHN. Brasil, Rio de Contas, 13°38′09″S, 41°39′08″W, 15 Mar 2007, *Urdampilleta & al. 379* (CORD) [Fig. 9A].

Cereus aethiops Haw.

2*n* = 22, *x* = 11, CHN. Argentina, Mendoza, Dpt. Lujan de Cuyo, 33°01′01″S, 68°11′07″W, 29 Jan 2006, *Chiarini 614* (CORD) [Fig. 10A].

Cleistocactus smaragdiflorus (F.A.C.Weber) Britton & Rose 2n = 22, x = 11, CHN. Argentina, Salta, Dpt. Guachipas, 25°32'14"S, 65°31'10"W, 15 Jan 2006, *Las Peñas & Uñates 279* (CORD) [Fig. 10B].



Fig. 9. FISH of 18S-5.8S-26S and 5S rDNA to metaphase chromosomes in Cactaceae simultaneously detected in all species. A, Arrojadoa rhodantha; B, Melocactus bahiensis; C, Pachycereus pringlei; D, Tacinga inamoena. — Scale bar = 5 μ m (all of the images are at the same scale).



Fig. 10. Photomicrographs of mitotic metaphases of Cactaceae species. A, Cereus aethiops; B, Cleistocactus smaragdiflorus; C, Denmoza rhodacantha; D, Echinopsis aurea var. dobeana; E, Echinopsis leucantha; F, Gymnocalycium monvillei subsp. achirasense; G, Harrisia pomanensis;
H, Harrisia hahniana; I, Lobivia saltensis; J, Maihueniopsis glomerata; K, Maihueniopsis darwinii; L, Maihueniopsis ovata; M, Oreocereus trollii;
N, Rhipsalis lumbricoides; O, Trichocereus arboricola; P, Trichocereus atacamensis; Q, Trichocereus thelegonus; R, Trichocereus macrogonus;
S, Pachycereus pringlei; T, Quiabentia verticillata. — Scale bar = 2 μm (all of the images are at the same scale).

Denmoza rhodacantha (Salm-Dyck) Britton & Rose

2n = 22, x = 11, CHN. Argentina, La Rioja, Dpt. Chilecito, 34°00′00″S, 68°54′22″W, 5 Dec 2005, *Las Peñas & Chiarini 212* (CORD) [Fig. 10C].

Echinopsis aurea var. dobeana (Dölz) J.Ullmann

2*n* = 22, *x* = 11, CHN. Argentina, Catamarca, Dpt. Ancasti, 28°48'13"S, 68°30'42"W, 30 Nov 2004, *Las Peñas 125* (CORD) [Fig. 10D].

Echinopsis aurea var. falax (Oehme) J.Ullmann

2n = 22, x = 11, CHN. Argentina, La Rioja, Dpt. Independencia, $28^{\circ}21'19''S$, $64^{\circ}11'08''W$, 5 Dec 2005, *Las Peñas & Chiarini 203* (CORD).

Echinopsis leucantha (Gillies ex Salm-Dyck) Walp.

2n = 22, x = 11, CHN. Argentina, San Juan, Dpt. Albardón, 31°27'58"S, 68°38'08"W, 14 Mar 2004, *Las Peñas & Uñates 87* (CORD) [Fig. 10E].

*Gymnocalycium andreae (Boed.) Backeb.

2*n* = 44, *x* = 11, CHN. Argentina, Córdoba, Dpt. San Alberto, 31°23′42″S, 64°46′48″W, 25 Jan 2004, *Las Peñas & Uñates 23* (CORD).

**Gymnocalycium monvillei* subsp. *achirasense* (H.Till & Schatzl) H.Till

2*n* = 22, *x* = 11, CHN. Argentina, Cordoba, Dpt. Rió Cuarto, 32°52′21″S, 64°53′11″W, 26 Aug 2007, *Las Peñas & al. 363* (CORD) [Fig. 10F].

* *Gymnocalycium monvillei* (Lem.) Britton & Rose subsp. *monvillei*

2n = 44, x = 11, CHN. Argentina, Córdoba, Dpt. San Alberto, $31^{\circ}23'41''S$, $64^{\circ}46'48''W$, 26 Aug 2007, *Las Peñas & Uñates 232* (CORD).

Harrisia hahniana (Backeb.) Kimnach & Hutchison ex Kimnach 2n = 22, x = 11, CHN. Paraguay, Dpt. Presidente Hayes, Irala Fernandez, 22°48′49.3″S, 59°34′25.3″W, 10 Oct 2014, *Las Peñas 588* (CORD) [Fig. 10H].

Harrisia pomanensis (F.A.C.Weber ex K.Schum.) Britton & Rose 2n = 44, x = 11, CHN. Argentina, San Luis, Dpt. Ayacucho, 32°14′52″S, 65°46′11″W, 30 Nov 2004, Las Peñas & Uñates 100 (CORD) [Fig. 10G].

Lobivia saltensis (Speg.) Britton & Rose

2n = 22, x = 11, CHN. Argentina, Salta, Dpt. Cafayate, 26°05'54.77"S, 65°57'02.50"W, 10 Dec 2004, *Las Peñas138* (CORD) [Fig. 101].

Maihueniopsis darwinii (Hensl.) F.Ritter

2*n* = 55, *x* = 11, CHN. Argentina, Chubut, Dpt. Biedma, 42°45′57″S, 64°08′46″W, 30 Jan 2003, *Las Peñas & Uñates 349* (CORD) [Fig. 10K].

Maihueniopsis glomerata (Haw.) R.Kiesling

2n = 44, x = 11, CHN. Argentina San Juan, Dpt. Iglesia, 29°45′57″S, 69°26′16.77″W, 16 Apr 2005, *Las Peñas & Uñates 178* (CORD) [Fig. 10J].

Maihueniopsis ovata (Pfeiff.) F.Ritter

2n = 44, x = 11, CHN. Argentina San Juan, Dpt. Iglesia, 29°45′57″S, 69°26′16.77″W, 16 Apr 2005, *Las Peñas & Uñates 181* (CORD) [Fig. 10L].

**Melocactus bahiensis* (Britton & Rose) Luetzelb. 2*n* = 44, *x* = 11, CHN. Brasil, Rio de Contas, 13°20'19"S, 41°29'52"W, 15 Mar 2007, *Urdampilleta & al. 387* (CORD) [Fig. 9B].

Opuntia quimilo K.Schum. 2n = 22, x = 11, CHN. Argentina, La Rioja, Dpt. Chamical, $30^{\circ}09'14''S$, $66^{\circ}28'44''W$, 18 Sep 2012, Las Peñas & al. 539 (CORD).

Oreocereus celsianus (Lem. ex Salm-Dyck) Riccob. 2n = 44, x = 11, CHN. Argentina, Jujuy, Dpt. Tumbaya, 23°51′28″S, 65°28′08″W, 15 Jan 2006, *Las Peñas & Uñates 274* (CORD).

Oreocereus trollii (Kupper) Backeb. 2n = 22, x = 11, CHN. Argentina, Jujuy, Dpt. Tumbaya, 23°51′28″S, 65°28′08″W, 15 Jan 2006, Las Peñas & Uñates 339 (CORD) [Fig. 10M].

**Pachycereus pringlei* (S.Watson) Britton & Rose 2*n* = 22, *x* = 11, CHN. México, Dpt. La Paz, 24°01′26″S, 110°33′39″W, 10 Sep 2015, *Las Peñas 714* (CORD) [Fig. 9C, Fig. 10S].

Quiabentia verticillata (Vaupel) Vaupel ex A.Berger 2*n* = ca. 187, *x* = 11, CHN. Bolivia, Dpt. Mizque, 56°16'04"S, 65°18'39"W, 28 Apr 2011, *Urdampilleta & al. 542* (CORD) [Fig. 10T].

**Rhipsalis lumbricoides* (Lem.) Lem. ex Salm-Dyck 2*n* = 22, *x* = 11, CHN. Argentina, Misiones, Dpt. Posadas, 14 Apr 2004, *Las Peñas & Uñates 56* (CORD) [Fig. 10N].

Tacinga inamoena (K.Schum.) N.P.Taylor & Stuppy 2n = 44, x = 11, CHN. Brasil, Rio de Contas, 13°49'53"S, 41°35'05"W, 15 Mar 2007, *Urdampilleta & al. 385* (CORD) [Fig. 9D].

Trichocereus arboricola Kimnach.

2n = 22, x = 11, CHN. Argentina, Salta, Dpt. Capital, El Baritú, 24°55'32"S, 65°38'30"W, 15 Mar 2017, *Las Peñas & Chiarini 715* (CORD) [Fig. 100].

* *Trichocereus atacamensis* (Phil.) W.T.Marshall & T.M.Bock 2*n* = 22, *x* = 11, CHN. Argentina, Salta, Dpt. Cachi, 25°05'57"S, 66°16'44"W, 15 Jan 2006, *Las Peñas & al. 319* (CORD) [Fig. 10P].

Trichocereus macrogonus Salm-Dyck) Riccob. 2*n* = 22, *x* = 11, CHN. Perú, Dpt. La Paz, 16°28′09″S, 67°55′58″W, 25 Dec 2005, *Las Peñas 122* (CORD) [Fig. 10R].

Trichocereus thelegonus (F.A.C.Weber) Britton & Rose 2*n* = 22, *x* = 11, CHN. Argentina, Salta, 25°38′11″S, 65°39′11″W, 15 Jan 2006, *Las Peñas & al. 283* (CORD) [Fig. 10Q].

The chromosome numbers of 28 and karyotype features of 17 taxa of Cactaceae are presented. Preparation of metaphase chromosomes was done from adventitious root tips pretreated with 2 mM 8-hydroxyquinoline for 24 h at 4°C and fixed in 3:1 ethanol: acetic acid, washed in distilled water, digested 45 min at 37° C with Pectinex SP ULTRA (Novozymes, Villa Regina, Rio Negro, Argentina),

and squashed in a drop of 45% acetic acid. Slides were stained with Giemsa. At least ten metaphases of each species were photographed with phase contrast in a Zeiss Axiophot microscope. Photographs were used to take the following measurements for each chromosome pair: s (short arm), l (long arm), and c (average total chromosome length). The arm ratio (r = 1/s) was then calculated and used to classify the chromosomes as recognized by Levan & al. (1964). In addition, total haploid chromosome length of the karyotype (TL) based on the mean chromosome lengths was calculated. Karyotype asymmetry was estimated using Romero Zarco's (1986) indices ($A_1 = intrachro$ mosomal asymmetry index, and $A_2 =$ interchromosomal asymmetry index). Figure 10 shows the range of chromosomes morphologies found. Idiograms representing the haploid complement were based on the mean values for each species. In situ hybridization followed Schwarzacher & Heslop-Harrison (2000) with modifications. For details of elaboration and labelling of rDNA probes see Las Peñas & al. (2017).

Table 1. Chromosomal data of	the Cactaceae	species studied
------------------------------	---------------	-----------------

Our chromosome counts are the first report for 18 species and 2 varieties, while the remaining counts confirm previously published data. Chromosome features are summarized in Table 1. Almost all species were diploid, with 2n = 22, except for 11 taxa which exhibited polyploidy with 2n = 44, 55 and 187 ca. (Table 1, Fig. 10). Our results confirmed that the basic chromosome number for Cactaceae is x = 11(Pinkava, 2002; Goldblatt & Johnson, 2013; Las Peñas & al., 2017). Polyploidy has been reported for ca. 25% of the cacti investigated so far (Pinkava, 2002; Las Peñas & al., 2009); particularly, it seems to have played an important role in the evolution of Opuntioideae where it reaches a percentage of 64% (Pinkava, 2002; Las Peñas & al., 2017; our data). The chromosomes were small, 2.51 µm being the average chromosome length for all taxa, and species analyzed here fall within the range expected for the family (Las Peñas & al., 2009, 2014). The karyotypes of 17 taxa studied are reported for the first time. Karyotypes were symmetrical, considering both centromere position (most are m with 1 or 2 sm pairs). Subtelocentric (st) chromosomes are

Taxa	2 <i>n</i>	KF	TL	с	r	A ₁	A ₂
Arrojadoa rhodantha	22*	10 m + 1 sm	25.40	2.30	1.21	0.20	0.18
Cereus aethiops	22	10 m + 1 sm	27.37	2.51	1.31	0.22	0.19
Cleistocactus smaragdiflorus	22	10 m + 1 sm	28.70	2.54	1.40	0.20	0.21
Denmoza rhodacantha	22	9 m + 1 sm + 1 st	19.93	1.81	1.29	0.19	0.09
Echinopsis aurea var. dobeana	22	10 m + 1 sm	26.90	2.45	1.29	0.2	0.24
Echinopsis aurea var. falax	22	10 m + 1 sm	26.70	2.30	1.20	0.19	0.21
Echinopsis leucantha	22	10 m + 1 sm	27.65	2.55	1.29	0.22	0.12
Gymnocalycium andreae	44*	_	_	_	_	_	-
Gymnocalycium monvillei subsp. achirasense	22*	10 <i>m</i> +1 <i>sm</i>	34.66	3.15	1.22	0.17	0.07
Gymnocalycium monvillei subsp. monvillei	44*	_	_	_	_	_	-
Harrisia hahniana	22	10 m + 1 sm	22.73	2.06	1.25	0.18	0.17
Harrisia pomanensis	44	_	_	_	_	_	-
Lobivia saltensis	22	10 <i>m</i> +1 <i>sm</i>	37.25	3.33	1.24	0.17	0.15
Maihueniopsis darwinii	55	_	_	_	_	_	_
Maihueniopsis glomerata	44	_	_	_	_	_	-
Maihueniopsis ovata	44	_	_	_	_	_	-
Melocactus bahiensis	44*	_	_	-	_	_	-
Opuntia quimilo	22*	11 <i>m</i>	23.95	2.17	1.21	0.15	0.09
Oreocereus celsianus	44	_	_	_	_	_	-
Oreocereus trollii	22	10 m + 1 sm	31.25	2.80	1.20	0.18	0.19
Pachycereus pringlei	44	_	_	_	_	_	-
Quiabentia verticillata	187 ca.	_	_	_	_	_	_
Rhipsalis lumbricoides	22*	10 m + 1 sm	24.25	2.20	1.42	0.28	0.17
Tacinga inamoena	44	_	_	_	_	_	-
Trichocereus arboricola	22	9 m + 2 sm	38.58	2.95	1.48	0.26	0.28
Trichocereus atacamensis	22*	9 m + 2 sm	28.15	2.58	1.42	0.25	0.24
Trichocereus macrogonus	22	10 <i>m</i> +1 <i>sm</i>	26.55	2.41	1.42	0.28	0.17
Trichocereus thelegonus	22	9 m + 2 sm	28.46	2.58	1.26	0.18	0.12

2n = somatic chromosome number; KF = karyotype formula; TL = total haploid chromosome length of the karyotype in μ m; c = average total chromosome length in μ m; A₁ = intrachromosomal asymmetry index; A₂ = interchromosomal asymmetry index. * Previous chromosome count.

not common in Cactaceae (Las Peñas & al., 2008); with a single pair seen in *Denmoza rhodacantha*. This is expressed by the asymmetry indices: A_1 ranges from 0.15 to 0.28 and A_2 from 0.09 to 0.28. The r value varies from 1.20 to 1.48 (Table 1). More karyotypes, however, are needed in order to establish hypotheses of chromosome evolution across this ornamental, and important family.

Physical mapping by FISH for four species (*Arrojadoa rhodacantha, Melocactus bahiensis, Pachycereus pringlei, Tacinga inamoena*) were here performed for the first time in metaphase chromosomes (Fig. 9). Sites of 18S-5.8S-26S genes located in terminal regions of short chromosome arms are frequent in angiosperms (Acosta & al., 2016; Chiarini & al., 2017). The number of 18S-5.8S-26S rDNA increases with the ploidy level (except *Tacinga inamoena*, Fig. 9). The number of 5S sites was one per haploid genome and were proximally located only on the short arms of the chromosomes. In addition, the 5S rDNA loci have a variable position in the family (Las Peñas & al., 2014, 2017; Moreno & al., 2015).

ACKNOWLEDGEMENTS

The authors thank Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Agencia Nacional de Promoción Científica y Tecnológica (FONCyT), SECyT (Universidad Nacional de Córdoba, Argentina), and FCEFy N for finantial support. Diego Uñates helped with the illustrations and Juan Urdampilleta provided some samples from Brasil and Bolivia.

LITERATURE CITED

- Acosta, M.C., Moscone, E.A. & Cocucci, A.A. 2016. Using chromosomal data in the phylogenetic and molecular dating framework: Karyotype evolution and diversification in *Nierembergia* (Solanaceae) influenced by historical changes in sea level. *Pl. Biol.* 18: 514–526. https://doi.org/10.1111/plb.12430.
- Chiarini, F., Moreno, N., Moré, M. & Barboza, G.E. 2017. Chromosomal changes and recent diversification in the Andean genus *Jaborosa* (Solanaceae). *Bot. J. Linn. Soc.* 183: 57–74. https://doi. org/10.1111/boj.12493.
- Goldblatt, P. & Johnson, D.E. 2013. Index to plant chromosome numbers. Missouri Botanical Garden. Available from http:// www.tropicos.org/Project/IPCN
- Las Peñas, M.L., Bernardello, G. & Kiesling, R. 2008. Karyotypes and fluorescent chromosome banding in *Pyrrhocactus* (Cactaceae). *Pl. Syst. Evol.* 272: 211–222. https://doi.org/10.1007/ s00606-007-0611-5.
- Las Peñas, M.L., Urdampilleta, J.D., Bernardello, G. & Forni-Martins, E.R. 2009. Karyotypes, heterochromatin, and physical mapping of 18S-26S rDNA in Cactaceae. *Cytogenet. Genome Res.* 124: 72–80. https://doi.org/10.1159/000200090.
- Las Peñas, M.L., Urdampilleta, J.D., López-Carro, B., Santiñaque, F., Kiesling, R. & Bernardello, G. 2014. Classical and molecular cytogenetics and DNA content in *Maihuenia* and *Pereskia* (Cactaceae). *Pl. Syst. Evol.* 300: 549–558. https://doi.org/10.1007/ s00606-013-0903-x.
- Las Peñas, M.L., Oakley, L., Moreno, N.C. & Bernardello, G. 2017. Taxonomic and cytogenetic studies in *Opuntia* ser. *Armatae* (Cactaceae). *Botany* 95: 101–120. https://doi.org/10.1139/cjb-2016-0048.
- Levan, A., Sandberg, A. & Fredga, K. 1964. Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201–220.
- Moreno, N.C., Amarilla, L.D., Las Peñas, M.L. & Bernardello, G. 2015. Molecular cytogenetic insights into the evolution of the epiphytic genus *Lepismium* (Cactaceae) and related genera.

Bot. J. Linn. Soc. 177: 263-277. https://doi.org/10.1111/boj.12242.

- Pinkava, D.J. 2002. On the evolution of the continental North American Opuntioideae. Pp. 59–98 in: Hunt, D. & Taylor, N. (eds.), *Studies in the Opuntioideae (Cactaceae)*. Succulent Plant Research 6. Milborne Port: David Hunt Books.
- Romero Zarco, C. 1986. A new method for estimating karyotype asymmetry. *Taxon* 35: 556–530. https://doi.org/10.2307/1221906.
- Schwarzacher, T. & Heslop-Harrison, P. 2000. Practical in situ hybridization. Oxford: Bios Scientific.

Rohit N. Mane,¹ Kumar Vinod C. Gosavi,^{1,2*} Avinash R. Gholave^{1,3} & Shrirang R. Yadav¹

- 1 Department of Botany, Shivaji University, Kolhapur (MS) 416 004, India
- 2 Department of Botany, HPT Arts & RYK Science College, Nashik (MS) 422 002, India
- 3 Life Science Research Laboratory, Walchand College of Arts and Science, Solapur (MS) 413 006, India
- * Author for correspondence: kumarvinodgosavi@gmail.com
 - * First chromosome count for the species.
 - ** New chromosome number (cytotype) for the species.

Authors express thanks to Head, Department of Botany, Shivaji University, Kolhapur for providing laboratory facilities, SERB (File No.: SB/FT/LS-130/2012).

ACANTHACEAE

**Barleria gibsonii Dalzell

2*n* = 44, CHN. India, Maharashtra, Nandurbar District, Toranmal, 1000 m, 21°53′55.56″N, 74°28′00.64″E, 26 Oct 2014, *K.V.C. Gosavi* 126 (SUK) [Fig. 11A].

*Barleria prattensis Santapau

2*n* = 44, CHN. India, Maharashtra, Nandurbar District, Toranmal, Kalapani, 852 m, 21°51′09.44″N, 74°27′57.77″E, 26 Oct 2014, *K.V.C. Gosavi 125* (SUK) [Fig. 11B].

* Pleocaulus sessilis (Nees) Bremek.

2*n* = 30, CHN. India, Maharashtra, Satara, Kas Plateu, 1237 m, 17°43'14.99"N, 73°49'25.37"E, 10 Mar 2016, *R.N. Mane 56* (SUK) [Fig. 11C].

* Ruellia urens Roth

2*n* = 30, CHN. India, Maharashtra, Kolhapur, Shivaji University Campus, 608 m, 16°40′40.89″N, 74°15′19.93″E, 18 Apr 2016, *R.N. Mane* 54 (SUK) [Fig. 11D].

**Strobilanthes callosa Nees

2*n* = 30, CHN. India, Maharashtra, Nashik District, Anjaneri 880 m, 19°55'35.83"N, 73°34'32.88"E, 24 Apr 2016, *K.V.C. Gosavi* 452 (SUK) [Fig. 11E].

ARECACEAE

*Trachycarpus takil Becc.

2*n* = 36, CHN. India, Uttarakhand, Pithoragarh, 2000 m, 29°34′58.29″N, 80°13′05.47″E, 17 Jun 2016, *S.R. Yadav 226* (SUK) [Fig. 11F].

DIPTEROCARPACEAE

* Vatica chinensis L.

2*n* = 22, CHN. India, Karnataka, Mangalore, Pilikula Nisargadhama, Vamanjoor, 78 m, 12°55′38.57″N, 74°53′57.45″E, 30 Aug 2016, *R.N. Mane* 67 (SUK) [Fig. 11G].

FABACEAE

Clitoria biflora Dalzell

2n = 14, CHN. India, Maharashtra, Nandurbar District, Toranmal, Leghapani, 765 m, 21°06′41.85″N, 73°46′19.72″E, 25 Sep 2015, *K.V.C. Gosavi 186* (SUK) [Fig. 11H].



E26

POACEAE

Hemarthria protensa Nees ex Steud.

n = 9, CHN. India, Assam, Guwhati, Deepor Bheel, 52 m, 26°06'52.71"N, 91°39'03.71"E, 19 Apr 2014, *K.V.C. Gosavi 171* (SUK) [Fig. 111].

**Lasiurus scindicus Henrard

n = 18, CHN. India, Rajasthan, Jaisalmer District, Jaisalmer, 190 m, 18 Jan 2015, 26°57′42.24″N, 71°19′17.05″E, *K.N. Koli 167* (SUK) [Fig. 11J].

Manisuris myurus L.

n = 8, CHN. India, Tamil Nadu, Kanchipurum District, Wadajabad, 56 m, 12°48′28.89″N, 79°49′22.92″E, 10 Oct 2015, *K.V.C. Gosavi 196* (SUK) [Fig. 11K].

Mnesithea striata (Nees ex Steud.) de Koning & Sosef

n = 18, CHN. India, Meghalaya, Jowai, on the way, 1237 m, 25°27'23.57"N, 92°11'26.50"E, 12 Dec 2015, *K.V.C. Gosavi 235* (SUK) [Fig. 11L].

SAPOTACEAE

*Chrysophyllum roxburghii G.Don

2*n* = 26, CHN. India, Karnataka, Anmod Ghat, 629 m, 15°56′24.22″N, 73°59′41.66″E, 4 Jun 2017, *R.N. Mane 61* (SUK) [Fig. 11M].

José Ignacio Márquez-Corro,^{1#*} Modesto Luceño,^{1#} Pedro Jiménez-Mejías,² Marcial Escudero,³ Santiago Martín-Bravo,¹ Andrew L. Hipp,^{4,5} Kyong-Sook Chung,⁶ A. Muthama Muasya,⁷ Paul E. Rothrock,⁸ Jaime A. Weber⁴ & Robert Naczi⁹

- 1 Department of Molecular Biology and Biochemical Engineering, Universidad Pablo de Olavide, Carretera de Utrera km 1, 41013 Seville, Spain
- 2 Biology (Botany), Universidad Autónoma de Madrid, Ciudad Universitaria de Cantoblanco, Calle Francisco Tomás y Valiente, 7, 28049 Spain
- 3 Department of Plant Biology and Ecology, University of Seville, Reina Mercedes sn, 41012 Seville, Spain
- 4 The Morton Arboretum, 4100 Illinois Route 53, Lisle, Ilinois 60532, U.S.A.
- 5 The Field Museum, Integrative Research Center, 1400 S Lake Shore Dr., Chicago, Illinois 60605, U.S.A.
- 6 Department of Medicinal Plant Science, Jungwon University, Goesan-gun, Chungbuk 28024, Korea
- 7 Department of Biological Sciences, University of Cape Town, Private Bag X3, Rondebosch, 7701 Cape Town, South Africa
- 8 Department of Biology, Indiana University, Bloomington, Indiana 47405, U.S.A.
- 9 New York Botanical Garden, 2900 Southern Boulevard, Bronx, New York 10458-5126, U.S.A.
- [#] The two first authors have equally contributed to this article.
- * Author for correspondence: jimarcorr@gmail.com
 - * New cytotype for the species.
 - ** First chromosome count for the species.
 - *** First chromosome count for the genus.
 - ! First chromosome count for the subfamily.

Initials of the contributor(s) for each count appear as a superscript at the end of the corresponding species name. Meiotic configurations of chromosomes (I: monovalent, II: bivalent, III: trivalent, IV: tetravalent) are reported after the diploid number.

Here we report chromosome number counts from the sedge family (Cyperaceae), most of which correspond to genus *Carex* from the Nearctic and western Palearctic. In addition, counts are reported for some Neotropical and Afrotropical sedges. We provide the first known chromosome counts for the subfamily Mapanioideae, 5 genera (*Lagenocarpus, Afroscirpoides, Dracoscirpoides, Trianoptiles, Calyptrocarya*) and 45 species.

CYPERACEAE SUBFAMILY CYPEROIDEAE TRIBE ABILDGAARDIEAE

Fimbristylis spadicea (L.) Vahl ML

 $2n = 20 (10^{II})$, CHN. Brazil, Pernambuco, Porto de Galinhas, 15 Aug 1996, *Luceño & al. s.n.* (UFP).

TRIBE BISBOECKELEREAE

*** Calyptrocarya glomerulata (Brongn.) Urb. ML

 $2n = 20 (10^{II})$, CHN. Brazil, Pernambuco, Cabo, Ecological Reserve of Gurjaú, atlantic forest, 1995, *Luceño & al. s.n.* (UFP) [Fig. 12X].

TRIBE CARICEAE

**Carex acidicola Naczi RN

 $2n = 50 (25^{II})$, CHN. U.S.A., Georgia, Clarke County, 1 mi ESE of Athens, 0.5 mi S of end of Rock and Shoals Road, 19 Apr 1988, *Naczi 1783* (MICH, NY).

** Carex aethiopica Schkuhr ML

2n = 74 (37^{II}), CHN. South Africa, Western Cape Province, Cape Town, Table Mountain, 12 Sep 2009, *Luceño & al. s.n.* (BOL).

A chromosome number of 2n = 76 (38^{II}) [Fig. 12A] was also found in a specimen of the same population.

** Carex amphibola Steud. RN

 $2n = 56 (28^{II})$, CHN. U.S.A., Arkansas, Scott County, 2 mi N of Y City along W side of route 71, just S of Fourche LaFave River, 14 May 1991, *Naczi 2829* (MICH, NY).

** Carex asynchrona Naczi RN

 $2n = 52 (26^{II})$, CHN. Mexico, Tamaulipas, 7 km WNW of Gómez Farias, vicinity of Rancho del Cielo Biological Station, 10 Apr 1992, *Naczi 2949* (MICH).

**Carex bolanderi Olney KSC,ME,AH

 $2n = 50 (25^{II})$, CHN. U.S.A., Oregon, Clackamas County, south slope of Mt. Hood, above Timberline Lodge access road, $45^{\circ}18'24.00''$ N, $121^{\circ}54'42.00''$ W, 13 Sep 2009, *Zika 24689* (MOR).

2n = 58 (29^{II}), CHN. U.S.A., Greenhouse collection grown in Illinois, DuPage County, The Morton Arboretum, from material sent by collaborator; greenhouse voucher: *Norris 43* (MOR). Original population information: New Mexico, Catron County, Gila National Forest, Mogollon Mountains, Bursum Road above Mogollon, 33.370918°N, 108.721147°W, 23 Jul 2008, *Norris, Kleinman & Walkup 2008-7-23-43* (SNM).

** Carex borbonica Lam. ML

 $2n = 68 (34^{II})$, CHN. France, La Réunion, Plaine de Remparts, surroundings of Commerson crater, 21°13′15.00″S, 55°38′44.00″E, 2280 m, 1 Jan 2009, *Luceño & Guzmán 1ML09* (UPOS-3635) [Fig. 12B].

** Carex borbonica Lam. × C. boryana Schkuhr ML

2n = 68 (34^{II}), CHN. France, La Réunion, Cirque de Mafate, Le Maïdo, 21°04′29.40″S, 55°23′14.50″E, 2180–2200 m, 6 Jan 2009, *Luceño & Guzmán 20ML09* (UPOS-3724) [Fig. 12C].

Carex boryana and *C. borbonica* constitute two ecologically and morphologically well-defined species (Escudero & Luceño, 2011). However, genetic differentiation has been shown to be scarce (Escudero & Luceño, 2009) and their chromosome number is herein reported to be the same. As this specimen presents morphological traits clearly intermediate between both species, it is worth mentioning that there were no irregularities found in the meiotic pairing (shown in Fig. 12C). Future genomic studies are required on these species, which might shed light on the apparently contrasting pattern of high morphological vs. low genetic/cytogenetic divergence.

** Carex boryana Schkuhr ML

2*n* = 68 (34^{II}), CHN. France, La Réunion, Les Makes, La Fenêtre, 21°11′07.80″S, 55°25′57.80″E, 1500–1650 m, 5 Jan 2009, *Luceño & Guzmán 14ML09* (UPOS-3641) [Fig. 12D].

Carex brevior (Dewey) Mack. ex Lunell KSC,AH

2*n* = 60 (30^{II}), CHN. U.S.A., Iowa, Boone County, 42°01′24.348″N, 93°43′09.012″W, 322 m, 10 Jul 2008, *Norris & Thompson s.n.* (MOR).

** Carex brysonii Naczi RN

 $2n = 50 (25^{II})$, CHN. U.S.A., Alabama, Winston County, 10 mi N of Double Springs, Sipsey River Recreation Area of Bankhead National Forest, 10 Apr 1992, *Naczi 2950* (MICH, NY).

** Carex bulbostylis Mack. RN

 $2n = 54 (27^{II})$, CHN. U.S.A., Texas, Denton County, 4.2 mi W of junction of I-35 and Road 455 in Sanger, S of Road 455, along E side of Clear Creek, 26 Apr 1990, *Naczi 2328* (MICH, NY).

** Carex calcifugens Naczi RN

 $2n = 52 (26^{II})$, CHN. U.S.A., South Carolina, Dorchester County, 6 mi SW of Summerville, along E side of Ashley River, N of route 17A, 22 Apr 1989, *Naczi 2059* (MICH, NY, PH).

Carex cespitosa L. PJM,ML

2n = 78 (39^{II}), CHN. Spain, Navarra, Lesaka, Zalain, 43°16'33.50"N, 01°41'48.60"W, 30 m, 16 Jul 2006, *Jiménez-Mejías & al. 99PJM06* (UPOS-1930).

** Carex conoidea Schkuhr ex Willd. RN

2n = 68 (34^{II}), CHN. Canada, New Brunswick, York County, Kings Landing, above St. John River, 26 Apr 1990, *Naczi 2347* (MICH, NY).

** Carex corrugata Fernald RN

2n = 58 (29^{II}), CHN. U.S.A., Virginia, Sussex County, 2 mi W of Homeville, along E side of Nottoway River, 0.3 mi N of route 40 crossing, 10 Jun 1989, *Naczi 2238* (MICH, NY).

 $2n = 63 (30^{II}+1^{III})$, CHN. U.S.A., California, Butte County, Pat's Meadow, along Butte Creek, 0.6 mi W of Cherry Creek Campground, $40^{\circ}05'55.50''$ N, 121°30'33.54" W, 19 Jul 2008, *Janeway & Gog 3197C* (MOR).

Carex demissa Hornem. PJM

2n = 70, CHN. Portugal, Sintra, Colares, $38^{\circ}46'40.00''$ N, $09^{\circ}26'15.00''$ W, 351 m, 3 Jan 2005, *Jiménez-Mejías & al. 06PJM05* (UPOS); 2 May 2005, *Jiménez-Mejías & al. 185PJM05* (UPOS-2866); Spain, Ávila, Sierra de Gredos, trail to Laguna Grande, 1915 m, 5 Jul 2004, *Jiménez-Mejías & al. 201PJM04* (UPOS-3518); Spain, Cádiz, Tarifa, $36^{\circ}04'34.30''$ N, $05^{\circ}31'50.40''$ W, 364 m, 16 Nov 2003, *Jiménez-Mejías & al. 28PJM03* (UPOS).

Carex demissa Hornem. × C. lepidocarpa Tausch PJM

 $2n = 68 (34^{II})$, CHN. Spain, Asturias, Cangas de Onís, Ercina lake, 2 Sep 2004, *Marín 15504JMM* (UPOS-3577); Spain, Asturias, Somiedo mountain pass, 19 Jul 2005, *Jiménez-Mejías & al. 451PJM05* (UPOS-2615).

2*n* = 70 (35¹¹), CHN. Spain, Lleida, Val d'Arán, Baños de Tredós, 1800 m, 16 Aug 2005, *Luceño & al. 9205MLbis* (UPOS-1405).

** Carex densa (L.H.Bailey) L.H.Bailey KSC, ME, AH

2*n* = 54 (27^{II}), CHN. U.S.A., California, Butte County, Bidwell Park, 39°46'33.72"N, 121°45'47.04"W, 18 Jul 2008, *Janeway & Gog 3183A* (MOR).

Carex distans L. ML,SMB

2*n* = 70 (35^{II}), CHN. Turkey, Karaman, Sarıveliler, 36°41'32.40"N, 32°31'57.00"E, 1615 m, 29 Mar 2010, *Martín-Bravo & al. 112SMB10* (UPOS-7401) [Fig. 12F].

***Carex ecklonii* Nees ^{ML}

2*n* = 64 (32^{II}), CHN. South Africa, Western Cape, Gordons bay, 8 m, 12 Sep 2009, *Luceño & al. s.n.* (UPOS-6503) [Fig. 12E].

** Carex edwardsiana E.L.Bridges & Orzell RN

 $2n = 52 (26^{II})$, CHN. U.S.A., Texas, Bandera County, 4 mi N of Vanderpool, along E side of Sabinal River, just SE of Road 187, 26 Apr 1990, *Naczi 2321* (MICH, NY).

** Carex flaccosperma Dewey RN

 $2n = 46 (23^{II})$, CHN. U.S.A., Arkansas, Cleveland County, 2 mi NW of New Edinburg, at junction of routes 97 and 8, Marks' Mill Battleground historic monument, 9 Apr 1991, *Naczi 2733* (MICH, NY).

** Carex fracta Mack. KSC,AH

 $2n = 70 (35^{II})$, CHN. U.S.A., Oregon, Deschutes County, $44^{\circ}27'17.928''N$, $121^{\circ}46'43.463''W$, 3925 ft, 16 Aug 2008, *Wilson & Otting 15143* (MOR).

Carex glaucodea Tuck. ex Olney RN

 $2n = 44 (22^{II})$, CHN. Canada, Ontario, Essex County, 3.5 mi ENE of Harrow, Colchester South Township, Concession III, lot 17, 10 Apr 1992, *Naczi 2954* (MICH, NY).

Agrees with counts of $2n = 22^{11}$ reported by Wahl (1940) from two plants.



Fig. 12. Cyperaceae chromosomes. All images depict the metaphase of the meiosis I, except for *Carex rainbowii*, *Cyperus fuscus* and *Trianoptiles capensis*, taken from pollen grain mitosis. **A**, *Carex aethiopica*, 2n = 76 (38^{II}); **B**, *C. borbonica*, 2n = 68 (34^{II}); **C**, *C. borbonica* × *boryana*, 2n = 68 (34^{II}); **D**, *C. boryana*, 2n = 68 (34^{II}); **E**, *C. ecklonii*, 2n = 64 (32^{II}); **F**, *C. distans*, 2n = 70 (35^{II}); **G**, *C. oedipostyla*, 2n = 68 (34^{II}); **H**, *C. rainbowii*, n = 29; **I**, *C. reuteriana* subsp. *mauritanica*, 2n = 74 (37^{III}); **J**, *C. reuteriana* subsp. *mauritanica*, 2n = 74 (37^{III}); **J**, *C. reuteriana* subsp. *mauritanica*, 2n = 74 (37^{III}); **K**, *C. trinervis*, 2n = 82 (41^{II}); **C**, *C. vixdentata*, 2n = 62 (31^{II}); **M**, *Lagenocarpus guianensis*, 2n = 34 (17^{II}); **N**, *Eleocharis confervoides* (= *Websteria submersa*), 2n = 40 (2^{III}+17^{II}); **O**, *Cyperus fuscus*, n = 18; **P**, *C. salzmannianus* (= *Lipocarpha salzmanniana*), 2n = 38 (19^{II}); **Q**, *C. subsquarrosus* (= *Hemicarpha micrantha*), 2n = 52 (26^{II}); **R**, *Isolepis marginata*, 2n = 8 (4^{III}); **S**, *Dracoscirpoides ficinioides*, 2n = 104 (52^{III}); **T**, *Afroscirpoides dioeca* (= *Scirpoides dioeca*), 2n = 35 (17^{II+11}, 1^{III+16II}); **U**, *Rhynchospora brownii*, 2n = 36 (18II); **V**, *R. modesti-lucennoi*, 2n = 84 (42^{III}); **W**, *Trianoptiles capensis*, n = 26; **X**, *Calyptrocarya glomerulata*, 2n = 20 (10^{III}); **Y & Z**, *Chrysitrix capensis*, 2n = 46 (1^{IV+21II}, 2^{IV+19II}). — Scale bar = 10 µm.

** Carex glomerabilis V.I.Krecz. ^{JMC,ML}

 $2n = 60 (30^{II})$, CHN. South Africa, Western Cape, Wilderness, Brown Hooded Kingfisher trail, between $33^{\circ}59'01.30''$ S, $22^{\circ}39'03.60''$ E and $33^{\circ}58'36.90''$ S, $22^{\circ}39'12.80''$ E, between 24 and 52 m, 14 Oct 2017, *Márquez-Corro & al. 196JMC17* (UPOS-9968).

** Carex godfreyi Naczi RN

 $2n = 50 (25^{II})$, CHN. U.S.A., Florida, Clay County, 1 mi N of Green Cove Springs, Magnolia Springs, 0.4 mi N of route 17, W of Haven Avenue and S of stream, 13 Apr 1991, *Naczi 2740* (MICH, NY).

** Carex grisea Wahlenb. RN

2n = 56 (28^{II}), CHN. Canada, Québec, Deux Montagnes County, Oka, Parc Paul-Sauvé, floodplain of Lac des Deux Montagnes, near La Crête campground, 7 Apr 1988, *Naczi 1768* (MICH, NY).

The chromosome number report for *C. grisea* by Wahl (1940) is based on a misidentification. His voucher (*Wahl 15106*, BH, NCU, PAC) is *C. amphibola*.

Carex hitchcockiana Dewey RN

2n = 54 (27^{II}), CHN. Canada, Ontario, Hastings County, ca. 10 mi E of Belleville, W of Salmon River, along S side of route 401, 19 Apr 1988, *Naczi 1786* (MICH, NY).

Agrees with $2n = 27^{II}$ reported by Wahl (1940) from one plant.

** Carex impressinervia Bryson, Kral & Manhart RN

 $2n = 46 (23^{II})$, CHN. U.S.A., Mississippi, Marion County, ca. 17 mi SSE of Columbia, 1 mi N of Pearl River County border, 2.3 mi E of route 43, 10 Apr 1992, *Naczi 2956* (MICH, NY).

** Carex leporinella Mack. KSC, ME, AH

2n = 82 (41^{II}), CHN. U.S.A., Idaho, Valley County, Snowbank Mountain, near pond beside the road up mountain, 44°25′08.472″N, 116°07′54.911″W, 20 Aug 2009, *Wilson & al. 15858* (MOR).

** Carex mariposana L.H.Bailey ex Mack. KSC, ME, AH

 $2n = 81 (39^{II}+1^{III})$, CHN. U.S.A., California, Tuolumne County, Yosemite National Park, Gaylor Lake, $37^{\circ}52'50.484''$ N, $119^{\circ}23'53.483''$ W, 11 Aug 2009, *Taylor & al. 20752* (MOR).

***Carex mesochorea* Mack. KSC,ME,AH

2n = 48 (24^{II}), CHN. U.S.A., Michigan, Washtenaw County, 14 Jul 2011, *Reznicek s.n.* (MOR).

*Carex molesta Mack. ex Bright KSC,AH

2n = 72 (36^{II}), CHN. U.S.A., Wisconsin, Dane County, 43°02'21.54"N, 89°25'55.2"W, 896 m, 17 Jun 2008, *Hipp & Gog 3137* (MOR).

** Carex nervina L.H.Bailey KSC,ME,AH

 $2n = 52 (26^{II})$, CHN. U.S.A., California, Butte County, just E of Philbrook Reservoir, 40°01'20.28"N, 121°27'13.439"W, 18 Jul 2009, *Janeway & Gog 3195C* (MOR).

Carex nigra (L.) Reichard PJM, ML, SMB

 $2n = 83 (40^{II}+1^{III})$, CHN. Portugal, Guarda, Covilha, Serra da Estrela, 40°18′51.00″N, 07°37′27.00″W, 1885 m, 1 Aug 2006, *Martín-Bravo 216SMB06* (UPOS); Spain, Ávila, Pinar de Hoyocasero, 1200 m, 3 Jul 2004, *Marín & Luceño 40JMM04* (UPOS); 12 May 2006, *Jiménez-Mejías & al. 31PJM06* (UPOS-2253); Spain, Granada, Sierra Nevada National Park, 2860 m, 19 Aug 2006, *Jiménez-Mejías & Escudero 160PJM06* (UPOS); Spain, Zamora, Puebla de Sanabria, Cubillas lagoon, 42°11′40.00″N, 06°43′41.00″W, 1800 m, 28 Aug 2007, *Martín-Bravo & al. 153SMB07* (UPOS).

 $2n = 84 (42^{II})$, CHN. France, Corse, Ghisoni, $42^{\circ}04'16.80''N$, 09°11'12.10''E, 804 m, 4 May 2007, *Escudero & Luceño 89ME07* (UPOS); Greece, Epiro, Konitsa, Vasilitsa, 1580 m, 12 Jul 2004, *Vargas & Luceño 284PV04* (UPOS-831).

 $2n = 84 (1^{I}+40^{II}+1^{III}, 11 \text{ cells}; 39^{II}+2^{III}, 4 \text{ cells}; 42^{II}, 1 \text{ cell}),$ CHN. Portugal, Guarda, Covilha, Serra da Estrela, 40°18′51.00″N, 07°37′27.00″W, 1885 m, 1 Aug 2006, *Martin-Bravo 216SMB06* (UPOS).

 $2n = 84 (42^{II})$, CHN. Spain, Ávila, pathway to Circo de Gredos, 13 May 2006, *Jiménez-Mejías 36PJM06* (UPOS).

2n = 84 (42^{II}, 2 indiv.), CHN. Spain, Almería, Bayárcal, 37°06'14.00"N, 03°00'28.00"W, 2350 m, 7 Aug 2006, *Jiménez-Mejías* & *García 153PJM06* (UPOS); Spain, Burgos, Sierra de Neila, 18 Jul 2006, *Jiménez-Mejías & al. 136PJM06* (UPOS-2220); Spain, Granada, Sierra Nevada National Park, 3085 m, 19 Aug 2006, *Jiménez-Mejías* & *Escudero 157PJM06* (UPOS).

 $2n = 84 (39^{II}+2^{III}+f)$, CHN. Spain, Soria, between Duruelo and Covaleda, 18 Jul 2006, *Jiménez-Mejías & al. 137PJM06* (UPOS-2221).

 $2n = 85 (41^{II}+1^{III})$, CHN. Morocco, Marrakech, Oukaimedem, 31°11'04.00"N, 07°50'43.00"W, 12 May 2005, *Chaparro & al. 07AJC05* (UPOS-1632); Morocco, Rif, Djbel Tidighine, May 2007, *Jiménez-Mejías & al. 88PJM07* (UPOS-4750); Spain, Soria, Covaleda, 41°54'03.30"N, 02°49'27.60"W, 1134 m, 18 Jul 2006, *Jiménez-Mejías & al. 135PJM06* (UPOS-2218).

2*n* = 85 (41^{II}+1^{III}, 2 indiv.), CHN. Spain, Almería, Bayárcal, 37°06'14.00"N, 03°00'28.00"W, 2350 m, 7 Aug 2006, *Jiménez-Mejías & García 153PJM06* (UPOS).

 $2n = 86 (41^{II}+1^{IV})$, CHN. Morocco, Rif, Djbel Tidighine, May 2007, *Jiménez-Mejías & al. 88PJM07* (UPOS-4750).

** Carex oedipostyla Duval-Jouve ML

2*n* = 68 (34^{II}), CHN. Spain, Cádiz, Alcalá de los Gazules, pathway to Aljibe peak, 36°30'42.00"N, 05°37'30.00"W, 860 m, Jul 2011, *Luceño & al. 1ML11* (UPOS-6228); 36°33'53.66"N, 05°36'06.61"W, 450 m, 23 Feb 2013, *Maguilla & al. 1EMS13* (UPOS-5127) [Fig. 12G].

Carex oligocarpa Schkuhr ex Willd. RN

2n = 54 (27^{II}), CHN. U.S.A., Illinois, Kane County, Elgin, along W side of route 25, S of I-90, Trout Park, 14 May 1991, *Naczi 2902* (MICH, NY).

Agrees with count of $2n = 27^{\text{II}}$ reported by Wahl (1940) from one plant.

*Carex oronensis Fernald KSC,AH

 $2n = *68, *71, 74 (34^{II}, 30^{II}+2^{III}+1^{V}, 37^{II})$, CHN. U.S.A., Maine, Penobscot County, E side of Maine Hwy 2 across from Mt. Hope Cemetery in Veazie, opposite main entrance and just S of Mt. Hope Ave, 44°49′21″N, 68°43′16″W, 27 Jun 2009, *Reznicek 11968* (MOR).

** Carex ouachitana Kral, Manhart & Bryson RN

2n = 52 (26^{II}), CHN. U.S.A., Arkansas, Polk County, summit of Rich Mountain, N of route 88 across from fire tower, 7 Apr 1988, *Naczi 1774* (MICH, NY).

*Carex pachycarpa Mack. KSC,AH

2*n*=70 (35^{II}), CHN. U.S.A., Oregon, Lake County, 42°03'44.712"N, 120°09'35.28"W, 5734 ft, 10 Jul 2008, *Otting & Wilson 1945B* (MOR).

42°03'35.104"N, 120°09'53.083"W, 10 Jul 2008; *Wilson & Otting 14947* (MOR), *Wilson & Otting 14948* (MOR); 42°03'29.611"N, 120°10'4.29"W, 10 Jul 2008, *Wilson & N. Otting 14949* (MOR); 42°03'35.104"N, 120°09'53.082"W, 5808 ft, 24 Jul 2008, *Wilson & Otting 15017* (MOR).

*2*n* = 74 (37^{II}), CHN. U.S.A., Oregon, Lake County, 42°03'35.104"N, 120°09'53.083"W, 24 Jul 2008, *Wilson 15018* (MOR).

**Carex pachystachya* Cham. ex Steud. KSC,AH

2*n* = 82 (41^{II}), CHN. U.S.A., Oregon, Lake County, 43°36′50.586″N, 122°07′3.752″W, 4310 ft, 25 Jul 2008, *Wilson 15052* (MOR).

** Carex paeninsulae Naczi, E.L.Bridges & Orzell RN

 $2n = 48 (24^{II})$, CHN. U.S.A., Florida, Clay County, 1 mi N of Green Cove Springs, Magnolia Springs, 0.4 mi N of route 17, W of Haven Avenue and S of stream, 10 Apr 1992, *Naczi 2958* (MICH, NY).

** Carex petasata Dewey KSC,AH

2*n* = 74 (37^{II}), CHN. U.S.A., Oregon, Lake County, 42°03'45.775"N, 120°09'18.064"W, 5749 ft, 24 Jul 2008, *Wilson 15019* (MOR).

** Carex pigra Naczi RN

2n = 56 (28^{II}), CHN. U.S.A., Alabama, Morgan County, 2 mi N of Falkville along W side of route 31, 9 Apr 1991, *Naczi 2735* (MICH, NY).

** Carex planispicata Naczi RN

 $2n = 50 (25^{II})$, CHN. U.S.A., Alabama, Madison County, Huntsville, Monte Sano State Park, along hiking trail near the Sinks, 22 Apr 1989, *Naczi 2060* (MICH, NY).

Carex projecta Mack. KSC,AH

 $2n = 62 (31^{II})$, CHN. U.S.A., Michigan, Menominee County, *Rothrock* 4070 (MOR).

2*n* = 64 (32^{II}), CHN. U.S.A., Ontario, Simcoe County, *Rothrock* 3971 (MOR).

 $2n = 66 (33^{II})$, CHN. U.S.A., Maine, Washington County, *Rothrock* 3970 (MOR). U.S.A., Wisconsin, Price County, *Rothrock* 4289 (MOR).

One population of 2n = 66 and the counts 2n = 62 and 2n = 64 correspond to the unpublished data cited by Hipp (2007).

Carex radiata (Wahlenb.) Small KSC,AH

2n = 58 (29^{II}), CHN. U.S.A., Wisconsin, Dane County, 43°02'25.14"N, 89°25'50.58"W, 880 ft, 17 Jun 2008, *Hipp & Gog 3133* (MOR).

***Carex rainbowii* Luceño, Jim.Mejías, M.Escudero & Martín-Bravo ^{ME,ML,SMB}

n = 29 (pollen grain mitosis), CHN. South Africa, KwaZulu-Natal, Rainbow gorge, 28°57′35.00″S, 29°13′34.00″E, 1525 m, 13 Nov 2011, *Martín-Bravo & Luceño 120SMB11* (UPOS-5030) [Fig. 12H].

* Carex retroflexa Muhl. ex Willd. KSC, ME, AH

2n = 42 (29^{II}), CHN. U.S.A., Michigan, Washtenaw County, Huron Parkway Nature Area, 14 Jun 2009, *Reznicek & Walters s.n.* (MOR).

Carex reuteriana subsp. *mauritanica* (Boiss. & Reut.) Jim.Mejías & Luceño ^{PJM}

2n = 74 (37^{II}, 2 indiv.), CHN. Spain, Huelva, La Nava, Múrtiga river, 5 Jan 2005, *Jiménez-Mejías & al. 22PJM05* (UPOS) [Fig. 121]. 2n = 74 (37^{II}), CHN. Spain, Córdoba, Villaviciosa de Córdoba, 38°05'03.00"N, 04°57'15.00"W, 24 Mar 2005, *Escudero & al. 4ME05* (UPOS) [Fig. 12J].

** Carex simulata Mack. KSC, ME, AH

2*n* = 64 (32^{II}), CHN. U.S.A., Colorado, Boulder County, Redrock Lake, 40°04'53.40"N, 105°32'23.74"W, 2 Aug 2009, *Chung & Lederer 1039* (MOR).

** Carex subbracteata Mack. KSC, ME, AH

2n = 80 (40^{II}), CHN. U.S.A., Washington, San Juan County, 48°28'48.00"N, 123°03'42.00"W, 29 Aug 2009, *Zika 24641* (MOR).

**Carex subfusca* W.Boott KSC,ME,AH

2*n*=82 (4¹¹), CHN. U.S.A., Oregon, Lake County, 42°29'19.383"N, 120°44'08.89"W, 6078 ft, 4 Aug 2008, *Wilson & al. 15068* (MOR).

* 2*n* = 86 (43^{II}), CHN. U.S.A., Nevada, Washoe County, Tahoe Meadows, 1.5 air km SW of Mt. Rose Highway summit, on Route 341, northern Sierra Nevada Highlands, 39°18′00.00″N, 119°54′00.00″W, 20 Aug 2009, *Zika 24619* (MOR).

** Carex thornei Naczi RN

2n = 50 (25^{II}), CHN. U.S.A., Georgia, Decatur County, Army Corps of Engineers' East Bank Campground on Lake Seminole, 14 Apr 1988, *Naczi 1779* (MICH, NY).

*Carex tincta (Fernald) Fernald KSC,AH

2n = 72 (36^{II}), CHN. U.S.A., Maine, Hancock County, 44°35'35.02"N, 68°30'36.00"W, 4 Jul 1999, *Reznicek 10919A* (MOR).

Carex tribuloides Wahlenb. KSC,AH

 $2n = 70 (35^{II})$, CHN. U.S.A., Indiana, Grant County, *Rothrock* 4243 (MOR). U.S.A., Indiana, St. Joseph County, *Rothrock* 3839C (MOR).

2*n* = 72 (36^{II}), CHN.U.S.A., Indiana, St. Joseph County, *Rothrock* 3839A (MOR).

**Carex trinervis* Degl. ^{ML}

*2n = *82, 84 (41^{II}, 42^{II}), CHN. Portugal, Coimbra, Figueira da Foz, Lagoa da Vela, 4 Jan 2005, *Jiménez-Mejías & al. 10PJM05* (UPOS) [Fig. 12K].

** Carex vernacula L.H.Bailey KSC,ME,AH

 $2n = 52 (26^{II})$, CHN. U.S.A., Colorado, Gunnison County, summit of Cement Creek, $38^{\circ}58'16.536''$ N, $106^{\circ}46'30.144''$ W, 4 Aug 2009, *Chung 1063* (MOR).

Carex viridula Michx. ML

 $2n = 70 (35^{II})$, CHN. Greece, Pellas, Pamagista, Voras mountain, 2200 m, 15 Aug 2004, *Luceño 1404ML* (UPOS-691).

** Carex vixdentata (Kük.) G.A.Wheeler ML

2n = 62 (31^{II}), CHN. Argentina, Buenos Aires, Mar de Ajó, 3 Dec 2002, *Marín 2102JMM* (UPOS-849) [Fig. 12L].

TRIBE CARPHEAE

*** Trianoptiles capensis (Steud.) Harv. ML

 $2n = 52 (26^{II})$, CHN. South Africa, Western Cape Province, Cape Town, Kenilworth Race Course, Sep 2009, *Muasya & al. 4549* (BOL) [Fig. 12W].

TRIBE CRYPTANGIEAE

***Lagenocarpus guianensis Nees ^{ML}
2n = 34 (17^{II}), CHN. Brazil, Pernambuco, Dois Irmãos, Açude do Prata, 1995, *Luceño & al. s.n.* (UFP) [Fig. 12M].

TRIBE CYPEREAE

****Afroscirpoides dioeca* (Kunth) García-Madr. & Muasya (= *Scirpoides dioeca* (Kunth) J.Browning) ^{ML}

 $2n = 35 (17^{II}+1^{I}, 1^{III}+16^{II})$, CHN. South Africa, Western Cape Province, Yzerfontein, Sep 2009, *Muasya & al. 4557* (BOL) [Fig. 12T].

Cyperus fuscus L. ML

2n = 36 (18^{II}), CHN. Spain, Seville, Pablo de Olavide University campus, Jul 2009, *M. Luceño s.n.* (UPOS) [Fig. 12O].

***Cyperus salzmannianus* (Steud.) Bauters (= *Lipocarpha salzmanniana* Steud.) ^{ML}

2*n* = 38 (19^{II}), CHN. Brazil, Rio Grande do Norte, Genipabú, 1995, *M. Luceño & al. s.n.* (UFP) [Fig. 12P].

n = 18, 19, 20 (pollen grain mitosis), CHN. Brazil, Pernambuco, Porto de Galinhas, 1996, *Luceño & Guerra s.n.* (UFP).

** Cyperus sesquiflorus (Torr.) Mattf. & Kük.

(= Kyllinga odorata Vahl) ML

2n = 108 (54^{II}), CHN. Brazil, Pernambuco, Recife, Cidade Universitária, Universidade Federal de Pernambuco campus, 1995, *Luceño & al. s.n.* (UFP).

** Cyperus subsquarrosus (Muhl.) Bauters (= Hemicarpha micrantha (Vahl) Britton) ML

 $2n = 52 (26^{II})$, CHN. Brazil, Pernambuco, Recife, Cidade Universitária, Universidade Federal de Pernambuco campus, 1995, *Luceño & al. s.n.* (UFP) [Fig. 12Q].

2n = 54 (27^{II}), CHN. Brazil, Pernambuco, Recife, Cidade Universitária, Universidade Federal de Pernambuco campus, 1995, *Luceño & al. s.n.* (UFP).

The same chromosome number was found in other specimen of the same population.

***Dracoscirpoides ficinioides (Kunth) Muasya ML

 $2n = 104 (52^{II})$, CHN. South Africa, KwaZulu-Natal, Garden Castle, 29°45′01.10″S, 29°12′43.10″E, 1803 m, 16 Nov 2012, *Maguilla & al. 79EMS12* (UPOS) [Fig. 12S].

** Isolepis marginata (Thunb.) A.Dietr. MM

2n = 8 (4^{II}), CHN. South Africa, Cape Town, Rhodes Memorial, Sep 2009, *Muasya & al. 4516* (BOL) [Fig. 12R].

TRIBE ELEOCHARIDEAE

***Eleocharis confervoides* (Poir.) Steud. (= *Websteria submersa* (C.Wright) Britton) ^{ML}

 $2n = 38, 40, 41 (19^{II}, 1 \text{ indiv.}; 2^{III}+17^{II}, 1 \text{ indiv.}; 2^{III}+17^{II}, 1^{III}+18^{II}+1^{I}, 1^{III}+19^{II}, 1 \text{ indiv.}), n = 20-21 (in pollen grain mitosis), CHN. Brazil, Pernambuco, Dois Irmãos, Açude do Prata, 1995,$ *Luceño & al. s.n.*(UFP) [Fig. 12N].

TRIBE RHYNCHOSPOREAE

Rhynchospora brownii Roem. & Schult. ^{ML} 2*n* = 36 (18^{II}), CHN. South Africa, KwaZulu-Natal, Drakensberg, Cathedral Peak, 29°56'39.20"S, 29°13'29.50"E, 1377 m, 20 Dec 2008, *Luceño & al. 125ML08* (UPOS-3707) [Fig. 12U].

** Rhynchospora modesti-lucennoi Castrov. ML

2n = 84 (42^{II}), CHN. Portugal, Setúbal, Alcácer do Sal, 38°23'43.70"N, 08°40'05.10"W, 11 m, 30 Apr 2005, *Jiménez-Mejías & al. 113PJM05* (UPOS-1394) [Fig. 12V].

TRIBE SCHOENEAE

*Schoenus nigricans L. ML

 $2n = 76 (38^{II})$, CHN. Spain, Huelva, Doñana National Park, Punta del Malandar, 25 Feb 2007, *Jiménez-Mejías & al. 3PJM07* (UPOS-2857).

SUBFAMILY MAPANIOIDEAE TRIBE CHRYSITRICHEAE

 $! Chrysitrix capensis L. ^{\rm ML}$

 $2n = 46 (1^{IV}+21^{II}, 2^{IV}+19^{II}, 1^{III}+21^{II}+1^{I})$, CHN. South Africa, Western Cape Province, Cape Town, Table Mountain, Sep 2009, *Muasya & al. 4540* (BOL) [Fig. 12Y, Z].

LITERATURE CITED

- Escudero, M. & Luceño, M. 2009. Systematics and evolution of Carex sects. Spirostachyae and Elatae (Cyperaceae). Pl. Syst. Evol. 279: 163–189. https://doi.org/10.1007/s00606-009-0156-x
- Escudero, M. & Luceño, M. 2011. Taxonomic revision of the tropical African group of *Carex* subsect. *Elatae* (sect. *Spirostachyae*, Cyperaceae). *Anales Jard. Bot. Madrid* 68: 225–247. https://doi.org/10.3989/ajbm.2256
- Hipp, A.L. 2007. Nonuniform processes of chromosome evolution in sedges (*Carex*: Cyperaceae). *Evolution (Lancaster)* 61: 2175– 2194. https://doi.org/10.1111/j.1558-5646.2007.00183.x
- Wahl, H.A. 1940. Chromosome numbers and meiosis in the genus *Carex. Amer. J. Bot.* 27: 458–470. https://doi. org/10.1002/j.1537-2197.1940.tb14707.x

Antonina V. Natyaganova,¹ Elena V. Mincheva,¹ Yurii S. Bukin,¹ Lyubov S. Kravtsova,¹ Tatiana E. Peretolchina,¹ Tatiana I. Triboy¹ & Denis A. Krivenko^{2,3*}

- 1 Limnological Institute of the Siberian Branch of the Russian Academy of Sciences, Ulan-Batorskaya Str. 3, 664033 Irkutsk, Russia
- 2 Siberian Institute of Plant Physiology & Biochemistry of the Siberian Branch of the Russian Academy of Sciences, Lermontov Str. 132, 664033 Irkutsk, Russia
- 3 Western Baikal Protected Areas, Baikalskaya Str. 291B, 664050 Irkutsk, Russia
- * Author for correspondence: krivenko.irk@gmail.com

The research was supported by the state projects LIN SB RAS #AAAA-A16-116122110060-9.

HYDROCHARITACEAE

Elodea canadensis Michx.

2n = 48, CHN. Russia, Irkutskaya Oblast', Irkutsk city, Akademgorodok, left bank of Angara River, 52°15′03″N, 104°17′00″E, in water, 10 Oct 2017, *A.V. Natyaganova & Yu.S. Bukin 50785* (IRK) [Fig. 13A]; Russia, Irkutskaya Oblast', Irkutskii District, Lake Baikal, 2 km SW of Bol'shie Koty village, opposite of Zhilishche fold, 51°53′32″N, 105°02'46"E, in water, 7 Jun 2018, A.V. Natyaganova 50789 (IRK); Russia, Irkutskaya Oblast', Irkutsk city, Pervomaiskii microdistrict, pond near the public transport station Botanical Garden, 52°15'42"N, 104°14'35"E, in water, 10 Jul 2018, A.V. Natyaganova 50786 (IRK) [Fig. 13B]; Russia, Irkutskaya Oblast', Ol'khonskii District, Lake Baikal, Maloe More strait, Mukhor bay, 53°03'12"N, 106°48'09"E, in water, 12 Jul 2018, Yu.R. Zakharova & M.V. Bashenkhaeva 50791 (IRK); Russia, Irkutskaya Oblast', Irkutsk city, Novo-Leninskii lake-marsh complex, near the public transport station Uzlovaya, 52°19'43"N, 104°13'59"E, in water, 12 Jul 2018, A.V. Natyaganova 50787 (IRK) [Fig. 13C]; Russia, Irkutskaya Oblast', Irkutskii District, Lake Baikal, Listvyanka settlement, 51°51'01"N, 104°52'02"E, in water, 20 Jul 2018, E.V. Mincheva 50788 (IRK) [Fig. 13D]; Russia, Irkutskava Oblast', Ust'-Kut city, right bank of Lena River, in area of bridge across the river, 56°47'27"N, 105°46'59"E, in water, 24 Jul 2018, M.P. Tskhovrebova 50795 (IRK) [Fig. 13E]; Russia, Irkutskaya Oblast', Irkutskii District, Lake Baikal, Bol'shoe Goloustnoe settlement, at the pier of ships, 52°01'37"N, 105°24'24"E, in water, 29 Jul 2018, A.V. Natyaganova 50796 (IRK) [Fig. 13F]; Russia, Irkutskaya Oblast', Irkutsk city, Akademgorodok, left bank of Angara River, backwater, 52°15'00"N, 104°17'05"E, 423 m, in water, 7 Aug 2018, D.A. Krivenko & A.V. Natyaganova 50771 (IRK); Russia, Republic of Buryatiya, Barguzinskii District, Lake Baikal, Chivyrkuiskii bay, Zmeinaya cove in area of hot springs, 53°46'05"N, 109°01'20"E, in water, 2 Jun 2018, A.V. Natyaganova 50790 (IRK); Russia, Republic of Buryatiya, Severo-Baykal'skii District, Lake Baikal, Nizhneangarsk city, at the pier of ships, 55°45'59"N, 109°33'09"E, in water, 3 Jun 2018, A.V. Natvaganova 50792 (IRK); Russia, Republic of Buryatiya, Kabanskii District, Lake Baikal, Posolskii sor bay, 51°57'15"N, 106°08'12"E, in water, 25 Jul 2018, L.S. Kravtsova 50793 (IRK) [Fig. 13G]; Russia, Zabaikal'skii Krai, Chitinskii District, southern part of Vitim Plateau, Ivano-Arakhleyskaya system of lakes, Shakshinskoe Lake, 52°09'21"N, 112°43'46"E, in water, 17 Jun 2014, E.V. Mincheva 50794 (IRK); Russia, Moskovskaya Oblast', Pushchino town, Fianovskii pond, 59°49'12"N, 37°37'12"E, in water, 20 Oct 2018, Yu.S. Bukin 51398 (IRK).

Egeria densa Planch.

2n = 48, CHN. Russia, Moskovskaya Oblast', Lyubertsy urban district, Kraskovo suburban village, Pekhorka River opposite Karl Marx Str. 1, 115 m, 55°39'39"N, 37°58'15"E, in water, 22 Sep 2018, *A.V. Natyaganova 51053* (IRK).

Saroj Kumar Pradhan,^{1*} Raghbir Chand Gupta¹ & Rajesh Goel²

- 1 Department of Botany, Punjabi University, Patiala, Punjab, 147002, India
- 2 Department of Pharmaceutical Sciences and Drug Research, Punjabi University, Patiala, Punjab, 147002, India
- * Author for correspondence: pradhan_sarojkumar@yahoo.com
 - * First chromosome count for the species.
 - ** New chromosome number (cytotype) for the species.
 - ▼ New chromosome count from an Indian accession.

The study is supported by financial grant under DBT-IPLS Project (Reference number: BT/PR 4548/INF/22/146/2012).

ASTERACEAE

▼ Arctium lappa L.

n = 17, CHN. India, Uttarakhand, Uttarkashi, way to Har ki Doon, 31°07′28.71″N, 78°22′06.10″E, 2800 m, 21 Aug 2015, *S.K. Pradhan 32112* (PUN 60928) [Fig. 14A].

The present chromosome number is in agreement with the previous report of 2n = 34 from outside of India (Ge & al., 1989). From India, this chromosome number is reported here for the first time. For this species also other cytotypes were reported from India, namely n = 16(Bala & Gupta, 2011), n = 18 (Mehra, 1976) and 2n = 36 from outside of India (Májovský & al., 1970; Pagni & Corsi, 1979; Huang & al., 1988).

** Carduus edelbergii Rech.f.

n = 16, CHN. India, Uttarakhand, Uttarkashi, Bhaironghati, 31°01'40.83"N, 78°52'06.09"E, 2790 m, 15 Jun 2016, *S.K. Pradhan 32149* (PUN 60929) [Fig. 14B].

The present chromosome count is the first report of tetraploid, based on 2n = 16 reported from outside of India (Podlech & Dieterle, 1969). From India, n = 20 has also been reported (Malik & al., 2011).

* Cremanthodium arnicoides (DC. ex Royle) R.D.Good

n = 9, CHN. India, Uttarakhand, Uttarkashi, Bhojwasa, 30°57'06.78"N, 79°02'58.34"E, 3808 m, 17 Jun 2016, *S.K. Pradhan 32150* (PUN 60930) [Fig. 14C].



Fig. 13. Mitotic metaphases. **A-G**, *Elodea canadensis*, 2n = 48. — Scale bars = 10 µm.



Fig. 14. A, *Arctium lappa*, PMC at diakinesis, n = 17 (PUN 60928); **B**, *Carduus edelbergii*, PMC at diakinesis, n = 16 (PUN 60929); **C**, *Cremanthodium arnicoides*, PMC at anaphase I, n = 9 (PUN 60930); **D**, *Saussurea auriculata*, PMC at diakinesis, n = 16 (PUN 60715); **E**, *Saussurea costus*, PMC at metaphase I, n = 17 (PUN 58869). — Scale bars = 10 µm

Saussurea auriculata (DC.) Sch.Bip.

n = 16, CHN. India, Uttarakhand, Uttarkashi, Dayara Bugyal, 3100 m, 30°51'35.38"N, 78°35'00.65"E, 30 Aug 2015, *S.K. Pradhan 32138* (PUN 60715) [Fig. 14D].

Previously, 2n = 32 (Singh & al., 2016) was reported from India.

** Saussurea costus (Falc.) Lipsch.

n = 17; India, Uttarakhand, Uttarkashi, Gangar, 31°06′24.58″N, 78°18′55.66″E, 2400 m, 20 Aug 2015, *S.K. Pradhan 32104* (PUN 58869) [Fig. 14E].

Previously the chromosome number of 2n = 36 (Siddique & Wafai, 1993) was reported from India.

LITERATURE CITED

- Bala, S. & Gupta, R.C. 2011. [Reports]. In: Marhold, K. (ed.), IAPT/ IOPB chromosome data 12. *Taxon* 60: 1784–1786.
- Ge, C.J., Li, Y.K., Wan, P. & Hsu, P.S. 1989. Chromosome numbers of 31 medicinal plants from Shandong Province. Pp. 267–272 in: Hong, D. (ed.), *Plant chromosome research 1987: Proceedings of the Sino-Japanese Symposium on Plant Chromosomes Held at Beijing, China, October 6–11, 1987.* Beijing: Organizing Committee of the Symposium.
- Huang, S.-F., Wang, Y.-Q., Chen, Z.-Y. & Shi, X.-H. 1988. Plant chromosome counts (4). Subtrop. Forest Sci. Technol. 16: 25–30.
- Májovský J. (ed.) 1970. Index of chromosome numbers of Slovakian flora (Part 1). Acta Fac. Rerum Nat. Univ. Comen., Bot. 16: 1–26.
- Malik, R.A., Gupta, R.C. & Kumari, S. 2011. [Reports]. In: Marhold, K. (ed.), IAPT/ IOPB chromosome data 12. *Taxon* 60: 1790, E47–E49.
- Mehra, P.N. 1976. *Cytology of Himalayan Hardwoods*. Calcutta: Sree Saraswaty Press.
- Pagni A.M. & Corsi, G. 1979. Cariologia di alcune specie d'interesse officinale della flora italiana. *Atti Soc. Tosc. Sci. Nat. Pisa. Mem.*, ser. B, 86: 465–473.
- Podlech, D. & Dieterle, A. 1969. Chromosomenstudien an afghanischen Pflanzen. Candollea 24: 185–243.
- Siddique, M.A.A. & Wafai, B.A. 1993. New bases number for Saussurea lappa C.B. Clarke an important endangered medicinal plant of Kashmir Himalayas. Chromosome Infor. Serv. 54: 5–6.
- Singh, V., Gupta, R.C. & Kaur, K. 2016. [Reports]. In: Marhold, K. (ed.), IAPT/ IOPB chromosome data 21. *Taxon* 65: 676, E17–E19.

Angeline M.S. Santos,¹ Felipe Nollet,² Alessandro Rapini,³ Roxana C. Barreto,⁴ Joel M.P. Cordeiro,¹ Erton M. Almeida,¹ José L. Silva,¹ Fabiane R.C. Batista⁵ & Leonardo P. Felix¹

- 1 Laboratório de Citogenética Vegetal, Departamento de Ciências Biológicas, Centro de Ciências Agrárias, Universidade Federal da Paraíba, Campus II, 58397-000 Areia, Paraíba Brazil
- 2 Laboratório de Sistemática e Florística de Angiospermas, Departamento de Biologia, Universidade Federal Rural de Pernambuco, 52171-900 Recife, Pernambuco, Brazil
- 3 Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas, 44036-900 Feira de Santana, Bahia, Brazil
- 4 Universidade Federal de Pernambuco, Departamento de Botânica, 50670-901 Recife, Pernambuco, Brazil
- 5 Instituto Nacional do Semiárido, 58434700, Campina Grande, Paraíba, Brazil
- * Author for correspondence: angelinemssantos@gmail.com

Financial support from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), and INSA (Instituto Nacional do Semiárido).

Methods for chromosome analysis are according to Guerra & Souza (2002).

- * First chromosome count for the species.
- ** New cytotype for the species.

APOCYNACEAE

** Allamanda doniana Müll.Arg.

2*n* = 27, CHN. Brazil, Ceará, Guaraciaba do Norte, 04°19'55"S, 40°51'08"W, 20 Apr 2014, *E.M. Almeida 1137* (EAN) [Fig. 15A].

Aspidosperma pyrifolium Mart.

2*n* = 34, CHN. Brazil, Paraíba, Areia, 06°58'12"S, 35°42'47"W, 3 Feb 2015, *A.M.S. Santos 24* (EAN) [Fig. 15B].

Mandevilla bahiensis (Woodson) M.F.Sales & Kin.-Gouv. 2*n* = 20, CHN. Brazil, Bahia, Piatã, 13°09′24″S, 41°45′53″W, 29 Apr 2015, *L.P. Felix 15520* (EAN) [Fig. 15C]. **Mandevilla catimbauensis* Souza-Silva, Rapini & J.F. Morales 2*n* = 26, CHN. Brazil, Pernambuco, Buíque, 08°35′37″S, 37°12′20″W, 30 Jul 2014, *J.M.P. Cordeiro* 379 (EAN) [Fig. 15D].

Mandevilla dardanoi M.F.Sales, Kin.-Gouv. & A.O. Simões 2n = 20, CHN. Brazil, Alagoas, União dos Palmares, 09°02'06"S, 35°52'32"W, 30 Oct 2014, J.M.P. Cordeiro 481 (EAN) [Fig. 15E].

**Mandevilla leptophylla* (A.DC.) K.Schum. 2*n* = 18, CHN. Brazil, Ceará, Meruoca, 03°32′03″S, 40°26′11″W, 26 Aug 2014, *L.P. Felix 15060* (EAN) [Fig. 15F].

Mandevilla tenuifolia (J.C.Mikan) Woodson 2n = 20, CHN. Brazil, Ceará, São Benedito, 04°07'17"S, 40°51'02"W, 20 Apr 2014, *E.M. Almeida 1165* (EAN) [Fig. 15G].

Marsdenia caatingae Morillo

2*n* = 22, CHN. Brazil, Paraíba, Esperança, 07°04'19"S, 35°58'13"W, 16 Jun 2015, *A.S. Santos 26* (EAN) [Fig. 16A].

**Marsdenia hilariana* E.Fourn.

2*n* = 22, CHN. Brazil, Ceará, Meruoca, 03°32′15″S, 40°35′37″W, 26 Aug 2014, *L.P. Felix 15063* (EAN) [Fig. 16B].

*Marsdenia megalantha Goyder & Morillo

2n = 22, CHN. Brazil, Ceará, Quixadá, 03°48′29″S, 40°57′29″W, 21 Apr 2014, *E.M. Almeida 1169* (EAN) [Fig. 16C].

Peltastes peltatus (Vell.) Woodson

2*n* = 18, CHN. Brazil, Pernambuco, Taquaritinga do Norte, 07°53′28″S, 36°02′06″W, 31 Mar 2012, *L.P. Felix 13771* (EAN) [Fig. 16D].

*Petalostelma dardanoi Fontella

2*n* = 20, CHN. Brazil, Paraíba, Cabaceiras, 07°22'39″S, 36°17'35″W, 25 Jun 2014, *J.M.P. Cordeiro 258* (EAN) [Fig. 16E].

Tabernaemontana catharinensis A.DC.

2*n* = 22, CHN. Brazil, Ceará, Poranga, 04°45′17.5″S, 40°52′02″W, 18 Apr 2014, *L.P. Felix 14943* (EAN) [Fig. 16F].

Tabernaemontana solanifolia A.DC.

2*n* = 22, CHN. Brazil, Bahia, Jacobina, 11°11′35″S, 40°29′51″W, 15 Jan 2015, *J.M.P. Cordeiro 654* (EAN) [Fig. 16G].

*Temnadenia stellaris (Lindl.) Miers

2*n* = 18, CHN. Brazil, Paraíba, Mamanguape, 06°44′22″S, 35°10′30″W, 27 Nov 2014, *J.M.P. Cordeiro 591* (EAN) [Fig. 16H].



Fig. 15. Chromosome complements of species belonging to the family Apocynaceae occurring on inselbergs: **A**, Allamanda doniana, 2n = 27; **B**, Aspidosperma pyrifolium, 2n = 34; **C**, Mandevilla bahiensis, 2n = 20; **D**, M. catimbauensis, 2n = 26; **E**, M. dardanoi, 2n = 20; **F**, M. leptophylla, 2n = 18; **G**, M. tenuifolia, 2n = 20. — Scale bar in G = 10 µm.



Fig. 16. Chromosome complements of species belonging to the family Apocynaceae occurring on inselbergs: A, Marsdenia caatingae, 2n = 22; B, M. hilariana, 2n = 22; C, M. megalantha, 2n = 22; D, Peltastes peltatus, 2n= 18; E, Petalostelma dardanoi, 2n = 20; F, Tabernaemontana catharinensis, 2n = 22; G, T. solanifolia, 2n = 22; H, Temnadenia stellaris, 2n = 18. — Scale bar in F = 10 µm.

ARACEAE

**Anthurium affine Schott
2n = 30, CHN. Brazil, Paraíba, Areia, 07°00'31"S, 35°44'53"W,
25 Mar 2014, *E.M. Almeida 1069* (EAN) [Fig. 17A].

COMMELINACEAE

***Aneilema brasiliense* C.B.Clarke 2*n* = 24, CHN. Brazil, Paraíba, Matureia, 07°14'41"S, 37°25'10"W, 14 May 2016, *L.P. Felix 15908* (EAN) [Fig. 17B].

Commelina diffusa Burm.f.

2n = 30, CHN. Brazil, Paraíba, Esperança, $07^{\circ}04'19''S$, $35^{\circ}58'13''W$, 16 Jun 2015, *A.S. Santos 32* (EAN) [Fig. 17C].

2n = 60, CHN. Brazil, Paraiba, Areia, 07°00'31"S, 35°44'53"W, 25 Mar 2014, *E.M. Almeida 1067* (EAN) [Fig. 17D].

Commelina obliqua Vahl

2*n* = 60, CHN. Brazil, Pernambuco, Buíque, 08°35'37"S, 37°12'18"W, 1 Jul 2015, *L.P. Felix 15665* (EAN) [Fig. 17E].

Dichorisandra hexandra (Aubl.) Standl.

2n = 76, CHN. Brazil, Pernambuco, Bonito, 08°30′40″S, 35°43′26″W, 3 Jul 2015, *L.P. Felix 15736* (EAN) [Fig. 17F].

Tradescantia ambigua Mart.

2*n* = 24, CHN. Brazil, Paraíba, Campina Grande, 07°11′02″S, 35°59′01″W, 16 Jun 2015, *A.S. Santos 18* (EAN) [Fig. 17G].

COSTACEAE

**Chamaecostus subsessilis* (Nees & Mart.) C.D.Specht & D.W.Stev.

2*n* = 18, CHN. Brazil, Paraíba, Areia, 07°00′31″S, 35°44′53″W, 25 Mar 2014, *E.M. Almeida 1074* (EAN). [Fig. 18A]

EUPHORBIACEAE

**Cnidoscolus urens* (L.) Arthur 2*n* = 72, CHN. Brazil, Paraíba, Areia, 07°00'31"S, 35°44'53"W, 25 Mar 2014, *E.M. Almeida 1076* (EAN) [Fig. 18B].

MELASTOMATACEAE

* *Tibouchina heteromalla* (D.Don) Cogn.
2n = 36, CHN. Brazil, Paraiba, Areia, 07°00'31"S, 35°44'53"W,
25 Mar 2014, *E.M. Almeida 1077* (EAN) [Fig. 18C].

SOLANACEAE

**Solanum polytrichum* Moric. 2*n* = 24, CHN. Brazil, Bahia, Una, 15°10'37"S, 39°03'29"W, 10 Feb 2015, *E.M. Almeida 1409* (EAN) [Fig. 18D].



Fig. 17. Chromosome complements of species occurring on inselbergs: Araceae – A, Anthurium affine, 2n = 30; Commelinaceae – B, Aneilema brasiliense, 2n = 24; C, Commelina diffusa, 2n = 30; D, C. diffusa, 2n = 60; E, C. obliqua, 2n = 60; F, Dichorisandra hexandra, 2n = 76; G, Tradescantia ambigua, 2n = 24. — Scale bars in A and F = 10 µm.

VELLOZIACEAE

* Vellozia plicata Mart.

2*n* = 18, CHN. Brazil, Paraíba, Patos, 07°04′15.8″S, 37°13′48.2″W, 1 Nov 2013, *L.P. Felix 14525* (EAN) [Fig. 18E].

VITACEAE

Cissus decidua Lombardi

2*n* = 34, CHN. Brazil, Paraíba, Cabaceiras, 07°22'39"S, 36°17'35"W, 24 Jun 2014, *J.M.P. Cordeiro 260* (EAN) [Fig. 18F].

Inselbergs, also known as rock outcrops, are granitic or gneiss geological formations from pre-Cambrian that stand out in a flat landscape (Sarthou & al., 2010). These formations are characterized by the presence of severe edaphoclimatic conditions in relation to the surrounding environment, which are important selective pressures for plant evolution and speciation. Generally, those rock outcrops are isolated geographically in relation to other habitats, forming islands of plant species that can present morphological and chromosomal differentiation, especially polyploidy, generally related to the species diversification (Dodsworth & al., 2016). We analyzed here 27 species belonging to nine different angiosperm families, occurring on inselbergs from the Northeast region of Brazil. Emphasis in the present work was given to the families Apocynaceae and Commelinaceae, due to the number of species belonging to both families.

Root tips obtained directly from the cultivated material were pre-treated with 0.002 M 8-hydroxyquinoline (8HQ) for 24 h (except Commelinaceae that went by 4 h) in a refrigerator and subsequently fixed in Carnoy 3:1 (absolute ethanol:glacial acetic acid, v/v) for 24 h at room temperature and stored at -20°C. For fluorochrome staining, root tips were digested in an enzymatic solution containing 2% cellulase and 20% pectinase at 37°C for 1 h. Subsequently, they were squashed in a drop of 60% acetic acid. Coverslips were removed in liquid nitrogen. The slides were stained with DAPI (2 μ g/ml) for 30 min. Slides were mounted in glycerol/McIlvaine buffer medium. The best metaphases were captured on a Zeiss light microscope using an Axio Cam MRC5 video camera and Axiovision v.4.8 software. Images were processed in Photoshop CS3.

The species analyzed varied from 2n = 18 in *Mandevilla leptophylla* (Apocynaceae, Fig. 15F), *Peltastes peltatus* (Apocynaceae, Fig. 16D) and *Temnadenia stellaris* (Apocynaceae, Fig. 16H) to 2n =76 in *Dichorisandra hexandra* (Commelinaceae, Fig. 17F). The following previous counts were confirmed: In Apocynaceae by Brito & al. (2014), 2n = 34 for *Aspidosperma pyrifolium* (Fig. 15B), 2n =20 for *Mandevilla bahiensis* (Fig. 15C), *M. dardanoi* (Fig. 15E) and *M. tenuifolia* (Fig. 15G), 2n = 22 for *Marsdenia caatingae* (Fig. 16A), *Tabernaemontana catharinensis* (Fig. 16F) and *T. solanifolia* (Fig. 16G). In Commelinaceae, 2n = 30 and 60 (Pitrez & al., 2001) for *Commelina diffusa* (Figs. 17C, D), 2n = 60 (Pitrez & al., 2001) for *Dichorisandra hexandra* (Fig. 17F), 2n = 76 (Pitrez & al., 2010) for *Tradescantia ambigua* (Fig. 17G). In Vitaceae, 2n = 34 (Cordeiro & al., 2017) for *Cissus decidua* (Fig. 18F).

New counts are presented here for *Mandevilla catimbauensis* with 2n = 26 (Fig. 15D), *M. leptophylla* with 2n = 18 (Fig. 15F), *Marsdenia hilariana* with 2n = 22 (Fig. 16B), *M. megalantha* with 2n = 22 (Fig. 16C), *Petalostelma dardanoi* with 2n = 20 (Fig. 16E), *Temnadenia. stellaris* with 2n = 18 (Fig. 16H) (Apocynaceae); *Chamaecostus subsessilis* with 2n = 18 (Fig. 18A) (Costaceae); *Tibouchina heteromalla* with 2n = 36 (Fig. 18C) (Melastomataceae); *Vellozia plicata* with 2n = 18 (Fig. 18B). New cytotypes are presented for Apocynaceae: *Allamanda doniana* with 2n = 27 (Fig. 15A) and *Peltastes*



Fig. 18. Chromosome complements of species occurring on inselbergs: Costaceae – A, Chamaecostus subsessilis, 2n = 18; Euphorbiaceae – B, Cnidoscolus urens, 2n = 72; Melastomataceae – C, Tibouchina heteromalla, 2n = 36; Solanaceae – D, Solanum polytrichum, 2n = 24; Velloziaceae – E, Vellozia plicata, 2n =18; Vitaceae – F, Cissus decidua, 2n = 34. — Scale bar in F = 10 µm.

peltatus with 2n = 18 (Fig. 16D); Araceae: *Anthurium affine* with 2n = 30 (Fig. 17A); Commelinaceae: *Aneilema brasiliense* with 2n = 24 (Fig. 17B).

Apocynaceae is a globally distributed family consisting of about 400 genera and 5000 species (Endress & al., 2007). In Brazil, about 750 species occur in different kinds of habitat, including inselbergs (Rapini, 2012; Pitrez & al., 2014a). The family is characterized by a remarkable diversity in chromosome numbers, ranging from 2n = 10 in *Rhabdadenia madida* (Vell.) Miers to 2n = 230 in *Vinca difformis* Pourr. The chromosome number variation in Apocynaceae indicates several polyploid and disploid series, suggesting that these events are playing an important role for the diversification of the family.

Commelinaceae present 41 genera and 650 paleo- and neotropical species (Faden, 1983). A wide range of chromosome number variation characterizes the family, as well the scarcity of chromosome numbers revisions. To date, only a review of chromosome evolution has been performed for the family, indicating several cycles of polyploidy and disploid series, with basic chromosome numbers ranging from x = 4 to x = 20 (Keith Jones & Jopling, 1972).

The genus *Commelina* presents 170 species, being considered the largest of the family (Hassemer, 2017). The genus presents a nearly cosmopolitan distribution, occurring in a great diversity of habitats, with terrestrial, rupicolous and aquatic species. *Commelina* occurs in all regions of Brazil, although only nine species are recognized, of which three are endemic (Aona & Pellegrini, 2015). Based only on the chromosome number variation, the genus is characterized by a dysploid series with n = 10, 11, 13, 14, 15, and a subset of chromosome numbers possibly derived by polyploidy (n = 21, 24, 26, 28, 29, 30, 42, 45, 60, 75). Here, *Commelina diffusa* presented 2n = 30 and 2n = 60 for different inselbergs, while *C. obliqua* presented 2n = 60 for all analyzed populations. In the literature there are records for other species of *Commelina*, such as 2n = 60 for *C. erecta* L. and 2n = 30 for *C. rufipes* Seub., analyzed by Pitrez & al. (2001) in populations occurring in different kinds of habitat.

LITERATURE CITED

- Aona, L.Y.S. & Pellegrini, M.O.O. 2015. Commelinaceae in Lista de Espécies da Flora do Brasil. Rio de Janeiro: Jardim Botânico do Rio de Janeiro.
- Brito, J.G., Santos, R.S., Meira, P.R.H., Alves, L.I.F., Andrade, M.J.G., Rapini, A. & Felix L.P. 2014. [Reports] In: Marhold, K., ed., IAPT/IOPB Chromosome Data 17. *Taxon* 63: 1149–1150, E6–E9. https://doi.org/10.12705/635.34
- Cordeiro, J.M.P., Nascimento, S., Santos, A.M.S., Neto, E.M., Santos, A., Assis, F.N.M. & Felix, L.P. 2017. [Reports] In: Marhold, K. & Kučera, J. (eds.), IAPT/IOPB Chromosome Data 24, *Taxon* 66: 275, E1–E5. https://doi.org/10.12705/661.37
- Dodsworth, S., Chase, M.W. & Leitch, A.R. 2016. Is post-polyploidization diploidization the key to the evolutionary success of angiosperms? *Bot. J. Linn. Soc.* 180: 1–5. https://doi.org/10.1111/ boj.12357
- Endress, M.E., Liede-Schumann, S. & Meve, U. 2007. Advances in Apocynaceae: The enlightenment, an introduction. Ann. Missouri Bot. Gard. 94: 259–267. https://doi.org/10.3417/0026-6493(2007)94[259:AIATEA]2.0.CO;2
- Faden, R.B. 1983. Phytogeography of African Commelinaceae. Bothalia 14: 553–557. https://doi.org/10.4102/abc.v14i3/4.1207
- **Guerra, M. & Souza, M.J.** 2002. Como observar cromossomos: Um guia de técnicas em citogenética vegetal, animal e humana. Ribeirão Preto: FUNPEC.

- Hassemer, G. 2017. Taxonomic and nomenclatural notes on neotropical *Commelina* (Commelinaceae), and an identification key for Brazil, Guyana, Paraguay, Suriname and Uruguay. *Phytotaxa* 303: 101–117. http://sci-hub.tw/10.11646/phytotaxa.303.2.1
- Keith Jones, F.L.S. & Jopling, C. 1972. Chromosomes and the classification of the Commelinaceae. *Bot. J. Linn. Soc.* 65: 129–162. https://doi.org/10.1111/j.1095-8339.1972.tb00929.x
- Pitrez, S.R., Felix, L.P., Barreto, R. & Guerra, M. 2001. Números cromossômicos de espécies de Commelinaceae R. Br. ocorrentes no nordeste do Brasil. *Bol. Bot. Univ. São Paulo* 19: 7–14. http:// sci-hub.tw/10.11606/issn.2316-9052.v19i0p7-14
- Pitrez S.R., Andrade, L.A., Assis, F.N.M. & Felix, L.P. 2014a. Is there a relationship between polyploidy and stressful environments? A case study of inselbergs in northeastern Brazil. *Genet. Molec. Res.* 13: 8353–8366. https://doi.org/10.4238/2014. October.20.11
- Pitrez, S.R., Ramalho, F.C., Andrade, L.A., Cordeiro, J.M.P., Assis, F.N.M. & Felix, L.P. 2014b. IAPT/IOPB Chromosome Data 18. Taxon 63: 1387–1393. https://doi.org/10.12705/636.37
- Rapini, A. 2012. Taxonomy "under construction": Advances in the systematics of Apocynaceae, with emphasis on the Brazilian Asclepiadoideae. *Rodriguésia* 63: 75–88. https://doi.org/10.1590/ S2175-78602012000100007
- Sarthou, C., Larpin, D., Fonty, E., Pavoine, S. & Ponge, JF. 2010. Stability of plant communities along a tropical inselberg ecotone in French Guiana (South America). *Flora* 205: 682–694. https:// doi.org/10.1016/j.flora.2010.04.005

Julia V. Shner, Tatiana V. Alexeeva & Michael G. Pimenov*

Botanical Garden, Moscow State University, 119234 Moscow, Russia

* Author for correspondence: mgpimenov@mail.ru

This study was supported by the grants 13-04-00648 and 16-04-00525 from the Russian Foundation for Basic Research (RFBR).

* First chromosome count for the species.

** New chromosome count for the species.

UMBELLIFERAE/APIACEAE

Athamanta sicula L.

2n = 22, CHN. Italy, Sicily, prov. Trapani, Erice, $38^{\circ}02'10''$ N, $12^{\circ}34'58''$ E, 720 m, rocks, 6 Jul 2012, *M.G. Pimenov 12-4* (MW); Italy, Sicily, prov. Messina, Forza d'Agro, $37^{\circ}54'50.47''$ N, $15^{\circ}20'08.11''$ E, 350 m, 22 Jul 2012, *M.G. Pimenov s.n.* (MW); Italy, Sicily, prov. Palermo, near Palermo, $38^{\circ}04'21.23''$ N, $13^{\circ}31'15.02''$ E, 220 m, 13 Jul 2012, *M.G. Pimenov s.n.* (MW) [Fig. 19A].

These determinations of chromosome numbers correspond to four previous ones for the plants from Italy (2n = 22) and Albania (n = 11). Five other studied species of the genus *Athamanta* L. have the same chromosome number (Pimenov & al., 2002).

Berula erecta (Huds.) Coville

n = 9, CHN. Greece, Ionic Islands Reg., Corfu (Κερκυρας), Benitses (Μπενιτσες), 39°31'17.24"N, 19°54'46.98"E, 29 Jun 2015, *M.G. Pimenov 15-30* (MW) [Fig. 20A].

The same number of chromosomes was reported in the majority of publications with some exceptions -2n = 20 was found for the plants from Sweden (Lövkvist, 1963) and Kazakhstan and Tajikistan (Retina & Pimenov, 1977; Shner & al., 2016). There was also a rather strange earlier determination of n = 6 for plants from Illinois (U.S.A.) (Bell & Constance, 1957) which was not confirmed later.

Bupleurum glumaceum Sm.

n = 8, CHN. Greece, Central Greece, Nomos Boeotia (Βοιωτια), Arachova (Αραχωβα), 38°29'01.90"N, 22°35'20.03"E, 800 m, 24 Jun 2015, *M.G. Pimenov 15-7* (MW) [Fig. 20H].

Chromosome numbers of this species were determined earlier at least six times; all determinations were from Greece (see Pimenov & al., 2002) and all showed n = 8, or 2n = 16, which indicates the karyological stability of the species.

Chaerophyllum temulum L.

n = 7, CHN. Greece, North Greece, Epirus (Ηπείρος), Pindos Mts., Nomos Ioannina (Ιωαννίνων), Metsovo (Μετσβο), 39°46'25.09"N, 21°10'49.17"E, 1170 m, 3 Jul 2015, *M.G. Pimenov 15-39* (MW) [Fig. 20B].

This chromosome number was found for the most of the studied populations from different European countries. Other counts (n = 11, 2n = 22, 24) were determined considerably rarer (from Germany, the Netherlands, Bulgaria, Spain, Czech Republic, and Turkey) (see Pimenov & al., 2002).

Conium divaricatum Boiss. & Orph.

2*n* = 22, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), Delphi (Δάλφι), 38°28′54.35″N, 22°39′05.41″E, 560 m, 23 Jun 2015, *M.G. Pimenov 15-4* (MW) [Fig. 19B].

The same chromosome number was reported by Constantinidis & al. (1997) from Greece too, B-chromosomes not revealed in our study, were found in some karyotypes.

Daucus carota subsp. gummifer (Syme) Hook.f. (= D. carota subsp. hispanicus (Gouan) Thell.)

n = 9, CHN. France, Provence, Marseille, Vieux Port, Bas Fort Saint-Nicolas, 43°17'30"N, 05°21'50"E, 10 m, 3 Jun 2017, *M.G. Pimenov s.n.* (MW) [Fig. 20E].

There are at least seven reports for *Daucus carota* subsp. *gummifer* and subsp. *hispanicus* (from Portugal, Spain and France) that all, with one exception, are 2n = 18 or n = 9 (see Pimenov & al., 2002).

Daucus involucratus Sm.

2n = 22, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), municipality Amfissa (Αμφισσα), Sernikaki (Σερνικακι), Mt. Partas, 38°29'17.66"N, 22°23'44.72"E, 113 m, 25 Jun 2015, *M.G. Pimenov 15-17* (MW) [Fig. 19D].

Two chromosome numbers were previously determined for this species, namely 2n = 22 or n = 11 from Greece and Cyprus (Engstrand, 1970; Vogt & Aparicio, 1999) and 2n = 20 from Turkey (Pimenov & al., 1998).

Daucus maximus Desf. (≡ *D.carota* subsp. *maximus* (Desf.) Ball) 2*n* = 18, CHN. Turkey, Canakkale, ruines of Troya, 39°57'32.26"N, 26°14'45.23"E, 7 Aug 2014, *M.G. Pimenov s.n.* (MW) [Fig. 19E].

The chromosome number 2n = 18, typical not only for the species in its strict sense, but also for the whole species-aggregate *D. carota*, studied from different parts of its primary and secondary area as opposed to the majority *Daucus* species that have 2n = 20 or 22.

Eryngium campestre var. virens (Link) Weins

n = 14, CHN. Greece, Central Greece, Nomos Boeotia (Βοιωτια), Arachova (Αραχωβα), 38°29'01.90"N, 22°35'20.039"E, 800 m, 24 Jun 2015, *M.G. Pimenov 15-9* (MW) [Fig. 20D].

Two cytotypes, diploid (2n = 14) and tetraploid (2n = 28), are known for *E. campestre* (see Pimenov & al., 2002). The only other chromosome number determination under the name of var. *virens* from Turkey (Pimenov & al., 1996) showed 2n = 28. There are no other determinations for Greek material of the species. Some botanists, however, doubt the taxonomic status of the variety (Breton, 1962; Wörz, 2011), due to broad overlapping in characters of the variety and *E. campestre* sensu stricto.

Eryngium creticum Lam.

n = 7, CHN. Greece, NW Greece, Ionic Islands Reg., Corfu (Κερκυρας), Benitses (Μπενιτσες), 39°31'17.24"N, 19°54'46.98"E, 29 Jun 2015, *M.G. Pimenov 15-29* (MW) [Fig. 20C].

Our record corresponds to earlier determinations for plants from Bulgaria (Česchmedjiev, 1976), Iran and Greece (Perdigo i Ariso, 1981) and Israel (Baltisberger & Widmer, 2004).

Ferula tingitana L.

2*n* = 22, CHN. Turkey, Muğla, Marmaris, among city houses, 36°50′53.35″N, 28°16′01.84″E, 30 m, 27 Jul 2014, *M.G. Pimenov 14-4* (MW) [Fig. 19H].

This chromosome number was reported for the species from Spain (Aparicio & Garcia Martin, 1986), Morocco (El Alaoui Faris,



Fig. 19. Mitotic chromosomes. A, Athamanta sicula, 2n = 22; B, Conium divaricatum, 2n = 22; C, Ferulago humilis, 2n = 22; D, Daucus involucratus, 2n = 22; E, Daucus maximus, 2n = 18; F, Ferulago trojana, 2n = 22; G, Ferulago mughlae, 2n = 66; H, Ferula tingitana, 2n = 22; I, Tordylium apulum, 2n = 20; J, Helosciadium nodiflorum, 2n = 22; K, Orlaya grandiflora, 2n = 20; L, Ridolfia segetum, 2n = 22. — Scale bars = 5 µm.

2006) and Turkey (Pimenov & al., 1998; Shner & al., 2010). It is not surprising that the species has the same number 2n = 22, as all *Ferula* L. species (more than 100), having been studied up to now.

Ferulago humilis Boiss.

2*n* = 22, CHN. Turkey, Muğla, road from Marmaris to Datca, 40 km from Datca, 36°45′42.96″N, 27°52′12.37″E, 300 m, 28 Jul 2014, *M.G. Pimenov 14-8* (MW) [Fig. 19C].

Two previous records for the species (Strid, 1987; Pimenov & al., 1998) from different localities in Turkey showed also 2n = 22, the most common chromosome number for the genus *Ferulago* W.D.J.Koch.

*Ferulago mughlae Peşmen

2*n* = 66, CHN. Turkey, Muğla, road from Marmaris to Datca, 40 km from Datca, 36°45′42.96″N, 27°52′12.37″E, 300 m, 28 Jul 2014, *M.G. Pimenov 14-9* (MW) [Fig. 19G].

This is the first chromosome number determination for the species, and the first record of hexaploidy in the genus *Ferulago*.

*Ferulago trojana Akalın & Pimenov

2*n* = 22, CHN. Turkey, Ballkesir, Kazdaği Milli Park, Kazdaği Mts., S slope, 39°38′28.17″N, 26°55′09.98″E, 240 m, 6 Aug 2014, *M.G. Pimenov & E. Akalın s.n.* (MW) [Fig. 19F].

This is the first chromosome number report for the recently described rare species, narrow endemic to the Kazdaği mountain massif (Mt. Ida) in the Western Turkey.

Helosciadium nodiflorum (L.) W.D.J.Koch

2n = 22, CHN. Italy, Sicily, prov. Agrigento, between Porto Empedoclo and S. Leone, sea shore, $37^{\circ}16'24''$ N, $13^{\circ}34'14''$ E, 15 Jul 2012, *M.G. Pimenov 12-13* (MW) [Fig. 19J].



Fig.20. Meiotic chromosomes. **A**, Berula erecta, n = 9; **B**, Chaerophyllum temulum, n = 7; **C**, Eryngium creticum, n = 7; **D**, Eryngium campestre var. virens, n = 14; **E**, Daucus carota subsp. gummifer, n = 9; **F**, Seseli varium, n = 9; **G**, Pimpinella cretica, n = 9; **H**, Bupleurum glumaceum, n = 8; **I**, Pastinaca yildizii, n = 11. — Scale bars = 10μ m; n – nucleolus.

There are at least 18 reports on chromosome number determination for the species, widely distributed from Macaronesia to Tanzania and Tajikistan. All of them characterize *H. nodiflorum* (most frequently studied under the name of *Apium nodiflorum* (L.) Lag.) as diploid with n = 11 (2n = 22).

Orlaya grandiflora (L.) Hoffm.

2*n* = 20, CHN. Greece, Ionic Islands Reg., Lefkada (Λευκαδα), near Ligia (Λυγιά) 38°47'10.06"N, 20°43'15.69"E, 20 m, 29 Jun 2015, *M.G. Pimenov 15-25* (MW) [Fig. 19K].

Our chromosome number determination, the first one for the plants of Greek origin, corresponds to the most previous ones (n = 10 or 2n = 20). Besides that, chromosome numbers 2n = 22 (Bulgaria), n = 9 (Romania), 2n = 16 (Morocco), and 2n = 14 (Croatia) are known for *O. grandiflora* (see Pimenov & al., 2002).

*Pastinaca yildizii Dirmerci

n = 11, CHN. Turkey, Ballkesir, Kazdaği Milli Park, Kazdaği Mts., upper forest borderline, 39°41′59.57″N, 26°51′26.66″E, 1700 m, 6 Aug 2014, *M.G. Pimenov & T. Dirmerci 14-10* (MW) [Fig. 201].

This is the first chromosome number determination for the species. It corresponds to all previous records for seven accepted species of *Pastinaca* L., excluding *P. pimpinellifolia* M.Bieb. (2n = 44 from Turkey, Pimenov & al., 1996) (see Pimenov & al., 2002).

Pimpinella cretica Poir.

n = 9, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), municipality Amfissa (Αμφισσα), Sernikaki (Σερνικακι), 38°29'22.71"N, 22°24'31.03"E, 23 Jun 2015, *M.G. Pimenov 15-3* (MW) [Fig. 20G].

2*n* = 18, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), municipality Amfissa (Αμφισσα), Sernikaki (Σερνικακι), Partas Mts., 38°29'17.66"N, 22°23'44.72"E, 113 m, 25 Jun 2015, *M.G. Pimenov s.n.* (MW) [Fig. 21A].

This is the third chromosome number determination for the species; the previous ones were slightly variable: n = 10; 2n = 20 (Jordan – Al-Eisawi, 1989; Crete, Greece – Pimenov & Shner, 2008) and 2n = 18 (Crete, Greece – Pimenov & Shner, 2008). This can be regarded as an evidence of dysploidy in karyotype evolution of the species, which is not rare in some other species of the genus (Yurtseva, 1988; Pimenov & al., 2002). 2n = 18 is the most widespread chromosome number in *Pimpinella* L.

Ridolfia segetum (Guss.) Moris

2n = 22, CHN. Italy, Sicily, prov. Agrigento, between Ribera and Seccagrande, fallow land, $37^{\circ}26'41''N$, $13^{\circ}14'27''E$, 50 m, 15 Jul 2012, *M.G. Pimenov 12-8* (MW) [Fig. 19L].

At least nine different records from Portugal, Spain, Morocco, and Algeria allow to characterize *R. segetum* as a rather stable diploid with n = 11 (2n = 22) (Pimenov & al., 2002). Only one investigation (El Alaoui Faris, 1989) showed intraspecific karyological variability of the species in Morocco (2n = 22 and 2n = 22+2B).

**Scaligeria moreana Engstrand

2n = 22, CHN. Greece, Peloponnisos, Nomos Messinias (Μεσσηνία), Taigetos, NW-part of Mani, above Langada, ravine, slope of SE-facing, maquis shrubland, rocks, under rocks, cracks of rocks, 36°47.05'N, 22°20.30'E, 600 m, 5 Jul 2012, *E.A. Zakharova & U.A. Ukrainskaya 48* (MW) [Fig. 21F].

This is the second chromosome number determination for *S. moreana*. It does not correspond to the previous record of 2n = 20 from the same area (Mt. Taigetos) (Kyriakopoulos & al., 2014).

Scaligeria napiformis (Willd. ex Spreng.) Grande

2n = 20, CHN. Greece, Ionic Islands Reg., Corfu (Κερkυρας), Benitses (Μπενιτσες), 39°31'17.24" N, 19°54'46.98"E, 29 Jun 2015, *M.G. Pimenov 15-32* (MW); Greece, Central Greece, Nomos Phokida (Φωκις), municipality Amfissa (Αμφισσα), Sernikaki (Σερνικακι), Mt. Partas, 38°29'17.66"N, 22°23'44.72"E, 113 m, 25 Jun 2015, *M.G. Pimenov 15-20* (MW); Greece, Peloponnisos, Nomos Lakonias/Messinias (Μεσσηνία), Taigetos, Langada gorge, slope of W-NW-facing, right bank of dry stream, between 42–43 km from Kalamata, near Climbing Centre, steep, limestone rocks, 37°04.88'N, 22°18.68'E, 810 m, 6 Jul 2012, E.A. Zakharova & U.A. Ukrainskaya 47 (MW) [Fig. 21E].

Our chromosome number determination, the fourth for the species and the second from Greece, confirms 2n = 20 as the most frequent number in the species (Engstrand, 1970; Vogt & Aparicio, 1999; Shner & al., 2013) and in the whole genus. One determination from Turkey 2n = 22 (Pimenov & al., 1998), however, showed variability in chromosome numbers in this species.

**Seseli varium Trevir. (= S. pallasii Besser)

n = 9, CHN. Greece, Central Greece, Nomos Boeotia (Βοιωτια), Arachova (Αραχωβα), 38°28′56.89″N, 22°34′56.39″E, ca. 1000 m, 24 Jun 2015, *M.G. Pimenov 15-23* (MW) [Fig. 20F].

This is a new chromosome number for the species, as previous determinations gave 2n = 16 (from Hungary), 2n = 20 (from the Czech Republic, Slovakia and Italy) and 2n = 22 (Ukraine) (see Pimenov & al., 2002). We also determined n = 11 for the plants from Italy. Karyological data were partly published under the name *S. pallasii*, a later synonym of *S. varium*. As to n = 9 (2n = 18), this chromosome number was determined many times (see Pimenov & al., 2002) for a closely related species, *S. elatum* L., distributed in southern and central Europe, and known, in particular (as *S. elatum* subsp. *gouanii* (W.D.J.Koch) P.W.Ball), from Italy to the Balkans.

Smyrnium olusatrum L.

2*n* = 22, CHN. Greece, Ionic Islands Reg., Lefkada (Λευκαδα), near Ligia (Λυγιά), 38°47'10.06"N, 20°43'15.69"E, 28 Jun 2015, 20 m, *M.G. Pimenov s.n.* (MW) [Fig. 21B]; Italy, Sicily, prov. Palermo, W of Cefalù, 08°01'39.31"N, 14°01'35.57"E, 120 m, 13 Jul 2012, *M.G. Pimenov s.n.* (MW); Turkey, Muğla, Marmaris, among city houses, 36°50'53.35"N, 28°16'01.84"E, 30 m, 27 Jul 2014, *M.G. Pimenov s.n.* (MW).

Karyologically invariant species (n = 11; 2n = 22), according to numerous previous chromosome number determinations (see Pimenov & al., 2002). Our three records, among them two, from Italy and Turkey, being the first for each of these countries, confirm chromosome number stability of the species.

Smyrnium rotundifolium Mill.

2*n* = 22, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), Delphi (Δάλφι), 38°28′54.35″N, 22°39′05.41″E, 560 m, 23 Jun 2015, *M.G. Pimenov s.n* (MW) [Fig. 21D]; Italy, Sicilia, prov. Catania, Mt. Etna (Mongbello), S slope, above Nicolosi, 37°41′34″N, 14°59′20″E, 1634 m, 17 Jul 2012, *M.G. Pimenov 12-15* (MW).

Previous records, all from Greece (see Pimenov & al., 2002; Pimenov & Shner, 2008) showed the same number 2n = 22 for the species. Generally, this chromosome number is the only one in *Smyrnium* L.

Thapsia garganica L.

2*n* = 22, CHN. Italy, Sicily, prov. Trapani, Segesta, dry slope near the temple, 37°56′24″N, 12°49′57″E, 350 m, 6 Jul 2012, *M.G. Pimenov 12-1* (MW) [Fig. 21C].

Thapsia garganica belongs to the group of species within the genus, in which chromosome number variability has not been found up to now, in contrast to *T. villosa* L. in which polyploid series (2n = 22, 44, 66) was revealed (see Pimenov & al., 2002).

Tordylium apulum L.

2n = 20, CHN. Greece, Central Greece, Nomos Phokida (Φωκις), municipality Amfissa (Αμφισσα), Sernikaki (Σερνικακι), 38°29'22.71"N, 22°24'31.03"E, 23 Jun 2015, *M.G. Pimenov 15-2* (MW) [Fig. 19I].

Our chromosome number determination corresponds to numerous previous data from Greece, Albania and Italy (see Pimenov & al., 2002; Pimenov & Shner, 2008).

Torilis triradiata Boiss. & Heldr.

2*n* = 12, CHN. Turkey, Muğla, between Içmeler and Turinç, 36°46'18.59"N, 28°11'50.38"E, 370 m, 27 Jul 2014, *M.G. Pimenov 14-3* (MW) [Fig. 21G].

This is the second chromosome number determination for the species, and it does not correspond to the previous report (2n = 16: Shner & al., 2013), made on the Turkish material too (from Isparta Province). In that publication 2n = 16 is an obvious misprint, omitted by the authors in the proofs, as only 12 chromosomes are clearly seen in the microphotograph (Shner & al., 2013: fig. 15). 2n = 12 is one of the usual counts in *Torilis* Adans., and, for instance, in the closely related species *T. arvensis* (Huds.) Link and *T. leptophylla* (L.) Rchb.f.



Fig. 21. Mitotic chromosomes. **A**, *Pimpinella cretica*, 2n = 18; **B**, *Smyrnium olusatrum*, 2n = 22; **C**, *Thapsia garganica*, 2n = 22; **D**, *Smyrnium rotundifolium*, 2n = 22; **E**, *Scaligeria napiformis*, 2n = 20; **F**, *Scaligeria moreana*, 2n = 22; **G**, *Torilis triradiata*, 2n = 12. — Scale bars = 5 µm.

LITERATURE CITED

- Al-Eisawi, D.M.H. 1989. Chromosome counts of Umbelliferae of Jordan. *Ann. Bot. (Rome)* 47: 201–214.
- Aparicio, A. & Garcia Martin, F. 1986. Nota sobre el genero *Ferula* (Umbelliferae) en Andalucia. *Acta Bot. Malac.* 11: 221–226.
- Baltisberger, M. & Widmer, A. 2004. Cytological data of some plant species from Israel. *Israel J. Pl. Sci.* 52: 171–176.
- Bell, C.R. & Constance, L. 1957. Chromosome numbers in Umbelliferae. *Amer. J. Bot.* 44: 565–572. https://doi.org/10.2307/2438928
- Breton, A. 1962. Révision des Eryngiums d'Afrique du Nord. Mém. Soc. Sci. Nat. Phys. Maroc, Bot. 2: 5–86.
- Česchmedjiev, J.V. 1976. [Reports]. In: Löve, A. (ed.), IOPB Chromosome Number Reports LIV. *Taxon* 25: 642–643.
- **Constantinidis, T., Kamari, G. & Phytos, D.** 1997. A cytological study of 28 phanerogams from the mountains of SE Sterea Ellas, Greece. *Willdenowia* 27: 121–142.
- El Alaoui Faris, F.E. 1989. Caryologie d'espèces de la famille des Umbelliferae. Ann. Sci. Nat., Bot. Biol. Veg., ser. 13, 10: 157–169.
- El Alaoui Faris, F.E. 2006. Nombre chromosomique de quelques espèces de férules marocaines (*Ferula*, Apiaceae). *Fl. Medit.* 16: 341–354.
- Engstrand, L. 1970. Studies in the Aegean flora. 18. Notes and chromosome numbers in Aegean Umbelliferae. *Bot. Not.* 123: 384–393.
- Kyriakopoulos, C., Bareka, P. & Kamari, G. 2014. Reports 1840– 1844. In: Kamari, G., Blanché, C. & Siljak-Yakovlev, S. (eds.), Mediterranean chromosome number reports - 24. *Fl. Medit.* 24: 287–291.
- Lövkvist, B. 1963. Berula erecta. P. 700 in: Weimarck, H. (ed.), Skanes flora. Lund: Bokförlaget Corona AB.
- Perdigo i Ariso, M.-T. 1981. Contribution à l'étude cytotaxinomique du genre *Eryngium* L. *Biol. Écol. Médit.* 8: 3–12.
- Pimenov, M.G. & Shner, Ju.V. 2008. Reports 1650–1663. In: Kamari, G., Blanché, C. & Siljak-Yakovlev, S. (eds.), Mediterranean chromosome number reports - 18. *Fl. Medit.* 18: 573–582.
- Pimenov, M.G., Daushkevich, Ju.V., Vasil'eva, M.G. & Kljuykov, E.V. 1996. Reports 716–748. In: Kamari, G., Felber, F. & Garbari, F. (eds.), Mediterranean chromosome number reports - 6. *Fl. Medit.* 6: 288–307.
- Pimenov, M.G., Alexeeva, T.V. & Kljuykov, E.V. 1998. Reports 905–935. In: Kamari, G., Felber, F. & Garbari, F. (eds.), Mediterranean chromosome number reports - 8. *Fl. Medit.* 8: 221–245.
- Pimenov, M.G., Vasil'eva, M.G., Leonov, M.V. & Daushkevich, J.V. 2002. *Karyotaxonomical analysis in the Umbelliferae*. Enfield, Plymouth: Science Publishers.
- Retina, T.A. & Pimenov, M.G. 1977. Chisla khromosom vidov semeistva Umbelliferae Srednei Azii. Nauchnye Dokl. Vysshei Shkoly Biol. Nauki 7(163): 90–95.
- Shner, Ju.V., Pimenov, M.G., Kljuykov, E.V. & Alexeeva, T.V. 2010. Umbelliferae/Apiaceae. In: Marhold, K. (ed.), IAPT/IOPB chromosome data 10. *Taxon* 59: 1937, E10–E12.
- Shner, Ju.V., Alexeeva, T.V., Pimenov, M.G. & Kljuykov, E.V. 2013. Reports 1768–1783. In: Kamari, G., Blanché, C. & Siljak-Yakovlev, S. (eds.), Mediterranean chromosome number reports - 23. *Fl. Medit.* 23: 256–263.
- Shner, Ju.V., Alexeeva, T.V., Kurbonov, A. & Pimenov, M.G. 2016. Apiaceae. In: Marhold, K. & Kučera, J. (eds.), IAPT/IOPB chromosome data 22. *Taxon* 65: 1206, E19–E23.
- Strid, A. 1987. Chromosome numbers of Turkish mountain plants: An annotated Iist of 34 taxa. *Notes Roy. Bot. Gard. Edinburgh* 44: 351–356.

- Vogt, R. & Aparicio, A. 1999. Chromosome numbers of plants collected during Iter Mediterraneum IV in Cyprus. *Bocconea* 11: 117–169. http://www.herbmedit.org/bocconea/11-117.pdf
- Wörz, A. 2011. Revision of Eryngium L. (Apiaceae Saniculoideae): General part and Palaearctic species. Bibliotheca Botanica 159. Stuttgart: Schweizerbart.
- Yurtseva, O.V. 1988. Tsitologicheskoe izuchenie nekotorykh vidov roda *Pimpinella* (Umbelliferae-Apioideae). *Nauchnye Dokl. Vysshei Shkoly Biol. Nauki* 11(301): 78–85.

Mayco Werllen dos Santos Sousa,^{1*} Marisa Toniolo Pozzobon,² José Francisco Montenegro Valls^{2,3} & Regina Célia de Oliveira¹

- 1 Programa de Pós-Graduação em Botânica da Universidade de Brasília, Campus Darcy Ribeiro, Brasília, DF, Brazil, CEP 70910-900
- 2 Embrapa Recursos Genéticos e Biotecnologia, Parque Estação Biológica - PqEB, Brasília, DF, Brazil, CEP 70770-917
- 3 Research Productivity Fellowship/CNPq, Brazil (Proc. 312215/2013-4)
- * Author for correspondence: maycowerllen@gmail.com

Financial support from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior/CAPES, Conselho Nacional de Desenvolvimento Científico e Tecnológico/CNPq (Proc. 312215/2013-4, 562340/2010-6 and 561768/2010-2), and Fundação de Amparo à Pesquisa do Distrito Federal/FAP-DF (Proc. 0193.000.979/2015), Brasília, Brazil.

- * First chromosome count for the species.
- ** New cytotype for the species.

POACEAE

Paspalum centrale Chase

2n = 20, CHN. Brazil, Acre, Rio Branco, 8 km from Rio Branco to Senador Guiomard, 10°01'45.9"S, 67°49'42.6"W, 10 Dec 2017, *R.C. Oliveira 3405* (UB).

Paspalum convexum Humb. & Bonpl. ex Flüggé

2n = 32, CHN. Brazil, Distrito Federal, Brasília, Fercal, 15°31'01.4"S, 47°50'14.6"W, 14 Mar 2018, *J.F.M. Valls & R.C. Oliveira* 16788 (CEN, UB), *J.F.M. Valls & M.W.S. Sousa* 16789 (CEN, UB); Brazil, Goiás, Flores de Goiás, 200 m after BR-020 Hwy., left, 14°54'46.7"S, 46°57'24.7"W, 18 Mar 2015, *R.C. Oliveira & M.W.S. Sousa* 3005 (UB).

*Paspalum cordaense Swallen.

2*n* = 40, CHN. Brazil, Bahia, Luís Eduardo Magalhães, BA-460 Hwy., ca. 13 km SE of Placas, 11°53′55.3″S, 46°10′15.3″W, 29 Mar 2017, *J.F.M. Valls & al. 16694* (CEN, UB).

* Paspalum macranthecium Parodi

2*n* = 20, CHN. Brazil, Bahia, Luís Eduardo Magalhães, BA-460 Hwy., ca. 13 km SE of Placas, 11°53′55.3″S, 46°10′15.3″W, 29 Mar 2017, *J.F.M. Valls & al. 16693* (CEN, UB).

* Paspalum marmoratum Kuhlm.

2n = 18, CHN. Brazil, Mato Grosso, Chapada dos Guimarães, Chapada dos Guimarães National Park, Morro São Jerônimo,

15°26'05.7"S, 55°51'25.4"W, 12 Feb 2016, *C.O. Moura & Y.F.F. Soares.* 98 (UB) [Fig. 22B].

Paspalum melanospermum Desv. ex Poir.

2n = 20, CHN. Brazil, Amazonas, Careiro de Várzea, BR-319 Hwy., 11 km after gas station, 03°20'11.0"S, 59°52'41.8"W, 27 Apr 2015, *R.C. Oliveira & J.M. Mendoza 3025* (UB) [Fig. 22C]; Brazil, Bahia, Feira de Santana, Universidade Estadual de Feira de Santana, 12°12'01.2"S, 38°58'17.9"W, 12 Oct 2017, *R.C. Oliveira 3362* (UB); Brazil, Goiás, Monte Alegre, 300 m from córrego das Lages, along GO-118 Hwy., on road to Monte Alegre, 13°19'34.0"S, 47°00'48.0"W, 28 Mar 2017, *Valls & al. 16671* (CEN, UB).

** 2n = 40, CHN. Brazil, Amazonas, Presidente Figueiredo, Balbina dam, 01°57'40.9"S, 59°29'23.5"W, 25 Apr 2015, R.C. Oliveira & J.M. Mendoza 3010 (UB); Brazil, Distrito Federal, Brasília, Farm Água Limpa, 15°56′55.5″S, 47°56′02.6″W, 7 Apr 2018, R.C. Oliveira & al. 3443 (UB); Brazil, Goiás, Flores de Goiás, 200 m after Formosa-Flores intersection, left, 14°54'46.7"S, 46°57'24.7"W, 18 Mar 2015, R.C. Oliveira & M.W.S. Sousa 3003 (UB); Brazil, Goiás, Alto Paraíso de Goiás, farm entrance, ca. 4.5 km S Alto Paraiso, 14°10'41.9"S, 47°30'40.5"W, 27 Mar 2017, J.F.M. Valls. & al. 16622 (CEN, UB); Brazil, Goiás, Flores de Goiás, GO-531 Hwy., ca. 1 km from BR-020, 14°54'37.8"S, 46°57'19.5"W, 16 Mar 2016, J.F.M. Valls & al. 16188 (CEN, UB); Brazil, Goiás, Flores de Goiás, 25 km south of Flores, GO-531 Hwy., 14°40'14.0"S, 46°59'44.0"W, 16 Mar 2016, J.F.M. Valls. & al. 16195 (CEN, UB); Brazil, Mato Grosso, Chapada dos Guimarães, Chapada dos Guimarães National Park, Morro São Jerônimo, 15°26'05.7"S, 55°51'25.4"W, 12 Feb 2016, C.O. Moura & Y.F.F. Soares 90 (UB); Brazil, Tocantins, Wanderlândia, BR-226 Hwy., towards Imperatriz, 06°50'57.5"S, 47°57'47.5"W, 10 Feb 2017, M.W.S. Sousa & E.C. Antunes 87 (UB).

Paspalum plicatulum Michx.

2*n* = 40, CHN. Brazil, Distrito Federal, Brasília, Bernardo Sayão Ecological Park, 15°50'10.5"S, 47°48'40.5"W, 18 Jan 2017, *A.P.M.L. Gouveia 232* (UB); Brazil, Distrito Federal, Brasília, Brasilia National Park, 15°58'43.2"S, 47°57'18.7"W, 10 Feb 2017, *R.C. Oliveira & M.W.S. Sousa 3423* (UB).

* Paspalum riparium Nees

2*n* = 40, CHN. Brazil, Amazonas, Manaus, INPA, 03°05'40.4"S, 59°59'21.3"W, 25 Apr 2015, *R.C. Oliveira & J.M. Mendoza 3008* (UB) [Fig. 22D]; Brazil, Presidente Figueiredo, Balbina dam, 01°54'37.4"S, 59°27'36"W, 25 Apr 2015, *R.C. Oliveira & J.M. Mendoza 3016* (UB).

Paspalum rojasii Hack.

2n = 40, CHN. Brazil, Distrito Federal, Brasília, Bernardo Sayão Ecological Park, 15°50'15.4"S, 47°48'42.5"W, 17 Jan 2018, *J.F.M. Valls.* & *M.W.S. Sousa 16777* (CEN, UB).

METHODS

Chromosome numbers were determined from root-tip cells (mitosis) or pollen-mother cells (meiosis), following Sousa & al. (2017). Young inflorescences were collected in the field and root tips were acquired from cultivated material.

The genus *Paspalum* L. includes nearly 350 species (Sánchez-Ken, 2010) mostly in America, some in Africa, Asia and Oceania, and three or four species are considered cosmopolitan (Rua & al., 2010).

Paspalum belongs to a clade where a basic chromosome number of x = 10 is plesiomorphic (Rua & al., 2010; Scataglini & al., 2013),

with few exceptions (Bonasora & al., 2015). Multiple intraspecific chromosome numbers frequently occur in *Paspalum* as an outcome of different ploidy levels (Quarín, 1992). The reproductive system of *Paspalum* is complex. Many species of this genus consist of sexual-diploid and apomictic-polyploid cytotypes, and several of them arose through hybridization (Quarín & Norrmann, 1990). Apparently, sexual-diploids are distributed in rather restricted areas, as well as certain hybrid combinations (Peñaloza & al., 2008).

We confirmed reported counts for *P. plicatulum* and *P. rojasii*, both with 2n = 40 chromosomes (Dandin & Chennaveeraiah, 1983, 1988; Honfi & al., 1990; Sinha & al., 1990; Norrmann & al., 1994; Pozzobon, 2000), and for *P. centrale*, with 2n = 20 (Davidse & Pohl, 1972).

The literature reports 2n = 20, 32, 40, and 60 chromosomes for *P. convexum* (Gould, 1966; Reeder, 1967, 1984; Pohl & Davidse, 1971; Davidse & Pohl, 1974; Selva, 1976). Our findings show that the chromosome number 2n = 32 [Fig. 22A] is typical of *P. convexum*, a species with broadly to very broadly obovate spikelets, distinct from all other annual species in the informal group Plicatula. Although no vouchers of diverging counts assigned to this name have been examined so far, we suggest that plants of allied species with obovate spikelets and x = 10 may have been taken for *P. convexum* by different authors.

The present work provides the first chromosome counts for the annual species *Paspalum cordaense*, 2n = 40, *P. macranthecium*, 2n = 20, and *P. riparium*, 2n = 40 [Fig. 22D]. *Paspalum riparium* is only known from river edges in the Amazon basin, while the other two are predominantly from the cerrado region.

The study showed an unusual chromosome count for *P. marmo*ratum, with 2n = 2x = 18 [Fig. 22B]. This number is only known, in the same genus, for *P. commutatum* Nees (Burton, 1942) and *P. reduncum* Nees ex Steud. (Peñaloza & al., 2008).

Two ploidy levels (2n = 2x = 20 [Fig. 22C] and 2n = 4x = 40) were documented for *P. melanospermum*, both levels on a broad geographic area, including several Brazilian states. Although the diploid level was previously known the tetraploid had not been reported in the past.



Fig. 22. Meiotic cells of *Paspalum* species. **A**, *P. convexum*, diakinesis, *J.F.M. Valls & al. 16187 (2n* = 32); **B**, *P. marmoratum*, diakinesis, *C.O. Moura & Y.F.F. Soares 98 (2n* = 18); **C**, *P. melanospermum*, metaphase, *R.C. Oliveira & J.M. Mendoza 3025 (2n* = 20); **D**, *P. riparium*, diakinesis, *R.C. Oliveira & J.M. Mendoza 3008 (2n* = 40). — Scale bar 10 μ m.

LITERATURE CITED

- Bonasora, M.G., Pozzobon, M.T., Honfi, A.I. & Rua, G.H. 2015. Paspalum schesslii (Poaceae, Paspaleae), a new species from Mato Grosso (Brazil) with an unusual base chromosome number. Pl. Syst. Evol. 301: 325–2339. https://doi.org/10.1007/s00606-015-1231-0
- Burton, G.W. 1942. A cytological study of some species in the tribe Paniceae. *Amer. J. Bot.* 29: 355–359. https://doi. org/10.1002/j.1537-2197.1942.tb14228.x
- Dandin, S.B. & Chennaveeraiah, M.S. 1983. Chromosome number and meiotic behaviour in interpretation of basic chromosome number in the genus *Paspalum*. J. Cytol. Genet. 18: 26–33.
- Dandin, S.B. & Chennaveeraiah, M.S. 1988. Cytological evidence for apomixis in some species of *Paspalum L. J. Cytol. Genet*. 23: 61–67.
- Davidse, G. & Pohl, R.W. 1972. Chromosome numbers and notes on some Central American grasses. *Canad. J. Bot.* 50: 273–283. https://doi.org/10.1139/b72-038
- Davidse, G. & Pohl, R.W. 1974. Chromosome numbers, meiotic behavior, and notes on tropical American grasses (Gramineae). *Canad. J. Bot.* 52: 317–328. https://doi.org/10.1139/b74-042
- Gould, F.W. 1966. Chromosome numbers of some Mexican grasses. Canad. J. Bot. 44: 1683–1696. https://doi.org/10.1139/b70-241
- Honfi, A.I., Quarín, C.L. & Valls, J.F. 1990. Estudios cariológicos em gramíneas sudamericanas. *Darwiniana* 30: 87–94.
- Norrmann, G.A., Quarín, C.L. & Killeen, T.J. 1994. Chromosome numbers in Bolivian grasses (Gramineae). Ann. Missouri Bot. Gard. 81: 768–774.
- Peñaloza, A.P.S., Côrtes, A.L., Pozzobon, M.T., Santos, S., Rua, G.H. & Valls, J.F.M. 2008. Unusual chromosome numbers in *Paspalum* L. (Poaceae: Paniceae) from Brazil. *Genet. Molec. Res.* 7: 399–406. https://www.ncbi.nlm.nih.gov/pubmed/18551406
- Pohl, R.W. & Davidse, G. 1971. Chromosome numbers of Costa Rica grasses. Brittonia 23: 293–324. https://doi.org/10.2307/2805632
- Pozzobon, M.T. 2000. Contagens cromossômicas em espécies brasileiras de Paspalum L. (Gramineae). Acta Bot. Brasil. 14: 151–162. https://doi.org/10.1590/S0102-3306200000200003
- Quarín, C.L. 1992. The nature of apomixis and its origin in Panicoid grasses. *Apomixis Newslett.* 5: 8–15.
- Quarín, C.L. & Norrmann, G.A. 1990. Interspecific hybrids between five *Paspalum* species. *Bot. Gaz.* 151: 366–369. https:// doi.org/10.1086/337837
- Reeder, J.R. 1967. Notes on Mexican grasses VI. Miscellaneous chromosome numbers. *Bull. Torrey Bot. Club* 94: 1–17.
- Reeder, J.R. 1984. [Report]. In: A. Löve (ed.), Chromosome number reports LXXXII. *Taxon* 33: 132–133.
- Rua, G.H., Speranza, P.R., Vaio, M. & Arakaki, M. 2010. A phylogenetic analysis of the genus *Paspalum* (Poaceae) based on cpDNA and morphology. *Pl. Syst. Evol.* 288: 227–243. https:// doi.org/10.1007/s00606-010-0327-9
- Sánchez-Ken, J.G. 2010. Two new species of *Paspalum* (Paniceae: Panicoideae: Poaceae), a preliminary checklist of the genus in Mexico, and the identity of *P. crinitum. Revista Mex. Biodivers.*81: 629–647. http://www.scielo.org.mx/pdf/rmbiodiv/v81n3/v81n3a4.pdf
- Scataglini, M.A., Zuloaga, F.O., Giussani, L.M., Denham, S.S. & Morrone, O. 2013. Phylogeny of New World Paspalum (Poaceae, Panicoideae, Paspaleae) based on plastid and nuclear markers. *Pl. Syst. Evol.* 300: 1051–1070. https://doi.org/10.1007/s00606-013-0944-1

- Selva, S.B. 1976. Some preliminary cytological observations on a new basic number in *Paspalum convexum* (Gramineae). *Canad. J. Bot.* 54: 385–394. https://doi.org/10.1139/b76-035
- Sinha, R.R.P., Bhardwaj, A.K. & Singh, R.K. 1990. SOCGI plant chromosome number reports-IX. J. Cytol. Genet. 25: 140–143.
- Sousa, M.W.S., Pozzobon, M.T., Fagg, C.W., Valls, J.F.M. & Oliveira, R.C. 2017. Poaceae (*Mesosetum*). In: Marhold, K. & Kučera, J. (eds.), IAPT/IOPB chromosome data. *Taxon* 66: 1498, E35–E39. https://doi.org/10.12705/666.30

Saeedeh Sadat Mirzadeh Vaghefi* & Adel Jalili

Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), P.O. Box 13185-116, Tehran, Iran Author for correspondence: Mirzadeh@rifr-ac.ir

The study was financially supported by Research Institute Forests and Rangeland of Iran (grant no. 12-09-09-7701-89001). The authors are grateful to several collectors for material donation.

- * First chromosome count from Iran.
- ** First chromosome count for the species.

APIACEAE

** Ferulago angulata (Schltdl.) Boiss.

2*n* = 22, CHN. Iran, Azarbayjan, Uromieh, Marmisho valley, 37°35′05.28″N, 44°38′20.49″E, 1760 m, 2 Oct 2013, *A. Ashrafi 106062* (TARI) [Fig. 23A].

ASTERACEAE

Achillea talagonica Boiss.

2*n* = 18, CHN. Iran, Tehran, Lalon-Bazrab, 35°59'45.32"N, 51°34'31.27"E, 2550–2650 m, 4 Aug 2013, *J. Mohebi 106054* (TARI) [Fig. 23B].

Anthemis tinctoria L.

2n = 18, CHN. Iran, Mazandaran, Befor Gachsar, $36^{\circ}05'21.37''$ N, $51^{\circ}18'53.55''$ E, 2100 m, 10 Oct 2013, *A. Ashrafi 106056* (TARI) [Figs. 23C, 24A] (for other chromosome number report for this species from Iran see Chehregani & Mehanfar, 2008).

** Tanacetum abrotanifolium (L.) Druce

2*n* = 36, CHN. Iran, Azarbayjan, Marand, Zozodagh and Koohkamar, 38°45′00.96″N, 45°30′04.96″E, 2195 m, 20 Jul 2013, *J. Mohebi 106059* (TARI) [Fig. 23D].

Tanacetum pinnatum Boiss.

2*n* = 18, CHN. Iran, Tehran, Lalon-Bazrab, 35°59′45.32″N, 51°34′31.27″E, 2550–2650 m, 21 Sep 2012, *J. Mohebi* 106057 (TARI) [Fig. 23E].

For other chromosome number report for this species from Iran, see Chehregani & Mehanfar (2008).

** Varthemia persica DC.

2*n* = 16, CHN. Iran, Alborz, Chalus, 35°55'37.19"N, 51°03'37.53"E, 1605–2025 m, 20 Sep 2012, *J. Mohebi 106058* (TARI) [Fig. 23F].

CAPRIFOLIACEAE

***Centranthus longiflorus* Steven 2*n* = 32, CHN. Iran, Azarbayjan, Khoy, Ghator village, 38°28'10.46"N, 44°24'24.25"E, 1550 m, 5 Sep 2012, *J. Mohebi 106047* (TARI) [Figs. 23G, 24D].

CARYOPHYLLACEAE

** Gypsophila bicolor Grossh.

2*n* = 34, CHN. Iran, Kermanshah, Ghorveh to Songhor, 34°46′38.93″N, 47°26′28.37″E, 2220 m, 2 Oct 2012, *A. Ashrafi 106063* (TARI) [Figs. 23H, 24C].



Fig. 23. Somatic metaphases. **A**, *Ferulago angulata*, 2n = 22; **B**, *Achillea talagonica*, 2n = 18; **C**, Anthemis tinctoria, 2n = 18; **D**, *Tanacetum abrotanifolium*, 2n = 36; **E**, *Tanacetum pinnatum*, 2n = 18; **F**, *Varthemia* persica, 2n = 16; **G**, Centranthus longiflorus, 2n = 32; H, Gypsophila bicolor, 2n = 34; I, Hippophae rhamnoides, 2n = 24; **J**, Lallemantia canescens, 2n = 14; **K**, *Phlomis olivieri*, 2n = 20; L, Salvia hypoleuca, 2n = 22; **M**, *Teucrium hircanicum*, 2n =32; **N**, *Thymus persicus*, 2n = 30; **O**, *Sida rhombifolia*, 2n = 14; **P**, *Papaver bracteatum*, 2n = 14; **Q**, *Astragalus mesoleios*, 2n = 16; **R**, *Oreophysa triphylla*, 2n = 16.

- Scale bars =10 μm.

ELAEAGNACEAE

* Hippophae rhamnoides L.

2n = 24, CHN. Iran, Mazandaran, Pole zangole to Yoush, $36^{\circ}14'09.68''$ N, $51^{\circ}26'37.65''$ E, 3130 m, 21 Oct 2015, *S. Mirzadeh 105207* (TARI) [Fig. 231].

LAMIACEAE

** Lallemantia canescens Fisch. & C.A.Mey.

2*n* = 14, CHN. Iran, Gazvin, Taleghan, southern side of dam, 36°11'06.11"N, 50°37'40.44"E, 1860 m, 14 Aug 2013, *A. Ashrafi 106048* (TARI) [Fig. 23J, 24E].

* Phlomis olivieri Benth.

2*n* = 20, CHN. Iran, Fars, Shiraz to Kazeron, 29°38'17.09"N, 52°11'40.57"E, 2005 m, 20 Sep 2011, *J. Mohebi 106050* (TARI) [Fig. 23K].

Salvia hypoleuca Benth.

2*n* = 22, CHN. Iran, Tehran, Chalus, entrance of Kondor, 35°50'37.37"N, 51°05'45.58"E, 1666–2220 m, 25 Sep 2016, *S. Mirzadeh 106052* (TARI) [Fig. 23L].

** Teucrium hircanicum L.

2*n* = 32, CHN. Iran, Alborz, Chalus, Kiasar village, 36°02′42.40″N, 51°13′36.94″E, 2273 m, 20 Sep 2012, *J. Mohebi 106051* (TARI) [Fig. 23M].

** Thymus persicus (Ronniger ex Rech.f.) Jalas

2*n* = 30, CHN. Iran, Azarbayjan, Marand, Zozodagh and kamar mountain, 38°45′00.96″N, 45°30′04.96″E, 2105 m, 20 Jul 2013, *J. Mohebi 106049* (TARI) [Fig. 23N, 24F].

MALVACEAE

* Sida rhombifolia L.

2*n* = 14, CHN. Iran, Azarbayjan, Khoy, Ghator village, 38°28'10.46"N, 44°24'24.25"E, 1550 m, 23 Jul 2013, *J. Mohebi 106061* (TARI) [Fig. 23O, 24B].

PAPAVERACEAE

Papaver bracteatum Lindl.

2n = 14, CHN. Iran, Tehran, Siah bisheh, $36^{\circ}12'59.37''$ N, $51^{\circ}19'00.04''$ E, 2040 m, 20 Sep 2012, *J. Mohebi 106060* (TARI) [Fig. 23P].

For other chromosome number report from Iran, see Goldblatt (1974).

PAPILIONACEAE/FABACEAE

** Astragalus mesoleios Boiss. & Hohen.

2*n* = 16, CHN. Iran, Karaj to Chalus, Sirachal, 36°01'44.97"N, 51°18'47.56"E, 2000 m, 10 Oct 2013, *A. Ashrafi 106053* (TARI) [Fig. 23Q].

** Oreophysa triphylla (Bunge ex Boiss.) Bornm.

2*n* = 16, CHN. Iran, Tehran, Chalus, on the way to Kondor village, 35°50'37.37"N, 51°05'45.58"E, 2020 m, 25 Sep 2016, *S. Mirzadeh 106055* (TARI) [Fig. 23R].

LITERATURE CITED

Chehregani, A. & Mehanfar, N. 2008. New chromosome counts in the tribe Anthemideae (Astraceae) from Iran. *Cytologia* 73: 189–196. https://doi.org/10.1508/cytologia.73.189

Goldblatt, P. 1974. Biosystematic studies in *Papaver* section Oxytona. Ann. Missouri Bot. Gard. 61: 264–296. https://doi. org/10.2307/2395056



Fig. 24. Species in habitat: A, Anthemis tinctoria; B, Sida rhombifolia; C, Gypsophila bicolor; D, Centranthus longiflorus; E, Lallemantia canescens; F, Thymus persicus.